74-21 to 25

THE


ANATOMY

OF-THE

HUMAN BONES

AND

NERVES, &c.



ANATOMY

OF THE

HUMAN BONES and NERVES.

WITH

An Account of the Reciprocal Motions of the HEART,

AND

A Description of the Human Lac-

Corrected and enlarged in the SIXTH EDITION,

By ALEXANDER MONRO, Senior, M. D. and F. R. S. Fellow of the Royal College of Physicians, and Professor of Medicine and of Anatomy in the University of Edinburgh.

EDINBURGH:

Printed for G. HAMILTON, and J. BALFOUR. Sold by them and other Booksellers there.

At LONDON, by Meff. HITCH and HAWES, and A. MIL-LAR, J. NOURSE, R. BALDWIN, RIVINGTON and FLETCHER, and WILSON and DURHAM.

At DUBLIN, by J. SMITH.

MDCCLVIIL

A M.A.T.O.M.Y.

HUMAN DONES and NERVER



oooA oA

th.

tl.

te

de

Th.

A

W

mo

A Description of the Heman Lagran

tible .e.p. ... it is forth to the second annual second an

STUDENTS of ANATOMY

you cannot realizacly liame me, if Tlike

University of EDINBURGH.

GENTLEMEN,

WHEN this Ofteology was first printed in 1726, I did not know that Albinus, Winflow, and Palfyn, were to publish Descriptions of the Bones; otherwise my Papers probably would have remained yet undelivered to the Printers. I however flatter myself, that this Essay has been of Use to the Gentlemen who did me the Honour to attend my Lectures, by affisting them to understand my Sense and Representation of Things in this fundamental Part of Anatomy; and that it has bossibly been of more Advantage to them than a more compleat Work from an abler Hand, unless my Demonstrations had been in the Order and Method of fuch an Author.

This

This View of your Improvement, Gentlemen, is a prevailing Argument with me, to cause this Essay to be reprinted; and you cannot reasonably blame me, if I likewise acknowledge another Motive for it, which more particularly relates to myself. In a new Edition an Author has an Opportunity of making his Works more correct, compleat, and consequently acceptable to the Public, who may perhaps be indulgent enough to think this little Treatise not altogether useless; since more Reasoning on the Structure and morbid Phænomena of Bones is to be found in it, than in the other Writers, who have confined themselves almost entirely to the descriptive or proper anatomical Part of the Offeology.

I have here kept to the Plan of the former Editions, by first considering, in the Order that seemed to me most natural and methodical, every thing which I thought necessary to be known concerning Bones in general; and, in the second Part, I have described the several Bones composing the

Skeleton.

The Bones of Adults are what I principally endeavour to describe; but I have added added as much of the Osteogenea, as I think serviceable in the Practice of Phy-

fic and Surgery.

That little might be omitted of what was formerly done on this Subject, I have taken all the Affistance I could from Books; but have never afferted any anatomical Fact on their Authority, without confulting the Life, from which all the Descriptions are made; and therefore the Quotations from Such Books, serve only to do Justice to the Authors, who have remarked any Thing in the Structure of the Parts that was commonly omitted, and to initiate you in the History of Anatomy; which I once proposed to make complete, so far as related to this Subject; but not being able to procure Several Books, and being sensible bow many more may have never come to my Knowledge, I laid afide this Defign, of Purpose omitted many I could have inserted, and in some. Places I have changed an older Author for a later one, who has more fully or clearly described what I treated of. Beside Anatomists, I have also named several other Authors to confirm my Reasoning by pra-Etical Cases; of which it is not to be suppojed

posed my own Experience could furnish a

Sufficient Variety.

You'll readily observe, that I quote no Passages with a View to criticise or condemn them. This Precaution of giving no Offence, is very necessary in those who are sufficiently conscious of their being liable to lay themselves open to just Censure; and it prevents Occasions of useless Wrangling, in which generally both Parties are Losers, and the Public has little Advantage.

In this Treatise I always make use of the most common Name of each Part, and have put the synonymous Names to be met with in Books at the Foot of the Page, that the Reading might be smoother, and you might consult them at your Leisure to assist you in understanding different Authors.

in

b

be

The Descriptions and Reasoning are bere blended, without which I always find young Anatomists are soon disgusted with Authors: Their Imaginations cannot follow a long Chain of Descriptions, especially when they are not taught at the same time the Uses which the described Parts serve: Their Minds must have some Relaxation, by a Mixture of Reasoning, which never

never misses to strike the Fancy agreeably, and raises a strong Desire to understand

the Principles on which it depends.

The Phænomena of Difeases are all deduced in this Essay, from the Structure of the Parts, by way of Corollaries and Questions; which such an anatomical Work confined me to. And this Method has otherwise a good Effect: For, when one meets with an useful Proposition, and is obliged to employ a little Thought to find out its Solution, the Impression it makes is deeper, and be acquires a Fondness for it, as being in Part his own Discovery. My Pupils have frequently assured me, that they could, with very small Reflexion, trace out the whole Reasoning from which my Conclusions were drawn; I hope their Successors will also think this an agreeable Manner of being instructed.

Those Gentlemen who desired I would add the Lectures which I pronounce in my Colleges as a Commentary upon the Text, where the Diseases are mentioned, will, I persuade myself, excuse me for not complying with their Desire, when they consider the Design of this is to be a School-book, and how great the Touth in private, and pretending to inform the Public. Art. xxiv. Vol. v. of Medical Essays and Observations, published in this Place, is one of these Lectures which I gave as a Commentary on the Paragraph (p. 12.) concerning the different Kinds of Caries.

In this Edition, I have corrected the Mistakes and obscure Passages which I discovered in the former, and in some Places I have made the Descriptions more full and exact, aiming all I could to shun unnecessary Minuteness on the order: Whether I have bit that just Medium, is what you

and the Public must now judge.

I am still of opinion, that Figures of the Bones, would at any Rate have been unnecessary in a Book that is intended to be illustrated and explained by the Originals themselves; but would be much more so now, when my late ingenious Friend Mr. Cheselden, and Dr. Albinus, have published such elegant ones; and the Pictures in the Osteography published here, are disposed in the Order which I follow.

You

You have advantageous Opportunities in this Place, of studying all the Parts of Medicine, under the Professors of its different Branches in the University, and of seeing the Practice of Pharmacy, Surgery, and Physic, withour Surgeon-apothecaries, and in the Royal Infirmary, where the diseased Poor are carefully treated. These, your Interest, and, I hope, your Inclinations, will lead you, Gentlemen, so to improve, as that they may become the happy Means of your making a confiderable Figure in your several Stations. Whatever Assistance is in my Power towards such a defirable Event, shall be given with the greatest Pleasure by

Your humble Servant,

I

u

e

ibe ils

eied
in
if-

24

ALEX MONRO.

I the factor accounts greens Opportunities in the Loce, of havying airthe Ports of Medicine, under the Profefores of its differear Branches in the University, and of being the Pradice of Pharmedy, Surgery, and Phylic, will our Surgeon-apoth currer, and in the Royal Informery, where the discased Poor are carefully treated. These, your Interoft, and, I hope, your Inclinations, will had you, Gentlemen, fo to imprivate, and that they may become the hereby Means if you emailing a considerable Right in your feveral stations. Whatever Affic Same is in my 08 the SiSrds fich a definalle Event, facil be given with the greatest. Pleasure by

l our burble Servant,

Th

the que The different of

OHMOMENTAL!

ANATOMY

OF THE

HUMAN BONES.

PART I.

Of the Bones in general.

ONES are covered by a Membrane, named on that account Periosteum*, which is so necessary to them, that we must examine its Texture and Uses, before we can understand their Structure.

The Periosteum, as well as most other Membranes, can be divided into Layers of Fibres. The exterior ones, composed of the Fibres of the Muscles connected to the Bones, vary in their Number, Size and Direction, and consequently occasion a very great Difference in the Thickness and Strength of the Periosteum of different Bones, and even of the different Parts of the same Bone.—The internal Layer is every A where

[·] Membrana circumosfalis, omentum ossibus impositum.

where nearly of a fimilar Structure, and has its Fibres in the same Direction with those of the Bone to which they are contiguous. Ought not then the name *Periosteum* to be applied, strictly speaking, only to this internal Layer, to which the others are joined in an uncertain Manner and Number?

Some authors (a) endeavour to prove the internal Layer of Fibres of the Periosteum to be derived from the Dura Mater: For, fay they, fince the Membrane covering the Scull is plainly a Production or Continuation of the Dura Mater, which passes out between the Sutures; and fince there are Muscles on the Head, as well as in other Parts, which might furnish a Periosteum, it is needless to assign different Origins to Membranes which have the fame Texture and Uses. They add further, in proof of this Doctrine, that the Periosteum extends itself along the Ligaments of the Articulations from one Bone to another; and therefore is continued from its Origin over all the Bones of the Body.—While Anatomists were fond of the Hypothesis of all Membranes being derived from one or other of the two that cover the Brain, a Dispute of this kind might be thought of Consequence: But now that the Hypothesis is neglected as useless, it is needless to examine the Arguments for or against it.

Except where Muscles are inserted into the Periosteum, its external Surface is connected to the surrounding Parts by thin cellular Membranes, which can casily be stretched considerably, but shorten themselves whenever the stretching Force is removed. When these Mem-

branes

⁽a) Havers, Ofteolog. nov. Difc. 1. p. 16.

branes are cut off or broken, they collapse into such a small Space, that the Surface of the Pe-

riosteum seems smooth and equal.

When we attempt to tear off the Periosteum from Bones, we see a great Number of white Threads produced from the Membrane into them; and, after a successful Injection of the Arteries with a red Liquor, numerous Vessels are not only seen on the Periosteum (a), but most of the Fibres sent from the Membrane to the Bone, shew themselves to be Vessels entering it, with the injected Liquor in them; and when they are broken, by tearing off the Periosteum, the Surface of the Bone is almost covered with red Points.

The Veins corresponding to these Arteries are sometimes to be seen in Subjects that die with their Vessels full of Blood; though such numerous Ramissications of them cannot be demonstrated as of the Arteries, because sew of them have naturally coloured Liquors in them, and such Liquors can dissicultly be injected into them.

The great Sensibility of the Periosteum in the deep seated Species of Paronychia, in Exostoses, Nodi, Tophi, and Gummata, from a Lues venerea, or whenever this Membrane is in an instanced State, is a sufficient Proof that it is well provided with Nerves, though they are too small to be traced upon it; and therefore one cannot well determine, whether they are sent along with the Arteries in the common Way, or are

⁽a) Ruysch. Epist. 5. Tab. 5. Fig. 1.2. Epist. 8. Wab. 9. Fig. 1. 9.

derived from the tendinous Fibres of the Muf-

cles expanded on the Periosteum (a).

Vessels also pass through the Periosteum to the Marrow; of which more hereafter. And frequently Muscles pierce through the Periosteum, to be inferted into the Bones.

The chief Uses of the Periosteum are: 1. To allow the Muscles, when they contract or are stretched, to move and slide easily upon the Bones; the smooth Surface of this Membrane preventing any ill Effects of their Friction upon each other. 2. To keep in due Order, and to support the Vessels in their Passage to the Bones. 3. By being firmly braced on the Bones, to affift in fetting Limits to their Increase, and to check their Overgrowth. 4. To strengthen the Conjunction of the Bones with their Epiphyses, Ligaments and Cartilages, which are eafily separated in young Creatures, when this Membrane is taken away. 5. To afford convenient Origin and Infertion to feveral Mufcles which are fixed to this Membrane. And, lastly, to warn us when any Injury is offered to the Parts it covers; which, being infensible, might otherwise be destroyed without our Knowledge, or endeavouring to procure a Remedy.

When the cellular Substance connecting the Periosteum to the surrounding Parts is destroyed, these Parts are fixed to that Membrane, and lose the fliding Motion they had upon it; as we fee

daily

fe.

iel

to

luc

eac

uni

reg the

⁽a) See the Dispute about the Sensibility of this and of other Membranes in Zimmerman, Dissert de Irritabilit .-Act. Gotting. Vol. 2. Haller. Sur la nature sensible et irritable. - Whytt's physiolog. Essays II. - Reimar. Dinert. de Fungo Articulor. § 26. 34.

This

daily in Issues, or any other tedious Suppurations near a Bone.—When the Vessels which go from the Periosteum to the Bones, are broken or eroded, a Collection of Liquor is made between them, which produces a fordid Ulcer or rotten Bone. This often is the Case after Fractures of Bones, and Inslammations of the Periosteum, or after Small-pox, Measles, Spotted Fevers, and Erysipelas.—Do not the Disorders of the Periosteum, coming along with or soon after the cutaneous Diseases, indicate a Similarity of Structure in the Periosteum and Skin?

The Bones are the most hard and solid Parts of the Body, and, as all other Parts where large Vessels do not enter, are generally of a white Colour; only in a living Creature they are bluish, which is owing to the Blood in the small Vessels under their Surface. The less therefore and fewer the Vessels are, and the thicker and firmer the bony Surface covering the Vessels is, the Bones are whiter. Hence the Bones of Adults are whiter than those of Children; and, in both young and old, the white Colour of different Bones, or of the several Parts of the same Bone, is always in Proportion to their Ve'fels and Solidities; which Circumstances ought to be regarded by Surgeons, when they are to judge of the Condition of Bones laid bare.

Bones are composed of a great many Plates *, each of which is made up of Fibres or Strings united by smaller Fibrils (a); which being irregularly disposed, and interwoven with the other larger Fibres, make a reticular Work.—

e

i,

:0

y

of

m-

^{*} Squamæ, Bractez, Laminæ.

This Texture is plainly seen in the Bones of Fœtuses, which have not their Parts closely compacted, and in the Bones of Adults which have been burnt, long exposed to the Weather, or whose Composition has been made loose by Diseases.—The Chinks which are generally made according to the Direction of the larger Fibres of Bones that have undergone the Action of Fire, or of the Weather, shew the greater Strength of these than of the Fibres which connect them.—Numerous accurate Observations of the different Times in which Exsoliations are made from the Sides or Ends of similar Bones, might bid fair to determine what is the proportional Force of Cohesion in the two sorts of Fibres.

The Plates are faid (a) to be firmly joined to each other by a great Number of Claviculi, or small bony Processes, which, rising from the inner Plates, pierce through some, and are fixed into the more external ones. Of these Nails, sour Kinds, viz. the perpendicular, oblique, beaded, and crooked, have been described: But in Bones sitly prepared, I could only see numerous irregular Processes rising out from the Plates (b).

of

are

fer

Di

are

hui

Spa

celli

diffe

and diffe Som

Though the exterior Part of Bones is composed of firm compact Plates, yet they are all more or less cavernous internally. In some (e.g. middle thin Part of the Scapula and Os Iliúm) the solid Sides are brought so near, that little Cavity can be seen; and in others (middle of Os Humeri, Femoris, &c.) the Cavities are so large, that such Bones are generally esteemed

⁽a) Gagliard. Anat. offium nov. invent. illustrat. cap. 1.

⁽b) Malpigh. oper. posthum.

to be hollow or fistular. But the internal spongy Texture is evident in young Animals; and some of it may be seen to remain in those of greatest Age, when Bones are cautiously opened, after they have been kept so long as to be free of the Oil they contain, or after being burnt.

This spongy cavernous internal Part of Bones, is generally called their Cancelli, or LATTICEwork, and is formed in the following Manner. The Plates are firmly joined about the Middle of the Bone; but as they are extended towards its ends, the more internal Plates separate from the exterior, and stretch out their Fibres towards the Axis of the Bone, where they are interwoven with the Fibres of other Plates that have been fent off in the same Way. Seeing the Plates are thus constantly going off, the folid Sides of the Bones must become thinner, and the Lattice-work must be thicker and stronger, towards their Ends. This is evident in many of them, where the folid Sides of their Middle are very thick, and the Cancelli are scarce obfervable; whereas, at the Ends, where their Diameter is greatest, the folid Walls or Sides are not thicker than Paper, and the Cancelli are numerous and large enough to fill up the whole Space left between the Sides.

Ō

r

e

d

1

n

J.

n-

Ill

g.

m) tle

of fo

red

to

. 1.

The Twisting and Windings which these Cancelli make, and the Interstices which they leave, differ considerably in Figure, Number and Size, and therefore form little Cells, which are as different, but communicate with each other. Some Writers (a) minutely remark these dif-

ferent

⁽a) Gagliard. Anat. offium, cap. 1. Obf. 4. 5. 6. 7.

ferent Appearances of the Cancelli, after they begin to separate from the Plates; and from thence distinguish them into wrinkled, perforated, and Net-like.

The Cancelli sustain the membranous Bags of the Marrow which are stretched upon them, and thereby hinder these membranous Parts to be torn or removed out of their proper Places, in the violent Motions and different Postures which the Bones are employed in. This Support which the Cancelli afford the Marrow, also saves its Membranes and Vessels, in the lower parts of the Bones, from being compressed by the Weight of the Marrow above.

The Depressions between the Fibres of the external Plates of Bones appear like so many Furrows on their Surface, into each of which the Periosteum enters; by which the Surface of Contact, consequently the Cohesion, between it and the Bone, is considerably increased, and a greater Number of Vessels is sent from it into

the Bone, than if it was a plain Surface.

Both on the Ridges and Furrows, numerous little Pits or Orifices of Canals are to be seen, by which the Vessels pass to and from the Bones.

After a successful Injection, the Arteries can be traced in their Course, from the Pits to the Plates and Fibres; and, in sawing, cutting or rasping the Bones of living Creatures, these Vessels discover themselves, by the small Drops of Blood which then ouze out from the most solid Part of the Bones. But the clearest Demonstration of the intimate Distribution of these small Arteries, is, to observe the Effect of such a tinging Substance as can retain its Colour, when swallowed,

Art. 1739.

ŀ

t

t

0

to

qu

n pa

at

me

Pla

the

Ar

is e

lefs

2.

of

ject

Fro

diffi

swallowed, digested and mixed with the Blood of any living Animal, and at the fame time has Particles small enough to be conveyed into the Vessels of the Bones; such is Rubia tinctorum, Madder-root (a): For we see the gradual Advances which this Tincture makes from the Periosteum into the more internal Parts of the Bones, and how univerfally the Distribution of the Liquors is made, the whole bony Substance being tinged. Whether the Time in which this tinging Liquor passes from the outer to the internal Plates, till all the Plates are made of its Colour, and the Time which the disappearing of the Dye, after giving the Creature no more of this fort of Food, makes us think it takes to return, are the same in which the natural Liquors circulate, is uncertain; because this tinging Substance may move more slowly, or may pals more quickly, than the natural Liquors do. -The Arteries are larger near each End than at the Middle of the large Bones that are much moved; because they not only serve the bony Plates near the Ends, but pass through them to the Marrow.—As Animals advance in Age, the Arteries of the Bones become less capacious; as is evident, 1. From the Bones of Adults having less Blood in them than those of Children have. 2. From their becoming incapable in old Age of admitting the coloured Powders used in Injections, which cafily pass in Youth. And, 3. From the Bones of old Creatures being more difficultly tinged with Madder than those of

f

1

d

•

5

,

5.

n

C

r.

1-

of

d

2.

3-

d,

⁽a) Philosoph. Transact. Num. 442. Art. 8. Num. 443. Art. 4. Num. 457. Art. 4.— Mem. de l'Acad. des Sciences, 1739, 1742.

young ones.—If Authors have not mistaken, the Arteries of Bones have fometimes become very

n

fo

ri

N

F

ot

01

an

W

thi

ly

Sei

cau

Bor

Cir

and

al o

bres

Bod

the

Ren

in ac

the !

demo

incre

Matt

till o

-Tl

large (a).

Since the Veins of the Periosteum are difficultly filled with a coloured Injection, we are not to expect that their Branches, the Veins of the Bones, can be shewn; for they are too small to be feen without fuch Affistance: But we may conclude from Arteries being accompanied with Veins so far as we can trace them in every other Part of the Body, that there are also Veins in the Bones; and the disappearing of the Tin-Eture of Madder, after Bones of living Animals are coloured with it, could not be without fuch

Veins to carry it away.

The Bones of a living Animal are so insensible, that they can be cut, rasped, or burnt, without putting the Creature to Pain, and the Nerves distributed in their Substance cannot be shewn by Diffection; from which it might be inferred that they have no Nerves distributed to them: But the general Tenor of Nature, which bestows Nerves to all the other Parts, should prevent our drawing fuch a Conclusion. And if Sensibility is a fure Proof of Nerves entering into the Composition of any Part, as it is generally allowed to be, we have sufficient Evidence of Nerves here in the Bones; for the granulated red Flesh which sprouts out from them, after an Amputation of a Limb, or performing the 0peration of the Trepan, or after an Exfoliation, is exquisitely sensible: And, in some Ul-the co cers of Bones, where the Periosteum was all se. become parated,

⁽a) Diemerbroek. Anat. lib. 9. cap. 1 .- Mery. Hift. de l'Acad. des Sciences, 1704.

parated, the Patient suffered racking Pain, if the Bone was touched with a rough Instrument; nor was he free of Pain after the Bone was perforated (a).—The reason why the Nerves of rigid hard Bones become insensible, is, That all Nerves must have a considerable Degree of Flexibility at the Part where Objects are applied, otherwise it cannot be affected by their Impressions. We see this illustrated in a very common analogous Case, the Growth of a new Nail: When the former one has suppurated off, the thin Membrane which first appears, is exquisitely sensible; but gradually becomes dull in its Sensation, till it can be cut or scraped, without causing Pain, after it is formed into a hard Nail.

t

e

0

y

h

).

1

ls

h

e,

ut

1-

y

at

ut

WS

nt

oi-

he

al-

of

ted

an

0.

ati. Ul. fe.

ed,

. de

From what has been faid of the Vessels of Bones, it is evident, that there is a constant Circulation of Fluids in every Part of them; and that there is a perpetual Waste and Renewal of the Particles which compose the folid Fibres of Bones, as well as of other Parts of the Body; the Addition from the Fluids exceeding the Waste during the Growth of the Bones; the Renewal and Waste keeping pretty near par in adult middle Age; and the Waste exceeding the Supply from the Liquors in old Age; as is demonstrable from their Weight: For each Bone increases in Weight, as a Person approaches to Maturity; continues of nearly the same Weight till old Age begins, and then becomes lighter. The specifick Gravity of the solid Sides, on the contrary, increases by Age; for then they become more hard, compact and dense. In Confequence

⁽a) Nicol Massa lib. introd. Anat. cap. 30.

fequence of this, the Bones of old People are thinner and firmer in their Sides, and have larg-

er Cavities than those of your Persons.

The vafcular Texture of Bones must make them subject to Obstructions, Ecchymoses, Ulcers, Gangrenes, and most other Diseases which the softer Parts are affected with; and therefore there may be a greater Variety of Caries than is commonly described (a.)

Hence we can account for the following Ap-

pearances.

Hamorrhagies from fungous Flesh rising out from the most solid Part of a cut Bone (b.)

(

f

t

K

tl

fā

PO

na th

OV

fer the

IM

Vei con

ton

11 19

tran

feril alfo whic

1

The regular alternate Elevation and Subfiding or apparent Pulsation frequently to be seen in some of the Cells of a carious Bone.

Cells resembling Cancelli, sometimes seen in the Part of a Bone, which, in a natural State,

is the most solid and firm (c.)

A Bone as a Tube including another Bone

within it (d.)

On the internal Surface of the folid Parts of Bones, there are Orifices of Canals, which pass outwards through the Plates to open into other Canals that are in a longitudinal Direction, from which other transverse Passages go out to terminate in other longitudinal Canals; and this Structure is continued through the whole Substance of Bones, both these Kinds of Canals becoming similar gradually as they approach the outer Surface.

(b) Medical Essays, Vol. IV. Art. 21.

(d) Idem, ibid.

⁽a) Edinburgh Medical Essays and Obs. Vol. V. Art. 25.

⁽c) Ruysch. Thes. 8. Num. 8. Thes. 10. Num. 176.

Surface (a.) - These Canals are to be seen to the best Advantage in a Bone burnt till it is white: When it is broken transversely, the Orifices of the longitudinal Canals are in View; and when we separate the Plates, the transports ones are to be observed. Here however we are in Danger of believing both these Sorts of Canals more numerous than they really are; because the Holes made by the Processes connecting the Plates of Bones have the Appearance of the transverse (b), and the Pallages for the Blood-Veffels rejemble the longitudinal Canals. I don't know how we are to keep free of Error about the transverse Canals; but think we may distinguish between the two Kinds of longitudinal ones; for the Passages of the Vessels are largest near the external Surface of the Bone, and every transverse Section of them is circular; whereas the longitudinat Canals are largest near the Cancelli, and their transverse Sections appear to me of a flat oval Figure, which may be owing to the disferent Momentum of the Fluids conveyed in them. The Situation of the larger longitudimal Canals, and of the Passages of the larger Veffels, makes a Bone appear more dense and compact in the Middle of its folid Sides, than towards its outer and inner Surfaces, where it is spongy.

ıt

in

in

e,

ne

of

ass

her

om

mi-

tru-

nce

ing

uter

ace.

25.

We see Marrow contained in the larger transverse and longitudinal Canals just now described, and from thence judge that it passes also into the smaller ones. The Drops of Oil which we discover with a Microscope every where

⁽a) Havers, Ofteolog. nov. p. 43.

⁽b) Morgagn. Advers. 2. Animad. 25.

where on the Surface of a recent Bone fractured transversly, and the outing of Oil through the most folid Bones of a Skeleton, which renders them greafy and yellow, are a Confirmation of the Use of these Canals. Of what Advantage this Distribution of the Marrow. through the Substance of Bones is, will be mentioned when the Nature and Use of this animal Oil is inquired into.

Most Bones have one or more large oblique Canals formed through their Sides for the Paffage of the medullary Veslels, which are to

be described afterwards.

Bones exposed to a strong Fire in Chemical Vessels, are resolved, in the same Manner as the other Parts of Animals, into Phlegm, Spirit, volatile Salt, fetid Oil, and a black Caput mornum. But the Proportion of these Principles varies according to the Age, Solidities, and other Circumstances of Bones. Young Bones yield the largest Proportion of Phlegm; spongy Bones afford most Oil, and solid ones give most Salt and black Residuum. - Though this Residuum can scarce be changed by the Force of Fire while it is in close Vessels; yet, when it is burnt in an open Fire, the tenacious Oil, to which it owes its black Colour, is forced away, and a white Earth is left that has little or no fixed Salt in it; on which Account these Albes are the most proper Materials for forming into Vessels that are to undergo the greatest Force of Fire, being scarce capable of vitrifying by Fire, as most other earthy Substances do. This Earth seems to be the proper constituent solid Part of Bones, and

the other Principles give it Firmnels and Tenacity: For the Quantity of the Earth is fogreat, that, after all the other Principles are separated from a Bone, its former Shape and Size remain (a); but it is very brittle till it is moistened with Water or Oil, when it recovers some Tenacity. - The Increase of the Proportion of Earth in old People's Bones is one Reason of their being more brittle than those of young People are.

Lest any imagine the Salts and Oils of Bones, while in a natural State, to be of the lame acrid Kind with those obtained from them by the Chemical Analysis, it is to be observed, that these Principles may be extracted from Bones in the Form of a very mild Jelly, by

boiling them in Water.

The Bones fultain and defend the other

Parts of the Body.

ıl.

13

1ut

1-

S,

ıg.

1;

es

gh

he

et,

C1-

18

1at

Ac-

te-

rgo

pa-

thy

the

and

the

Bones are lined within, as well as covered externally with a Membrane; which is therefore commonly called Periosteum Internum.

The internal Periosteum is an extremely fine Membrane; nay frequently it has a loofe reticular Texture; and therefore it is compared by fome to the Arachnoid Coat of the spinal Mar. row: So that we cannot expect to divide it in, to Layers, as we can divide the external Perio. seum. We can however observe its Processes. entering into the transverse Pores of the Bones, where probably they are continued to form the immediate Canals for the Marrow distributed thro the Substance of the Bones; and along with them Vessels may be sent as from the external tra sou haven is a mil 20 meted a Pario.

⁽a) Havers, Ofteolog. nov. Difc. I. p. 31.

Perioseum, into the Bone (a). These Processes being of a very delicate Texture, the Adhesion of this Membrane to the Bone is so small, that it separates commonly more easily from the Bone than from the Marrow which it contains: Wherefore, one might call it the common Membrane of the Marrow, rather than by the Name it now has. But whether one or tother Designation ought to be given it, is not

worthy a Dispute.

From the internal Surface of the internal Periofteum, a great Number of thin Membranes are produced; which, passing a-cross the Cavity, unite with others of the same Kind, and form so many distinct Bags, which communicate with each other; and thele again are subdivided into communicating yeficular Cells, in which the Marrow is contained. Hence it is, that the Mairow, when hardened, and viewed with a Migroscope, appears like a Cluster of small Pearl; and that the hardened Marrow of Bones buried long under Ground, or laid fometime in Water, and then dried, is granulous (!). This Texture is much the same with what obtains in the other cellular Parts of the Body, where Fat is collected; only that the Cells containing the Marrow are smaller than those of the Times adiposa or cellulosa else. where, which probably is owing to their being inclosed in the Bones, where they are not to much stretched or extended as in other Parts.

in

ni

F

fe]

fre

ne

Ch

Ch

thi

wh

theic

The Markow is the oily Part of the Blood, feparated by small Arteries, and deposited in

⁽a) Winflow, Exposition Anat. des Os frais, § 82. 83.
(b) Ruysch. Thesaur. 9. Num. 2. & Advers. Dec. 111.
Obs. 9.

these Cells. Its Colour and Consistence may therefore vary according to the State of the Vessels, and their Distribution on the Membranes of the Cells.

Besides the Arteries, which I mentioned already, p. 9. to be sent from the Bones to the Marrow, there is at least one Artery for each Bone; several Bones have more, whose principal Use is to convey and secent this oily Matter. After these Arteries have pierced the solid Side of a Bone, they are divided into selection where on the internal Periosteum, and afterwards spread their Branches on the medulary Cells.

The Blood, which remains after the Secretion of the Marrow, is returned by proper Veins, which are collected from the Membranes into one or two large Trunks, to pais out at the same Holes or Passages at which the

Artery or Arteries enter.

h

e

n e-

s.

d,

in,

1.

The general Rule of the small Vessels decreasing in their Capacities as Animals advance in Age, to which many Phænomena in the Animal Occonomy are owing, obtains here: For though the Trunks of the medullary Vessels enlarge as Animals turn older; yet the small Branches become smaller; as is evident from Injections, which cannot be made to pass near so far in these Vessels of Adults as of Children. Hence the Marrow is bloody in Children, oily and balmy in middle Age, and thin and watery in old People.

By Experiments made on the Marrow, when Bones of living Animals are opened or

B 3

Cut

cut through (a), and from the racking Pain with which Suppurations within Bones are frequently attended, we have sufficient Proof, that the Membranes here are sensible, and consequently have Nerves distributed to them. Hippocrates (b), might therefore say justly, that a Wound penetrating into the Cavity of a Bone may produce a Delirium.

The Vessels of the Marrow, wrapt up in one common Coat from the Periosteum, pass through the Bones by proper Canals; the most considerable of which are about the Middle of each Bone, and are very oblique. Sometimes these Vessels continue at a little Distance in their Passage, when the Canal is divided by a small

bony Partition or two as daily boold ad I

From the Structure of the Contents of the Bones, we may judge how these Parts, as well as others, may be subject to Oidema, Phlegmon, Erysipelas, Schirrhus, &c. and may thence be led to a Cure of each, before the common Consequence, Putrefaction, takes Place, and frequently occasions the Loss of the Limb, if not of the Patient.

I

a

n

b

n

m

is

fu

th

fif

of

cel

qu

toi

Hay

The Marrow is of very considerable Use to the Bones; for, by entering their transverse Canals, and passing from them into the longitudinal ones, it is communicated to all the Plates, to soften and connect their Fibres, whereby they are preserved from becoming too brittle; as we see they do in burnt Bones, or those long exposed to the Air; in People labouring under old Age, Pox or Scurvy: In all which

(5) Aphoritin, § 7. Aph. 24.

⁽a) Du Verney, Memoirs de l'Acad. des Sciences, 1700.

which Cases, the Oil is either in too little Quantity, or has its natural good Qualities changed for worse jones. some over some bas

Belides this Advantage which the Substance of Bones has from the Marrow, their Articulations are said (a) to receive no less Benefit from it: For it is thought that the Marrow passes into the articular Cavities, through the Holes which are in the Bones near the large Joints. And, as a Proof of this, it is alledged, that Butchers, upon seeing the greater or less fer Quantity of Marrow in the Bones of Cows, can tell whether they have travelled far or little before they were saughtered.

1

e;

1

1

5-

y

le

e,

b,

to

11

he es,

00

ov la+

all

ch

00.

When the Marrow, after having served the Uses mentioned, is reasumed into the Mass of Blood (as it is continually, in common with all other fecreted Liquors that have not Paffages formed for conveying them out of the Body) it corrects the too great Acrimony communicated to the faline Particles of our Fluids. by their Circulation and Heat; in the fame manner as lixivial Salts are blanted by Oil in making Soap. Hence, in acute Difeases, the Marrow, as well as the other Fat of the Body, is quickly wasted, but must be immediately supplied by Liquors from the Vessels, seeing the Cells within the Bones, which have no Alfiftance to their Contraction from the Pressure of the Atmosphere, cannot collapse, as the Tela cellularis under the Skin does, when the Liquor in its Cells is abforbed, the Bones therefore are always full.

Since

⁽a) Joan. de Muralto Vade mecum, Anat. Exercit. V. § 3. Havers, Ofteolog. nov. Difc. 3. p. 179.

Since it is the Nature of all Oil to become thin and rancid when exposed long to Heat, and Bones have much Oil in their firm hard Substance, we may know why an ungrateful Smell, and dark coloured thin Ichor, proceed more from corrupted Bones, than from other Parts of the Body; and we can understand the Reason of the Changes of Colour which Bones undergo, according to their different Degrees of Mortification. Hence likewife we may learn the Caufe of a Spina ventofa, and of the Difficulty of curing all Caries of Bones proceeding from an Obstruction, and consequent Putrefaction of the Marrow; and of the quick Palfe, Thirst, and bettic Paroxysms, to often attending these Diseases. These Phanomena alfo teach us the Reason of the fatal Prognosis taken from black fetid Urine in Fevers.

Though Bones fo far agree in their Structure and annexed Parts, yet we may observe a considerable Difference among them in their Magnitude, Figure, Situation, Substance, Connexion, Wies, &c. From which Authors have taken Occasion to diffinguish them into as many Classes as they could enumerate of these different Circumstances. But these being obvious to every Person that looks on Bones, I shall only mention one of them; which comprehends very near the whole Bones of the Body, and at the same time leads us to examine the most considerable Variety that is to be found in the Disposition of their constituent Parts, and in their Uses. It is this, that some Bones are broad and flat, while others are long and

round.

The

U

0

111

th

no

to To

yei the

hav

and

ries

exti

too

geo Mid

to in

es t Stre

then

estin

cylin

(4)

The broad Bones have thin Sides, by the Plates being loop and equally fent off to form the Lattice-work; which therefore is thicker, and nearly of an equal Form all through. By this Structure, they are well adapted to their Uses, of affording a large enough Surface for the Muscles to rife from, and move upon, and of defending sufficiently the Parts which they inclose.

The round Bones have thick strong Walls in the Middle, and become very thin towards their Ends, which is owing to year few Platen Sepal rating at their Middle, where, on that account, the Concelli are so fine and small, that they are not taken notice of : But such Bones are faid to have a large Reservoir of Oil in this Place! Towards their Ends the Lattice work becomes very thick, and rather more compleat than in the other fort of Bones.—Their round Bones having strong Forces naturally applied to them) and being otherwise exposed to violent Injul ries, have need of a cylindrical Figure to refift external Pressure, and of a considerable Quantity of Oil to preferve them from becoming too brittle. Belides which, they ware advantageously provided with thick Sides towards their Middle, where the greatest Forces are applied to injure them; while their Hollowness increase es their Diameter, and consequently their Strength, to resist Forces applied to break them transversely (a) a Thus, for Instance, in estimating the proportional Resistance of two ylindrical Bones of unequal Diameters, but gaithines and in whitever Manner their his

are dilco.ed, their Strength muit aim up h

t

5

2

1

y

Ú.

is Il

2.3

7,

0

d

S,

23

id

16

⁽a) Galilei. Mechanic. Dialog. 1.

consisting of an equal Number of similar Fibres uniformly disposed round each, it is plain,

Bones is equal, because they confist of equal

Numbers of fimilar Fibres.

10 2. That the absolute Forces of all the Fibres in each Bone have the same Effect in resisting any Power applied to break them, as if the Sum of all their Forces was united in the respective Centers of the transverse Sections where the Fractures are to be made. For, by Hypothesis, the Fibres being uniformly disposed in each, there is not any Fibre in either Bone that has not a corresponding Fibre; the Sum of both whose Distances from the Axis of Revolation (about which all the Parts of the Bone must revolve in breaking) is equal to two Semidiameters of the Bone: Confequently each Fit bre, and all the Hibres, may be regarded as refifting at the Distance of one Semidiameter of Radius from this Axis, that is, in the Center.

f

m

h

ol

fin

et.

w

Ol

ag

M

uni

Co

cel

the

one

stan

or 1

(a erfiel

(b)

resius,

ductio,

cesso

3. Since the united Force of all the Fibres is to be regarded as refishing at a Distance from the Center of Motion equal to the Semidiameter, it follows, that the total Resistance of all these Fibres, or the Strength of the Bone is proportional to its Semidiameter, and conse-

quently to its' Diameter! of the mode of int

I have here taken for an Example one of the most simple Cases for calculating the proportional Forces of Bones. But, was it not too so reign to the present Design, it might be universally demonstrated, that, of whatever Figure Bones are, and in whatever Manner their Fibres are disposed, their Strength must always be in a Ratio.

Ratio, compounded of the Area of their transverse Sections, or of their Quantity of bony Matter, and of the Distance of the Center of Gravity of these Sections from the Center of Motion or Fulcrum, on which the Bone is supposed to be broken (a).

0

e

e-

ns

DY

ed

ne of

0-

ne ni-

Fil

re-

10

is

om ne-

all

is ife-

the

fo.

ret.

ure

ores in a

Since therefore the Strength of Bones depends on their Number of Fibres, or Quantity of Matter, and the Largeness of their Diameters, one may conclude, that the Part of a Bone formerly fractured, and reunited by a Callus, must be stronger than it was before the Fracture happened; because both these Advantages are obtained by a Callus: Which is a wife Provision, fince Bones are never fet in fuch a good Direction as they were naturally of; and then where-ever a Callus is formed, there is such an Obstruction of the Vessels, that if the Bone was again broken in the same Place, the offifick Matter could not so easily be conveyed to reunite it. This Callus may indeed; for Want of Compression, be allowed to form into a spongy cellular Substance (b); but even in this Case the Strength of the Bone is here increased by one or both the Causes above mentioned.

Many Bones have Protuberances, or Processes, rising out from them. If a Process stands out in a roundish Ball, it is called Caput, or Head.—If the Head is flatted, it obtains the

Appellation

⁽a) See the Demonstration of this Theorem by Dr. Porersield in the Edinburgh Medical Essays, Vol. 1. Art. 10.

⁽b) Ruysch. Thesaur. 8. n. 49. Mus. Anat. Thec. B. Re-

posit. 2. n. 2.

* 'Απόρυσες, ἔκουσες, ἔξοχη, προβολη, πρόβλημα, Εχcessus, Explanatio, Tuberculum, Gibbus, Eminentia, Productio, Extuberantia, Projectura, Enascentia.

Appellation of Condyle.—A rough unequal Protuberance, is called Tuberosity.—When a Process rises narrow, and then becomes large, the narrow or small Part is called Cervix, or Neck.—Long Ridges of Bones, are called Spines.—Such Processes as terminate in a sharp Point, have the general Name of Coronæ*, or Coronoid, bestowed on them, tho' most of them receive particular Names from the Resemblance they have, or are imagined to have, to other Substances, e. g. Massoid, Styloid, Anchoroid, Coracoid, Spinal, &c.—Such Processes as form Brims of Cavities, are called Supercilia+

Processes serve for the advantageous Origin and Insertion of Muscles, and render the Arti-

t

tl

T

in

th

To

th

pe

lete

the

En

mo ann

culations firm and stable. et author s 1979-9

Before leaving this Subject, we must remark, that much the greater Number of what are called Proceses in adult Bones, discover themfelves in Children to be Epiphyses, or distinct Bones, which are afterwards united to the other Parts; such are the Styloid Processes of the remporal Bones, Processes of the Vertebra, Trochanters of the Thigh, &c. However, as I delign to infift chieffy of the Description of the adult Skeleton, in which the Union of these Parts is so intimate, that scarce any Vestige remains of their former Separation, - I shall retain the common Appellation of Apophyse, or Process, to all such Protuberances; but shall remark the principal ones that have cler to a ap. 1 m. Anat 11. c lt Ros-

Roffra, Glandes.

[†] Trues, οφρυες, αμβονες, χέλη, Labra.

no just Title to this Name, when they occur

in the Description of particular Bones.

1

a

1

d

à *,

e-

e,

n-

es

cl-

in

ti-

k,

al-

m-

nct

ner

the ræ,

as of

1 of

Ve-

, - I

Apo-

ces;

ave

no

On the Surfaces of a great many of the Bones, there are Cavities, or Depressions: If these are deep, with large Brims, Authors name them Cotylæ*. — If they are superficial, they obtain the Defignation of Glenæ, or Glenoid. These general Classes are again divided into several Species: - Of which Pits are small roundish Channels funk perpendicularly into the Bone. - Furrows, are long narrow Canals, formed in the furface; - Nitches, or Notches, small Breaches in the Bone; Simuosities, broad, but superficial Depressions without Brims; — $Foff\alpha$, large deep Cavities, which are not equally furrounded by high brims;—Simuses, large Cavities within the Substance of the Bones, with small Apertures; - Foramina, or Holes, Canals that pierce quite through the Substance of the Bones. - When this last fort of Cavity is extended any long Way within a Bone, the middle Part retains the Name of Canal, and its Ends are called Holes.

The Cavities allow the Heads of Bones to play in them; they lodge and defend other Parts; they afford fafe Passage to Vessels, Muscles, &c. To mention more would engage us too much in the History of particular Bones, which more properly belongs to the demonstration of the Skeleton, where we shall have Occasion to observe

these several Species of Cavities.

To far the greater Number of Bones, whose Ends are not joined to other Bones by an immoveable Articulation, there are smaller ones annexed, which afterwards become scarce distinguishable

Κοτυλίδες, όξυβάροι, Acetabula, Pixides, Buccellæ.

some Bones have one, others have two, three or four of these Appendices annexed by the Means of Cartilages, which are of a considerable Thickness in Children, but by Age become thinner; the Ossisication proceeding from the End of the Bone on one Side, and from the Epiphyses on the other, till at last, in Adults, the Place of their Conjunction can scarcely be seen on the external Surface; and it is only sometimes that we can then see any Mark of Distinction in the Cancelli (a).

Several Processes (e. g. Trochanters of the Thigh, Spine of the Scapula, &c.) have Epiphyses; and Processes frequently rise out from Epiphyses; for Example, at the lower End of

the Femur, Ulna, Tibia, &c. (b).

The Epiphyses are united chiefly to such Bones as are destin'd for frequent and violent Motion; and for this Purpose they are wisely framed of a larger Diameter than the Bone they belong to; for by this Means, the Surface of Contact between the two Bones of any Articulation being increased, their Conjunction becomes firmer, and the Muscles inserted into them act with greater Force, by reason of their Axes being surther removed from the Center of Motion. These Advantages might indeed have been obtained by the Expansion of the End of the Bone itself, to a Thickness equal to

1

of

th

* Applantatio, Additamentum, Adnascentia, Adnexum, Perone.

(b) Vefal. De Human. Corp. fabrica, lib. 1. cap. 3.

⁽a) Winstow, Exposition Anatomique de Corps Humain Traite des Os secs, § 116.

that of the Epiphyses; but then the constant Separation of new Plates to form so wide a cellular Structure, must have left the solid Sides of the Bones so thin, as to yield easily, either to the Action of the Muscles fixed to them, and passing over them, to the Weight several of them are obliged to support, or to the Appli-

cation of any other external Force.

n

h

ıt

y

1e

ce

r-

110

to

ar

er

ed

he

to

at

um,

ain

Several Anatomists (a) thought that the Epiphyfer ferve other Purposes; such as, securing the Ligaments of the Articulations which rife out from between the Bones and them; for, as foon as these Parts are intimately joined, the Ligainents infinuated betwixt them must have a much stronger Connexion than they could have to the smooth Surface of the Bones. Such an Interception of the Ligament between the Body of the Bone and its Epiphyfe is not to be feen, but as at this Place, the Bone remains longer fost than any where else, and the Adhesion of the Periosteum, and of Ligaments to Bones, is always stronger in Proportion to the Bones being nearest to the Consistence of their Parts, which is being foftest, their Opinion of the stronger Connexion of the Ligaments, where the Bones and Epiphyses join, is not without Foundation.

Possibly too, by the Fibres of Epiphyses not extending themselves so longitudinally as those of the Bones, there may be less Chance of the former running into each other, than of the latter.

Ca

The

⁽a) Columb. De re Anatomica, lib. 1. cap. 2. Fallop. Expos. de Ossibus, cap. 11.

The Softness of the Ends of Bones may be of some Advantage in the Womb, and at Birth, after which the Offification begins at different Points to form *Epiphyses*, before the Offification can extend from the Middle to the Ends of

the Bones (a).

However folid and compact adult Bones are, yet they were once Cartilages, Membranes, nay, a mere Gelly. This needs no further Proof, than repeated Observations of Embryos when diffected: And how much more tender must the Bones be before that Time, when neither Knife nor Eye is capable to discover the least Rudiments of them? By Degrees they become more folid, then assume the Nature of Griftles, and at last offify; the Cohesion of their Plates and Fibres always increasing in Proportion to their increased Solidities; as is evident from the Time necessary to unrayel the Texture of Bones of People of different Ages, or of dense and of spongy Bones, or of the different Parts of the same Bone, and from the more tedious Exfoliations of the Bones of Adults than of Children.

After any Part of a Bone is fully offified, its Fibres are extended little more in Length at that Part, tho' they increase there in Thickness, and tho' their softer Parts continue to become longer (b).

As the Solidity of Bones increases, their Perioseum more easily separates from them. When Bones are membranous, the Perioseum

and

m

is

fre

pa

of

the

add

nul

the

bec

Thi

(a) Haller de studio medic. p. 267.

(b) Hale's vegetable Staticks, p. 393.—DuHamel Memoires de l'Acad. des Sciences 1742.

and they cannot be distinguished; they appear to be the same Substance. When they are Cartilages, their Membrane adheres so firmly to them, that it is difficult to separate them. Where the rigid bony Fibres are, the Periofteum is easily taken off.—Is the Similarity of Structure and consequent greater Attraction of the Membrane and Substance it incloses, while they are both flexible, the Cause of their greater Adhesion then? or is it owing to the Vessels, that go from the one to the other, being then larger? or do both these Causes combine to produce this Effect? Or is the Membrane or Cartilage, which becomes Bone afterwards, to be confidered as the same Substance with the Periosteum (a)? and must all these Plates of Bones be therefore faid to be Layers of the Periosteum hardened (b)?

The Offication of Bones depends principally on their Vessels being so disposed, and of such Diameters, as to separate a Liquor, which may easily turn into a bony Substance, when it is deprived of its thinner Parts; as seems plain from the Observation of the callous Matter, separated after Fractures and Ulcers, where Part of the Bone is taken out: For in these Cases, the Vessels extending themselves, and the Liquors added to them, are gradually formed into granulated Flesh; which sills up all the Space where the Bone is taken from; then hardens, till it becomes as firm as any other Part of the Bone. This happens frequently, even when the Ends

t

e

n

ts

at

K-

e-

e.

m.

um nd

Me-

⁽a) Memoires de l'Acad. des Sciences, 1744.

⁽b) Memoires de l'Acad. des Sciences, 1743.

of the diseased Bone are at a considerable Di-

stance from each other (a).

The Induration of Bones is also greatly asfifted by their being exposed, more than any other Parts, to the strong Pressure of the great Weights they support, to the violent Contraction of the Muscles fixed to them, and to the Force of the Parts they contain, which endeavour to make Way for their own further Growth. By all this preffing Force, the folid Fibres and Vessels of Bones are thrust closer; and such Particles of the Fluids conveyed in these Vessels as are fit to be united to the Fibres, are fooner and more firmly incorporated with them, while the remaining Fluids are forcibly driven out by the Veins, to be mixed with the Mass of Blood. In Consequence of this, the Vessels gradually diminish as the Bones harden. From which again we can understand one Reason, why the Bones of young Animals sooner reunite after a Fracture, than those of old, and why Cattle that are put too foon to hard Labour, feldom are of fuch large Size as others of the fame Brood who are longer kept from Labour.

That the offifying of Bones greatly depends on Pressure, seems to be evinced from the frequent Examples we meet with of other Parts turning bony, when long exposed to the pressing Force of the surrounding Parts, or when they are subjected to the like Circumstances by their own frequent and violent Contrac-

tion.

t

li

VI

ea

Ac

loc

med

de l'

(6

205.

⁽a) Hildan. de Vuln. graviss.—Med. Essays. Vol. I. Art. 23.—Job. a Meckren Obs. 69.—Mem. de l'Acad. des Sciences, 1742.—See a Collection of such Cases in Boehmer de Ossium Callo.

tion. Witness, the Bones found frequently near the Base of the Heart in some old Men (a), and in several other Creatures. Nay, the muscular Substance of the Heart has been offissed in such (b), and the Arteries of old Men often become bony.—The Cartilages of the Larynx are generally offissed in Adults.—In Beasts of Burden, the Cartilages between the Vertebræ of the Back very often change into compleat Bones; and, being intimately united with the Vertebræ, the whole appears one continued Bone—Nor is the Periosteum exempted from such an Induration (c).

S

e

it

of

a-

m

n,

te

17

elhe

ur.

ds

re-

rts ef-

ces ac-

Art.

ces,

To confirm this Argument Itill further, we may observe, That Bones begin their Offification at the Places where they are most expofed to these Causes, viz. in the cylindrical Bones from a middle Ring, and in the broad ones, at or near their Center, from one or more distinct Points. The Reason of which is, That these Parts are contiguous to the Bellies of the Muscles annexed to the Bones, where the Swelling of these moving Powers is greatest. What the Effects of this may be, let any judge, who view some of the Bones, as the Scapula, and Offa Iliam, which are covered with Muscles on each Side; how compact and thin they are in Adults, where the Bellies of the Muscles were lodged; whereas in Children they are thicker.

But

⁽a) Riolan. Comment. de Offib. cap. 32.—Bartholin. Hist. medic. Cent. 1. Hist. 50.—Ibid. Cent. 2. Hist. 45.

⁽b) Chefelden, Anatomy, Book 1. Introd.—Garengeot, Hist. de l'Acad. des Sciences, 1726.

⁽c) Peyer. Ephemerid. German. Decur. 2. Ann. 7. Observ.

But this being the middle Part of these Bones. where the greatest Number of Fibres is, this particular Place would have been much thicker in Adults, had not this forcible Cause been applied, which has not had fuch Effects in Children, whose Muscles have not been much exercifed.—Besides, if we allow that all the Parts of a Bone are equally increased by the constant Supply of new Particles, each Fibre, and every Particle of a Fibre, endeavours to make Way for its own Growth; by pulling the one next to it; and consequently by far the greatest Presfure is on the Middle, to make the Particles firm, and therefore to begin their Offification there. Lastly, the Pulsation of the medullary Arteries, which enter the Bones near to this middle Part, may, as Authors have alledged, contribute perhaps somewhat to this Induration.

From the Effects of Pressure only it is, that we can account for the Bones of old People having their Sides so much thinner, yet more dense and solid, while the Cavities are so much larger, than in those of young People; and for the Prints of Muscles, Vessels, &c. being so much more strongly marked on the Surfaces of the former than of the latter, if they belong to People of near the same Condition in Life.—Pressure must likewise be the Cause, which, in People of equal Ages, makes these Prints stronger in the Bones of those who had much Labour and Exercise, than they are in People who have

led an indolent unactive Life.

Perhaps both the Causes of Offisication above mentioned, may be affished by the Nature of the Climate People live in, and the Food

they

ha

du

Bo

the

the

plia

froy,

Num

1742 (c) Gagli

Decui

(6)

they use. Whence, in hot Countries, the Inhabitants sooner come to their Height of Stature, than in the Northerly cold Regions: And thence seems to have arisen the common Practice among the Ladies, of making Puppies drink Brandy or Spirit of Wine, and of bathing them in these Liquors, to prevent their growing big. Nay, it has been observed, that much Use of such Spirits has occasioned Parts, naturally soft, to petrify in some, and to offisy in other People of no great Age (a).

From the foregoing Account of the Structure of Bones, and of their Offification, we may understand the Reasons of the following Phæno-

mena.

0

S

n

19

d,

17.

at

le

re

ch

Ol'

fo

of

to

in

ng-

our

ave

a -

ure

ood

hey

How Bones may be fostened or dissolved.

How the natural Colour of Bones may be

changed by some Sorts of Food (b).

Why the Bones of some People are so long in hardening, and in others never compleatly indurate.

Why, in such whose Offisication is slow, the Bones are generally thicker in Proportion to their Lengths, especially at their Ends; as in the Rickets.

How hard firm Bones have become fost and pliable by Diseases (a).

How

(a) Littre, Histoire de l'Acad. des Sciences, 1706.—Geoffroy, Memoires de l'Acad. des Sciences, 1706.

(b) Philosoph. Transact. 442. Art. 8. Num. 443. Art. 2. Num. 457. Art. 4. Mem. de l'Acad. des Sciences, 1739,

1742.

⁽c) Histoire de l'Acad. des Sciences, 1700.—Mem. 1722.—
Gagliardi, Anatom. ossium, cap. 2. observ. 3.—Ephem. Germ.
Decur. 1. ann. 1. obs. 37. & Schol. Decur. 2. ann. 7. obs.
212. 235. Decur. 3. ann. 2. obs. 3.—Philos. Trans. No.
470. § 3. ibid. Vol. 48. § 4 and 44.

How Bones exfoliate by the rifing of granulated Flesh from their Surface.

How Bones may waste and diminish (a).

How Bones may become folid all through,

without any Appearance of Cancelli (b).

Why in some Diseases Epiphyses separate from Bones (c), and the End of Bones come afunder many Years after their Fractures appeared to be cured (d).

How Nodes, Tophi, and Exostoses, happen after the Erosion of the external Plates of Bones in the Lues Venerea, Scurvy, Rheumatism, and

· How and from what Callus is formed after a

Fracture (e).

Why Callus appears to be rather the continued Substance of the Periosteum than of the Bone, while it continues foft and flexible; but feems continued with the Bone after it offifies whening, and in others never completel, (A)

Why Callus is fensible, while it is foft, but

becomes infensible when it hardens. It was

What occasions sometimes such Difficulty in curing fractured Bones; or why they never reunite, though they are reduced, and all proper but hot enforce event cannot must brish Means

ishtely District (a).

0

ar

th

th

(e)

Su

W

by

ver

(a

Adve

Philo

rad.

(6)

(c)

(d)

(e)

(b) Ruysch. Thesaur. 2. Arc. 5. Thes. 3. Loc. 1. Num. 5. Thel. 9. Num. 2. not. 3.—Boehmer de Callo Offium.

(e) Memoires de l'Acad. des Sciences, 1741. Dehtleef de

Offium Callo. loli 12--- . . . ido . : . . ida

⁽a) Chefelden, Anat Book to Introd .- Hift. de l'Acad. des Sciences, 1700. 2017 antining

⁽c) Memoires de l'Acad. des Sciences, 1699. - Diemerbrock Anat. lib. gocap. 19. - Cowper's Anat. Explic. Tab. 96. Fig. 1. . (d) Aufon's Voyage. We and maile of the

⁽f) Mem. de l'Acad. 1741.

Means towards a Cure are used (a).—Are the Bones of a Woman with Child more tedious in reuniting, than those of others (b)?

Why Calluses, after Fractures, are sometimes

very thick and protuberant.

What Difference there ought to be in the Application of Bandages to Fractures of the

Bones of old and of young Patients.

How Bones, remaining long unreduced after a Luxation, may have their Form so changed, as to make their Reduction very difficult, if not

impossible (c).

d

2

].

10

ut

es

ut

in

re-

per

ans

des

n. 5.

rock

g. 1.

E de

Whoever is desirous to know, in what Time and Order each Bone, and its several Parts, begin to assume a bony Nature, let him consult Kerckringius (d), who gives us the Delineations of Abortions from three Days after Conception, and traces the Ossisication of the Bones from three Weeks, and a Month, till the Time of the Birth: To whom should be added Coiterus (e), and Eyssimus (f). An Account of this Subject might also be collected out of Ruysch's Works, where some of the Mistakes committed by the former Authors are corrected: And several more Particulars to make the History of the

(b) Hildan. Centur. 5. Obs. 87. & Cent. 6. Obs. 68.— Philos. Transact. No. 494. § 21.

(d) Anthropograph. Ichnograph. & Osteogenea Fætuum.

(e) De offibus fœtûs abortivi.

⁽a) Meckren. Observ. Medico-Chirurg. Obs. 71.—Ruysch. Advers. Dec. 2. § 2. Observ. Anat. Chir. obs. 4.

⁽c) Saltzman. Observ. Decur. Obs. 6.—Memoires de l'Acad de Chirurgic. Tome 2. p. 155.

⁽f) De offibus Infant. cognoscend. & curand.

the Osteogenea more accurate, have fince been

added by Nesbitt (a) and Albimis (b).

I must refer to the Authors now quoted for the more curious Part of the human Osteogeny; not having Preparations enough to give such a full History of it as is done by them. But I shall endeavour to explain the more useful and necessary Part of the Osteogeny, by subjoining to the Description of each Bone of an Adult, its Condition in ripe Children; that is, in such as are born at the ordinary Time; and shall point out what Parts of each are afterwards joined in Form of Epiphyses. This, with the following general Rules, seem to me sufficient for understanding what of this Subject is necessary in the Practice of Physick and Surgery.

1. Where-ever I mention any Parts being cartilaginous, or their being still separable from the other Parts of the Bone to which they belong, I would be understood to hint, that, about seven or eight Years of Age, such Parts are offissed and united to their proper Bones, unless when it is said, that they are afterwards formed

into Epiphyses.

2. Such as become Epiphyses, are generally offissed at seven or eight Years of Age; but, being for the most Part moistened by Synovia, their external Surface is still somewhat cartilaginous, and they are not yet united to their Bones.

3. At eighteen or twenty Years of Age, the Epiphyses are entirely offified, and have blended

their

E

tl

di

W

Ca

fer

mu

per

we the

miss cal

Claf

riety nece

the v

and t

exic,

a, Cor

(a) Human Osteogeny explained.
(b) Icones ossium sœtús humani accedit osteogeniæ brez historia.

their Fibres fo with the Body of the Bone, as to make them inseparable without Violence.

The Knowledge of this Part of the Ofteogeny, I think necessary to prevent dangerous Mistakes, in the Cure of several Diseases. As for example: Without this Knowledge, the Separation of an Epiphyse might be mistaken for a Fracture or Luxation. The interstice of two Parts of a Bone not yet joined, might be judged to be a Fissure. — A Diastasis, or Separation of such disjoined Pieces of a Bone, might be thought a Fracture. — The Protrusion of one Piece, or its overloping any other, could be mistaken for an Excrescence or Exostosis.—Such Errors about the Nature of a Disease, would give one very different Indications of Cure, from what he would have, if he really understood his Patient's Case. And very often the Knowledge of the different Inequalities on the Surfaces of Bones, must direct us in the Execution of what is proper to be done to cure several of their Diseases.

ľ

g

m

e-

a.

ess

red

ally

be.

via, tila

heir

the

ded

revi

Having thus confidered the Bones when single, we ought next to shew the different Manner of their Conjunctions †. To express these, Anatomists have contrived a great Number of technical Terms; about the meaning, Propriety and Classing of which, there has unluckily been Variety of Opinions. Some of these Terms it is necessary to retain, since they serve to express the various Circumstances of the Articulations, and to understand the Writers on this subject.

D The

[†] Σύνταξις, σύνθεσις, συμβολή, όμιλία, Compositio, Contexic, Articulatio, Conjunctio, Nodus, Commissura; Structua, Compages.

The ARTICULATIONS are most commonly divided into three Classes, to wit, Symphysis,

Synarthrofis, and Diarthrofis.

Symphysis, which properly signifies the Concretion or growing together of Parts, when used to express the Articulations of Bones, does not seem to comprehend, under the Meaning generally given to it, any thing relating to the Form or Motion of the conjoined Bones; but by it most Authors only denote the Bones to be connected by some other Substance; and as there are different Substances which serve this Purpose, therefore they divide it into the three following Species:

1. Synchondrosis*, when a Cartilage is the connecting Substance: Thus the ribs are joined to the Sterman; thus the Bodies of the Vertebræ are connected to each other; as are likewise

T

L

na

112

th

tic

nec

wid

et,

Cav

the

(6)

the Offa Pulis.

2. Symeurosis, or Syndesmosis, when Ligaments are the connecting Bodies, as they are in all the moveable Articulations.

3. Sysfarcosis, when Muscles are stretched from one Bone to another, as they must be

where there are moveable Joints.

The fecond Class of Articulations, the Synarthrofis, which is said to be the general Term by which the immoveable Conjunction of Bones is

expressed, is divided into three Kinds.

two Bones are mutually indented into each of ther, or as if they were fowed together, and is formed by the Fibres of two Bones meeting while

^{*} Amphiarthrofis.

[†] Papn,

while they are yet flexible and yielding, and have not come to their full Extent of Growth; fo that they mutually force into the Interstices of each other, till, meeting with fuch Refistance as they are not able to overcome, they are stopped from sprouting out farther, or are reflected; and therefore these Indentations are very different both in Figure and Magnitude: Thus the Bones of the Head are joined; thus Epiphyles are joined to the Bones, before their full Connexion and Union with them.

Under this Title of Suture, the Harmonia of the Antients may be comprehended; scarce any unmoved Bones being joined by plain Surfaces and Caviles. m. (a)

t

C

e

18 is

ee

ne ed

te-

ife

rain

ned

be

lar. by

3 18

rere

1 0.

nd is

ting hile

2. Gomphofis * is the fixing one Bone into another, as a Nail is fixed in a Board: Thus the Teeth are secured in their Sockets.

3. Schindylesis or Ploughing (b), when a thin Lamella of one Bone is received into a long narrow Furrow of another: Thus the Processis azygos of the Sphenoid, and the nafal Process of the ethmoid Bone, are received by the Vomer.

The third Clais, or Diarthrofis |, is the Articulation where the Bones are fo loofely con-This is fubdinected as to allow large Motion.

vided into three Kinds.

The first is Enarthrosis, or the Ball and Socket, when a large Head is received into a deep Cavity; as the Head of the Os Femoris is intothe Acetalulum Coxendicis.

> D 2 The

⁽a) Vefal. Observ. Fallop. Examen. Conclavatio.

⁽b) Keil, Anat. chap. 5. § 3. Απαρθρώσις, Dearticulatio, Abarticulatio.

The fecond is Arthrodia, when a round Head is received into a superficial Cavity; as in the Articulation of the Arm-bone and Scapula. These two Species of Diarthrosis allow Motion to all Sides.

The third is Ginglimus &, which properly fignifies the Hinge of a Door or Window; in it the Parts of the Bones mutually receive and are received, and allow of Motion two Ways: Workmen call it Charnal.

The Ginglimus is generally divided into three Kinds, to which fome (a) give the Names of

To

Ĭe

Sy fir

an

tic

th

Ex

ga

ma

Syn

hay

cies

the

in t

tedi

nio

Det

t is

too vhi

W

or a

contiguous +, distant 1, and compound |.

The first Kind of Ginglianus is, when a Bone has several Protuberances and Cavities, which answer to as many Cavities and Processes of the other Bone, with which it is articulated; as in the Conjunction of the Femur with the Tibia.

The second Species is, when a Bone receives another at one End, and is received by the same Bone at the other End; as in the Radius and

Ulna.

The last Sort is, when a Bone receives another, and is received by a third; as in the ob-

lique Processes of the Vertebræ.

When I first mentioned the Articulations of Bones, I said there were different Opinions concerning the Use of their technical Names, e.g.; It has been said, that Symphysis should be the Name for the immoveable Articulations, and that

§ Articulatio mutua.
(a) Baker, Curf. Offeolog. Demonstr. s.

[†] Proximus. † Longus. † Compositus.

that by Synarthrofis should be understood the Conjunction of Bones by some connecting Medium.—Those who have taken Symphysis in the Sense I did, of its expressing the Conjunction of Bones with a connecting Substance, have difagreed in their Definition of it; some inserting, and others leaving out, its allowing Motion .-Where they have agreed in their Definition, they have not been of the same Mind concerning the Species of it. For feveral think the Syfarcosis and Syndesmosis applicable to so many Joints which are univerfally allowed to be claffed under the Diarthrofis, that it must create Confusion to name them by any Species of the Symphysis. Few keep to such a general Definition of the Synchondrosis as I have done; and, whether they determine it to allow no Motion, or an obscure or a manifest one, bring themselves into Difficulties, because there are Examples of all these three Kinds.—Some again, by too nicely distinguishing obscure and manifest Motions of Bones, have blended the Synarthrofis and Diarthrofis, and from thence have branched out the different compound Species of Articulations that may be formed of them so far, that they could find no Examples in the Body to illustrate them by.—It would be tedious to enumerate more of the jarring Opinions, and it would be far more fo to give a Detail of the Arguments used by the Disputants. t is sufficient for my Purpose, that it is underfood in what Sense I take these technical Terms; which I do in the following Manner.

f

h

e

n

es

91

nd

0-

b-

of

n-

j.;

nd

at

When I mention the Symphysis or Synarthrosis, or any Species of them, I shall always under-

D 3 stand

stand them according to the Explication already given of them. But though the preceding Account of the Diarthrosis, or Articulation of moveable Bones, has been almost universally received; yet seeing it does not comprehend all the moveable Articulations of the Body, and one of its Species does not answer to any Notion we can have of the Conjunction of two Bones, I must beg Leave to change the Definitions and

Kinds of these Joints.

I would call Diarthrosis that Conjunction of Bones, whereby they are fitted for Motion, being each covered with a smooth Cartilage, connected by one or more common Ligaments, and lubricated with Liquor at the conjoined Parts. In which Definition, I have no Regard to the Quantity of Motion which they really do perform; the Motion being often confined or enlarged by some other Cause not immediately depending on the Frame of the two Surfaces of the Bones, forming the particular Joint, which then is considered.

The first Species of the Diarthrosis, viz. the Enarthrosis or Ball and Socket, I would define more generally than above, That Articulation where a round Head of one Bone is received into a Cavity of another, and consequently, without some foreign Impediment, is capable of Motion to all Sides. Examples of this Kind are to be seen in the Articulation of the Thigh-bone and Ossa innominata; Arm-bone and Scapular Astrogalus and Os Naviculare; Magnum of the Wrist, with the Scaphoides and Lunare; first Bone of the Thumb with the second, &c.

The

0

fa

ec

tic

w

do

No

fec

mu

Art

low

Wit] Kne cof

y

ıll

nd ti-

es.

nd

of

e-9c

n-

nd

ts.

the

er.

en-

de-

of

ich

the

fine

tion

in.

ith.

Mo.

e to

one

ula

the

The

The fecond Sort, or the Arthrodia, differing from the Enarthrofis, in the preceeding Account, only in the Cavity's being more superficial, which makes no effential Difference, especially that, in the recent Subject, Cartilages or Ligaments supply the Deficiency of Bone, ought, in my Opinion, to be called, with Vesalius (a), that Articulation of two Bones adapted for Motion, where it is not at first Sight obvious which of the two has the Head or Cavity, or where they are joined by plain Surfaces, or nearly fo: such is the Conjunction of the Clavicle with the Scapula; Offa Cuneiformia with the Os Naviculare; Metatarsat Bones with the Ossa Cuneiformia, &c. From the Nature of this Sort of Joint, it is plain, that very great Motion cannot be allowed, without the Bones going farther out of their natural Situation, than is convenient or fafe.

Ginglimus, I would reckon that Articulation by the Form of which the Motion of the joined Bones must be chiefly confined to two Direc-

tions, which Hinges of Doors are.

The first Species of this is the Trochoides, when one Bone turns on another, as a Wheel does on its Axis: Thus the first Vertebra of the Neck moves on the Tooth-like Process of the fecond. This is the most proper Kind of Ginglimus.

The fecond Species should be esteemed that Articulation, where several prominent and hollow Surfaces of two Bones move on each other, within the same common Ligament; as in the Knee, Elbow, &c.

The

⁽a) De corp. human. fabrica, lib. 1. cap. 4.

The third Sort of Ginglimus is, when two Bones are articulated to each other at different Parts, with a distinct Apparatus of the motory Machines at each; fuch is the Articulation of the Os Occipitis with the first Vertebra of the Neck, of any two contiguous Vertebræ by their. oblique Processes, of the Ribs with the Bodies and transverse Processes of the Vertebræ, of the Radius with the Ulna, Tibia with the Fibula,

Astragalus with the Calcaneum, &c.

I would entirely throw out what is commonly called the third Kind of Ginglimus: For, in examining the Conjunction of a Bone with two others, as in the common Example of a Vertebra joined with the one above and below, the Connexion of the middle one with each of the other two, ought to be confidered separately; otherwise we might with the same Propriety esteem the Articulations that the long Bones, the Femult, Tibia, Humerus, &c. have at their different Ends, as one Articulation; which is abfurd.

If the moveable Bones were not connected and kept firm by some strong Substance, they would be luxated at every Motion of the Joints; and if their hard rough unequal Surfaces were to play on each other, their Motion would not only be difficult, but the Loss of Substance from Attrition would be great: Therefore Ligaments are made to obviate the first, and Cartilages to prevent the other Inconveniency. But because Ligaments and Cartilages turn rigid, inflexible and rough, unless they are kept moift, a sufficient Quantity of proper Liquors is supplied for their Lubrication, and to preserve them in a

flexible

a

e

0

W th

ca La

of

io

cuo

Tru ull

S

ies,

per

al nd:

iqu

prce

he C

Th

g :

ould

flexible State. Seeing then these Parts are so necessary to the Articulations, I shall next consider their Structure, Situation and Uses, so far as they are subservient to the Bones, and their Motions.

LIGAMENTS | are white flexible Bodies, thicker and firmer than Membranes, and not so hard or firm as Cartilages, without any remarkable Cavity in their Substance, difficultly stretched, and with little Elasticity; serving to connect one Part to another, or to prevent the Parts to which they are fixed from being removed out of that Situation, which is useful and safe.

After Maceration in Water, the Ligaments can easily be divided; and each ligamentous Layer appears composed of Fibres, the largest of which are disposed in a longitudinal Direction.

a

.

)-

le

f-

b-

ed

ey

ts;

ere

ot

mc

nets

to

use

ble ici-

for

a ble The Arteries of Ligaments are very conspimons after a tolerable Injection, and the larger Frunks of their Veins are sometimes to be seen full of Blood.

Such Ligaments as form the Sides of Caviies, have numerous Orifices of their Arteries
pening upon their internal Surface, which keep
t always moist: If we rub off that Moisture,
and then press the Ligament, we can see the
iquor outing out from small Pores; and we can
bree thin Liquors injected by the Arteries, into
the Cavities formed by Ligaments.

These exhalant Arteries must have corresponding absorbent Veins, otherwise the Cavities ould soon be too full of Liquor.

Ligaments

Σύνδισμοι, νεύροι, Copulæ, Vincula.

Ligaments then must be subject to the Diseases common to other Parts, where there is a Circulation of Fluids, Allowance always being made for the Size of Vessels, Nature of the Fluids, and Firmness of the Texture of each Part.

Authors generally fay, That Ligaments are infensible; and consequently it may be inferred, that they have no Nerves bestowed on them. But the violent racking Pain selt on the least Motion of a Joint labouring under a Rheumatism, the Seat of which Disease seems often to be in the Ligaments, and the insufferable Torture occasioned by Incisions of Ligaments, and by a Collection of acrid Matter in a Joint, or by Tophi in the Gout, would persuade us, that they are abundantly supplied with Nerves.

The Ligaments, which connect the moveable Bones, commonly rife from the Conjunction of the Epiphyses of the one Bone, and are inferted into the same Place of the other; or where Epiphyles are not, they come out from the Cervix, and beyond the Supercita of the articulated Bones; and after fuch a Manner, in both Cases, as to include the Articulation in a Purse or Bag, with this Difference, depending on their different Motions, that where the Motion is only to be in two Directions, the Ligaments are strongest on those Sides toward which the Bones are not moved; and when great Variety of Motions is designed to be al lowed, the Ligaments are weaker than in the former fort of Articulations, and are nearly the same Strength all round.

Part

b

C

fh

bu

16

1t

the

by

fev

the

Lig

Bon

Hol

Bon

nien

Part of the capfular Ligaments is composed of the Periosteum, continued from one Bone to another, as was observed P. 2. and their internal Layer is continued on the Parts of the Bone or Cartilage which the Ligament includes (a.)

Besides these common capsular Ligaments of the joints, there are particular ones in feveral Places, either for the firmer Connexion of the articulated Bones, or for restraining and confining the Motion to some one Side; fuch are the cross and lateral Ligaments of the

Knee, the round one of the Thigh, &c.

1.

A

a-

to

r.

nd

or

at

e-

n-

nd

er;

out

of

er,

in

nd.

the

the

irds

en a

al.

the

y of

Pari

From this Account of the Ligaments, we may conclude, that, cæteris paribus, in whatever Articulation, the Ligaments are few, long and weak, the Motion is more free and quick; but Luxations happen frequently: And, on the contrary, where the Ligaments are numerous, thort and strong, the Motion is more confined; but such a Joint is less exposed to Luxations (b).—Whence we may judge how necessary it is to attend to the different Ligaments, and the Changes which have been made on them by a Luxation, when it is to be reduced.

Ligaments also supply the Place of Bones in feveral Cases to Advantage: Thus the Parts in the Pelvis are more fafely supported below by Ligaments, than they could have been by Bone. The Ligaments, placed in the great Holes of the Offa innominata, and between the Bones of the Fore-arm and Leg, afford convenient Origin to Muscles.-Immoveable Bones

are

⁽a) Nesbit. Osteogen .- Philos. Transact. No. 470. §. 6. (b) Fabric. ab Aquapend. de Articul. Part. utilit. Pars 3.

are firmly connected by them; of which the Conjunction of the Os facrum and immoninatum is an Example.—They afford a Socket for moveable Bones to play in, as we see Part of the Astragalus does on the Ligament stretched from the Heel-bone to the Scaphoid.

Numerous Inconveniencies may arise from too long or short, strong or weak, lax or rigid

Ligaments.

CARTILAGES (a) are folid, smooth, white, elastick Substances, between the Hardness of Bones and Ligaments, and covered with a Membrane, named *Perichondrium*, which is of the same Structure and Use to them, as the Pe-

riosteum is to the Lines.

Cartilages are composed of Plates, which are formed of Fibres, disposed much in the same Way as those of Bones are; as might be reasonably concluded, from observing Bones in a cartilaginous State before they offify, and from seeing, on the other Hand, so many Cartilages become bony. This may be still farther confirmed, by the Exsoliation which Cartilages are subject to, as well as Bones.

The Perichondrium of several Cartilages, for Example, those of the Ribs and Larynx, has Arteries which can be equally well injected with those of the Periosteum, but the Vessels of that Membrane in other Parts, e. g. the articular Cartilages, do not admit the coloured Injections, and in none of them does the Injection enter deep into the Substance of the Cartilage; nay, Madder, mixed with the Food of

Animals,

d

b

tl

of

Bo

01

bu

offi

Mo

óf

fim

ed (

Enc

the

Bon

(a) No. 4. 742.- Animals, does not change the Colour of Car-

tilages, as it does of Bones (a).

The granulated Flesh which rifes from the Ends of metarcarpal or metatarfal Bones, when the Cartilage exfoliates, after a Finger or Toe has been taken off at the first Joint, is very fensible, from which the Existence of Nerves

in Cartilages may be inferred.

2.

re

10

a.

m

n-

arc

for

has

ted

s of

rti-

In-

etirtiof uls, While Cartilages are in a natural State, it is to be remarked, first, That they have no Cavity in their Middle for Marrow. Secondly, That their outer Surface is softest, which renders them more flexible. Thirdly, That they do not appear to change their Texture so much by Acids as Bones do. And, lastly, That, is the specifiek Gravity of Cartilages is near a third less than that of Bones; so the Cohesion of their several Plates is not so strong as in Bones: Whence Cartilages laid bare in Wounds or Ulcers, are not only more liable to corrupt; but exsoliate much sooner than Bones do.

Cartilages feem to be principally kept from offifying, either by being subjected to alternate Motions of Flexion and Extension, the Effects of which are very different from any Kind of simple Pressure, or by being constantly moistened (b): Thus, the Cartilages on the articulated Ends of the great Bones of the Limbs, and the moveable ones placed between the moving Bones in some Articulations, which are obliged

(a) Philos. Transact. No. 442. Art. 8. No. 443. Art. 4. No. 457. Art. 4.—Mem. de l'Acad. des Sciences 1739 & 742.—Dehtleef de Ossium Callo.

(b) Havers's Ofteolog. nov.

to fuffer many and different Flexions, and are plentifully moistened, scarce ever change into Bone; while those of the Ribs and Larynx are often offified .- The middle angular Part of the Cartilages of the Ribs, which is constantly in an alternate State of Flexion and Extension, by being moved in Respiration, is always the last of becoming bony.—In the Larynx, the Epiglottis, which is oftener bended and more moistened than the other four Cartilages, seldom is offified, while the others as feldom e-

fcape it in Adults.

Cartilages, which have little Flexion or Moisture, such as those of the Ribs or Larynx, of ten begin to offifyon their external Surface, and the Offification proceeds internally, till the Cancelli are at last formed, when Marrow is deposited into them. While this Change is a bringing about in the Substance of Cartilages, their Blood-vessels gradually appear bigger towards their internal Substance. This Change of a Cartilage from a natural State, to its becoming Bone, is at first View surprising; and, fo far as I know, none has attempted to explain it; which I mention only, to have fome Plea for an Apology, if I should be thought to have failed in accounting for this Phænomenou.

The Cartilages, being naturally of a firm Substance, and of a Composition a-kin to that of Bones, gradually acquire greater Soliding by constant Pressure; and this Change is made foonest and most remarkably where the Pres fure is greatest; that is, at their external Surface.—The exterior Plates, when offified, are more compact and dense than formerly, and transfe

therefore

n

ar

Po

Bo

the

do all

Pla

a ca

Mid had

Fibr

of th Cance

distri

fened

therefore have a stronger Power of attracting those in Contact with them; while the Branches of Vessels distributed to the first ossified Plate, and those that run in the Interstices of the Fibres of this and of the Plate next to it, being now more compressed than formerly, have a less Quantity of Fluids passing through them; and, confequently, the other Branches receive a proportionally larger Quantity; the Momentum of which, joined to that Power of Attraction or Force of Cohesion, increased by the greater Solidity of the Plates, increases the Preffure upon the more internal Plates, and hastens their Hardening; after which these last Plates produce the same Effects on the others contiguous to them: And thus the Offification goes on, till all are become bony.—The Body thus changed, has its former Dimensions, or nearly fo; because its external Part offified first, and being rigid, yields little or nothing to the Powers that draw it towards the Axis of the Bone. But feeing the new Particles, added from the circulating Fluids, during the Offification, do not compensate for the Condensation which all the Particles undergo, and thereby the Plates occupy less Space than they did while in a cartilaginous State, a Cavity is left in the Middle. And as all the Flates cohered, and had cross Fibres joining them, many of these Fibres are stretched irregularly from one Side of the Cavity to the other, and therefore form Cancelli.—The Branches of the Vessels formerly distributed to the Plates being now much lesfened, the remaining Branches, which run transversely, are consequently proportionally E 2

f.

ne

e.

2

es,

to-

ge

be-

nd,

ex.

t to

21011.

firm that

dity

ref.

Sur-

are

and fort increased, become very conspicuous, and are dispersed every where in the Cavity, to serve for the Secretion of Marrow.—Thus, this flexible elastick solid Substance, becomes a rigid in flexible cavernous Bone, with Marrow contain-

ed in its Cancelli.

The Cartilages subservient to Bones, are sometimes found on the Ends of Bones which are joined to no other; but are never wanting on the Ends, and in the Cavities of such Bones as are designed for Motion (a). Cartilages also are interposed between such other Cartilages as cover the Heads and Cavities of articulated Bones; nay, they are also placed between im-

I

S

fe

iı

as

th

til

th

en

pla

Ec

Bo

cou

whi

Body

(a)

No. 46

(b) I Thef.

Shor

moveable Bones.

The Uses of Cartilages, so far as they regard Bones, are, to allow, by their Smoothness, such Bones as are defigned for Motion, to slide ea fily without Detrition, while, by their Flexibi lity, they accommodate themselves to the several Figures necessary in different Motions, and by their Elasticity, they recover their natural Polition and Shape, as foon as the Pressure is removed, - This springy Force may also al fift the Motion of the Joint to be more expe ditious, and may render Shocks in running jumping, &c. less. - To these Cartilages we chiefly owe the Secrity of the moveable Articulations: For without them the bony Fibra would fprout out, and intimately coaleice with the adjoining Bone; whence a true Anchylofa must necessarily follow; which never fails to happen when the Cartilages are eroded by Crie

⁽a) Celf, de re Medic. lib, 8, cap. 1.

crid Matter, or offified from Want of Motion or Defect of Liquor, as we see often happens after Wounds of the Joints, Paidarthrocace, Scrophula, and Spina ventofa, or from old Age, and long Immobility of Joints (a).-Hence we may know what the Annihilation is, which is faid to be made of the Head of a Bone, and of the Cavity for lodging it, after an unreduced Fracture (b). The moveable Cartilages interposed in Joints, serve to make the Motions both freer and more fafe than they would otherwise be.—Those placed on the Ends of Bones that are not articulated, as, on the Spine of the Os Ilium, Base of the Scapula, &c. ferve to prevent the bony Fibres from growing out too far.—Cartilages fometimes ferve as Ligaments, either to fasten together Bones that are immoveably joined, such are the Cartilages between the Os facrum, and Offa Ilium, the Offa Pubis, &c. or to connect Bones that enjoy manifest Motion, as those do which are placed between the Bodies of the true Vertebra, &c. - Cartilages very often do the Office of Bones to greater Advantage than these last could; as in the Cartilages of the Ribs, those which supply Brims to Cavities, &c.

Too great Thickness or Thinness, Length or Shortness, Hardness or Suppleness of Cartilages, may therefore cause great Disorders in the

Body.

h

es

10

25

ed

m-

ard

ich

ea.

vend,

ara

e is

spe.

ing

W

Arti-

brewith

logu

s to

y a

crie

E 3

The

(a) Columb. de re anat. Lib. XV.—Dessandes Hist. de l'Acad. des Sciences 1716.—Phil. Transact. No. 215.—Ibid. No. 461. § 16.

(b) Hildan. de Ichor. et Melicer. acri Celfi, cap. 5.- Ruysch. Thes. 8. No. 103. - Saltzman in Act. Petropolit. Tom. 3.

P. 275.

The Liquor, which principally ferves to moisten the Ligaments and Cartilages of the Ar. ticulations, is supplied by Glands, which are commonly fituated in the Joint, after fuch a Manner, as to be gently prefled, but not destroyed by its Motion. By this Means, when there is the greatest Necessity for this Liquor. that is, when the most frequent Motions are performed, the greatest Quantity of it must be separated. These Glands are fost and pap. py, but not friable; In fome of the large Joints they are of the conglomerate Kind, or a great Number of small Glandules are wrapt up in one common Membrane. Their excretory Ducts are long, and hang loofe, like fo many Fringes, within the Articulation; which, by its Motion and Pressure, prevents Obstructions in the Body of the Gland, or its Excretories, and promotes the Return of this Liquor, when fit to be taken up by the absorbent Vesfels, which must be in the joints, as well as in the other Cavities of the Body; and, at the fame Time, the Preffire on the excretory Ducts hinders a superfluous unnecessary Secretion, while the fimbriated Disposition of these Excretories does not allow any of the secreted Liquor to be pulhed back again by these Canals towards the Glands (a).

n

ap

ee

la

oi

de

T

ne

hic

ie S

arai

ie

em In

iv S

ey h

Very often their Fountains of flimy Liquor appear only as a Net-work of Veslels. - Frequently they are almost concealed by cellular asses imall simple niucous Foliculi may be seen (b) em:

(a) Courper, Anatomy Explicat tab. 79. litt. E. E. (b) Morgagn. Adversat. 2. Animad. 23.

The different Joints have these Organs in different Numbers and Sizes; the conglomerate ones don't vary much, especially as to Situation, in the similar Joints of different Bo. dies, but the others are more uncertain.

Upon pressing any of these Glands with the Finger, one can squeeze out of their Excreories a mucilaginous Liquor, which fomewhat resembles the White of an Egg, or Serum of he Blood, but it is manifestly Salt to the Taste. It does not coagulate by Heat, as the Serum loes; but turns first thinner, and, when evaporated, leaves only a thin falt Film. Diferent Salts have much the same Effect on it as n many other Juices of our Body.

The Quantity of this Mucilage, constantly applied, must be very considerable, fince we ee what a plentiful troublesome Discharge of lary Matter follows a Wound or Ulcer of any oint; of which Liquor the Mucilage is a con-

derable Part.

.

e

1

46

n

r,

re

ift

p.

ge

or

pt

e-10

ch, Ai.

to-

or, Tel-

s in

the

ory

cre-

hele eted

nals

roup

Fre-

ular

mes

The Vessels which supply Liquors for making he Secretion of this Mucilage, and the Veins hich bring back the Blood remaining after te Secretion, are to be feen without any Prearation; and, after a tolerable Injection of le Arteries, the Glands are covered with em.:

In a found State, we are not conscious of ly Senfibility in those Glands; but, in some ties which I have feen, when they inflame imes ad suppurate, the most racking Pain is selt in (h) em: A melancholly, tho' a sure Proof that The ey have Nerves.

Thefe

These mucilaginous Glands are commonly lodged in a cellular Substance; which is also to be observed in other Parts of the Bag formed by the Ligaments of the Articulation; and contains a fatty Matter, that must necessarily be attenuated, and forced through the including Membranes into the Cavity of the Join, by the Pressure which it suffers from the move

ing Bones, and an to and we at taken

If then the Oil is conveyed from this cel lular Substance; and if the attenuated Man row passes from the Cancelli of the Bones by the large Pores near their Ends, or in their Can ties, and fweats through the Cartilages then into the Articulations; which it may, when affifted by the constant Heat and Action of the Body, more easily do, than when it escape through the compact Substance of the Bone in a Skeleton: If, I fay, this Oil is fent to: Joint, and is incorporated with the Mucilage and with the fine Lymph that is constantly outing out at the Extremities of the small A teries distributed to the Ligaments, one the fittest Liniments imaginable must be produc ed; for the Mucus diluted by the Lymph con tributes greatly to its Lubricity, and the 0 preferves it from hardening. How well fut a Mixture serves the purpose it is designed for Boyle (a) tells us he experienced in working his Air-pump; for the Sucker could be move with much less Force after being moister ed with Water and Oil, than when he use either one or other of these Liquors: And I h

0

00

t

Boi

he

ity

he

oir

o co

rid,

equ

ne I

nall

hen

oife

† Mu

ap. 5.

⁽a) Physico-mechanic. Experim.

y

n.

nd Lly

ıd.

nt,

07.

el.

ar.

the

wi.

lere her

the

ape

one

to

age

nth

Ar e o

duo

con e Oi

fue d for

rkin

ister

I be

lieve every one, at first View, will allow the diluted Mucilage to be much preferable to simple Water. The Synovia, + as this Liquor composed of Oil, Mucilage and Lymph, is commonly now called, while in a found State, effectually preserves all the Parts concerned in the Articulations soft and slexible, and makes them slide easily on each other, by which their mutual Detrition and Overheating is prevented, in the Manner daily practised in Coach and Cart Wheels, by betmearing them with Grease and Tar.

After the Liquor of the Articulations becomes too thin and unferwiceable, by being constantly bounded and rubbed between the moving Bones, t is reassumed into the Mass of Blood by the blorbent Vessels.

When the Synovia is not rubbed betwixt the Bones, it impissates. And sometimes, when he Head of a Bone has been long out of its Caity, this Liquor is said to fill up the Place of he Bone, and hinder its Reduction; or if a joint continues long unmoved, it is also said to dement the Bones, and occasion a true Anhylosis (a).—If the Synovia becomes too arid, it erodes the Cartilages and Bones; as requently happens to those who labour under the Lues Venerea, Scurvy, Scrophulæ, or Spina entosa.—If this Liquor is separated in too nall Quantity, the Joint becomes stiff; and then with Difficulty it is moved, a crackling soile is heard, as People advanced in Years frequently

[†] Μυζα, Mucus, Axungia.
(a) Pare, Chirurgie. livre 15. chap. 18. & livre 16.
ap. 5.

frequently experience (a). - If the Mucilage and Lymph are deposited in too great Quantity, and the absorbent Vessels do not perform their Office sufficiently, they may occasion a Drop fy of the Joints (b). - From this fame Caule also the Ligaments are often so much relaxed as to make the Conjunction of the Bones ven weak: Thence arise the Luxations from an in ternal Cause, which are easily reduced, but di ficultly cured (c). - Frequently, when fuel a superfluous Quantity of this Liquor is pent up it becomes very acrid, and occasions a great Train of bad Symptoms; fuch as Swelling and Pain of the Joints, long finuous Ulcers and R stulæ, rotten Bones, Immobility of the Joint Marcor and Atrophia of the whole Body, held Fevers, &c. (d). From a Depravity in the Blood or Diseases in the Organs that furnish the Synovia of the Joints, it may be greatly chang ed from its natural State, it may be purule after Inflammation, mucous in the white Swell ings, gelatinous in the Rheumatism, chall from the Gout, &c. hence a great Variety Disorders in the Joints (e).

mently happens to the called the action

erodes the Cartisties and

gn

her Glar

A

he]

one

hicl

hose

only

(b) Hildan, de Ichore & Meliceria acri Celsi.

il agaict, an

⁽a) Galen. de usu Part, lib. 12. eap. 2. — Fabric. Aquapend. de Articul. part. utilitat. pars 3. — Barthof Hist. medic. Cent. 3. Hist. 11.

⁽c) Hippocrat. de locis in homine, S. 14. & de Articul (d) Hildan. de Ichore & Meliceria acri Celsi.

⁽e) See Reimar Dissert. de fungo Articular.

THE

ANATOMY

OFTHE

HUMAN BONES.

PART II.

Of the SKELETON.

HOUGH any dry Substance may be called Skeleton, yet, among Anatomists, this Word is universally understood to ignify the Bones of Animals connected togeher, after the Teguments, Muscles, Bowels, slands, Nerves, and Vessels are taken away *.

A Skeleton is faid to be a natural one, when he Bones are kept together by their own Liaments; and it is called artificial, when the ones are joined with Wire, or any other Subsance which is not Part of the Creature to hich they belonged. Small Subjects, and such those Bones are not fully offissed, are comsonly prepared the first Way; because, were

p.

111-

ich

up, eat

nts

the the anguler

vel

alk

ty

Th

E.

bric.

arthol

rticul

^{*} Cadaveris crates.

all their Parts divided, the nicest Artist could not rejoin them, by reason of their Smallness, and of the Separation of their unossified Parts; whereas the Bones of large adult Animals are soonest and most conveniently cleaned when single, and are easily restored to, and kept in their natural Situation.—Sometimes the Skeleton of the same Animal is prepared in both these Ways; that is, the smaller Bones are kept together by their natural Ligaments, and the larger ones are connected by Wires, or

some such Substances.

Before we proceed to the Division and particular Description of the Skeleton, it is worth while to remark, that when the Bones are put into their natural Situation, scarce any one of them is placed in a perpendicular Bearing to another; though the Fabrick composed of them is fo contrived, that, in an erect Posture, a perpendicular Line, from their common Center of Gravity, falls in the Middle of their common Base (a). On this Account, we can support ourselves as firmly, as if the Axis of all the Bones had been a streight Line perpen dicular to the Horizon; and we have much greater Quickness, Ease and Strength in seve ral of the most necessary Motions we perform It is true indeed, that where-ever the Bones on which any Part of our Body is sustained decline from a streight Line, the Force re quired in the Muscles to counteract the Gravi ty of that Part, is greater than otherwise neede

of

Cri

fift

Car

Cer

Ne

to i

Size

it is

galea,

⁽a) Cowper. Anat. of human Bodies, Explic. of Tab. 8.

needed to have been: But then this is effectually provided for in such Places, by the Number and Strength of the Muscles. As long therefore as we remain in the same Posture, a considerable Number of Muscles must be in a constant State of Contraction; which we know, both from Reason and Experience, must soon create an uneasy Sensation. This we call, being weary of one Posture: An Inconvenience that we should not have had in standing erect, if the Bearing of all the Bones to each other had been perpendicular; but is more than compensated by the Advantages above mentioned.

The human Skeleton is generally divided into the HEAD, the TRUNK, the SUPERIOR and

the INFERIOR EXTREMITIES.

h

nd

ar-

put

of to of

ire,

heir can

S O

pen

nuch

feve.

orm

ones

ined

e re

ravi

fe

ede

b. 8 A

Of the HEAD.

BY the HEAD is meant all that spheroidal Part which is placed above the first Bone of the Neck. It therefore comprehends the Cranium and Bones of the Face.

The Cranium *, Helmet, or Brain-case, consists of several Pieces; which form a vaulted Cavity, for lodging and defending the Brain, Cerebellum, with their Membranes, Vessels, and Nerves.

The Cavity of the Cranium is proportioned to its Contents. Hence such a Variety of its Size is observed in different Subjects; and hence it is neither so broad nor so deep at its Fore-

^{*} Κογχος, κύτος, κάδωα, σκαρίον, calva, çavaria, cerebra galea, theca & olla capitis, testa capitis, scutella capitis.

part, in which the anterior Lobes of the Brain are lodged, as it is behind, where the large posterior Lobes of the Brain, and the whole

Cerebellum, are contained.

The roundish Figure of the Scull, which makes it more capacious, and better able to defend its Contents from external Injuries, is chiefly owing to the equal Pressure of these contained Parts as they grow and increase before it is entirely offified.—It is to be observed, however, that in Adults the Sides of the Cranium are depressed below a spherical Surface by the strong temporal Muscles, whose Action hinders here the uniform Protrusion of the Bones, which is more equally performed in other Parts, where no fuch large Muscles are. In Children, whose Muscles have not acted much, and confequently have not had great Effects on the Bones, this Depression is not so remarkable: and therefore their Heads are rounder than in Adults. These natural Causes, differently difposed in different People, produce a great Variety in the Shapes of Sculls, which is still in creased by the different Management of the Heads of Children when very young: So that one may know a Turk's Scull by its globular Figure, a German's by its Breadth and Flatnes of the Occiput, Dutch and English by their of long Shapes, &c. (a). Two Advantages are reaped from this Flatness of the Sides of the Cranium, viz. the Enlargement of our Sphere com of Vision, and more advantageous Situation of OUI

t

in

shi

in

1.0.

ter

of

fere

be .

mo

Wh

her

ed .

s fu

er in he I

nore

from

our Ears, for receiving a greater Quantity of Sound, and for being less exposed to Injuries.

The external Surface of the upper Part of the Cranium is very smooth, and equal, being only covered with the Perioseum, (common to all the Bones, but, in the Scull, distinguished by the Name of Pericranium), the thin frontal and occipital Muscles, their tendinous Aponeurosis, and with the common Teguments of the Body; while the external Surface of its lower Part has numerous Risings, Depressions and Holes, which afford convenient Origin and Infertion to the Muscles that are connected to it, and allow safe Passage for the Vessels and Nerves

that run through and near it.

h

0

n-

re d,

n-

by

on

es, ts,

en,

onthe

le;

in dif.

Va-

In-

the

hat

ular

ness ob-

are

the

n of

OU

The internal Surface of the upper Part of the Scull is commonly smooth, except where the Vessels of the Dura Mater have made Furrows in it, while the Bones were foft.—Surgeons should be cautious when they trepan here, lest, in faving or raising the Bone where such Furrows are, they wound the Vessels of the Dura Mater.—In the upper Part of the internal Surface of several Sculls, there are likewise Pits of different Magnitudes and Figures, which feem to be formed by some Parts of the Brain being more luxuriant and prominent than others. Where these Pits are, the Scull is so much thinher than any where elie, that it is often renderd diaphanous, the two Tables being closely compacted without a Diploe; the Want of which s supplied by Vessels-going from the Dura Maer into a great many small Holes observable in he Pits. These Vessels are larger, and much nore conspicuous than any others that are sent

from the Dura Mater to the Scull; as evidently appears from the Drops of Blood they pour out, when the Scull is raifed from the Dura Mater in a recent Subject; and therefore they may furnish a sufficient Quantity of Liquors necessary to prevent the Brittleness of this thin Part.-The Knowledge of these Pits should teach Surgeons, to faw cautiously and slowly through the external Table of the Scull, when they are performing the Operation of the Trepan; fince, in a Patient whose Cranium has these Pits, the Dura Mater and Brain may be injured, before the Instrument has pierced near the ordinary Thickness of a Table of the Scull-The internal Base of the Scull is extremely unequal, for lodging the feveral Parts and Appendices of the Brain and Cerebellum, and allowing Passage and Defence to the Vessels and Nerves that go into, or come out from these Parts.

t

i

Sp

the

bei

col

of

For n t

(a)

The

The Bones of the Cranium are composed of two Tables, and intermediate Cancelli, commonly called their Diploe †. The external Table is thickest; the inner, from its Thinness and consequent Brittleness, has got the Name of Vitrea. Whence we may see the Reason of those mischievous Consequences, which so often attend a Collection of Matter in the Diploe, either from an external or internal Cause, before any Sign of such a Collection appears in the Teguments that cover the Part of the Scullection is ladged (a)

where it is lodged (a).

† Meditullium, commissura. (a) Bonet. Sepulchret. Anat. lib. 1. § 1. obs, 96.—103 The Diploe has much the same Texture and Uses in the Scull, as the Cancelli have in other Bones.

ly

ry

ir-

he

er-

ce,

he

ore

ary

er-

for

the

and

ato,

d of

om-

rnal

nels

ame

n of

often

, el-

efore

the

Scull

The

-103

The Diploe of several old Subjects is so obliterated, that scarce any Vestige of it can be seen; neither is it observable in some of the hard craggy Bones at the Base of the Scull. Hence an useful Caution to Surgeons who trust to the Bleeding, Want of Resistance, and Change of Sound, as certain Marks in the Operation of the Trepan, for knowing when their Instrument has sawed through the first Table, and reached the Diploe (a). In other People, the Diploe becomes of a monstrous Thickness, while the Tables of the Scull are thinner than Paper.

The Cranium consists of eight Bones, six of which are said to be proper, and the other two are reckoned common to it, and to the Face.—
The six proper are the Os frontis, two Ossa parietalia, two Ossa temporum, and the Os occipitis.—The common are the Os Ethmoides and Sphenoides.

The Os frontis forms the whole Fore-part of the Vault; the two Osfa parietalia form the upper and middle Part of it; the Osfa temporum compose the lower Part of the Sides; the Os occipitis makes the whole hinder Part, and some of the Base; the Os Ethmoides is placed in the Fore-part of the Base, and the Os Sphenoides is in the Middle of it.

These Bones are joined to each other by sive Sutures; the Names of which are the Coronal, Lambdoid, Sagittal, and two Squamous.

F₃ The

⁽a) Bartholin. Anat. reform. Lib. IV. Cap. 4.

The Coronal + Suture is extended over the Head, from within an Inch or so of the external Canthus of one Eye, to the like Distance from the other; which being near the Place where the Antients wore their Vitta, Corona, of Garlands, this Suture has hence got its Name.—Though the Indentations of this Suture are conspicuous in its upper Part, yet an Inch or more of its End on each Side has none of them:

for it is fquamous and smooth there.

The Lambdoidal † Suture begins fome Way below, and farther back than the Vertex of Crown of the Head, whence its two Legs are stretched obliquely downwards, and to each Side, in Form of the Greek Letter A, and are now generally said to extend themselves to the Base of the Scull; but formerly Anatomists (a) reckoned the proper Lambdoid Suture to terminate at the Squamous Sutures, and what is extended at an Angle down from that on each Side, where the Indentations are less conspicuous than in the upper Part of the Suture, they called Additamentum suture Lambdoidis *.

01

th th

Off the

er

otl.

Scu

(a)

(b) (c)

(d)

Hip

ift. 5

(c) 1

This Suture is sometimes very irregular, being made up of a great many small Sutures, which surround so many little Bones that are generally larger and more conspicuous on the external Surface of the Scull, than internally. These Bones are generally called Triquetra of Wormiana; but some other Name ought to be

[†] Στιφαναΐα, Arcualis, Puppis. ‡ Laudæ, Proræ, Hypsyloides.

⁽a) Vefal. Anat. Lib. 1. cap. 6.
Lambdoides harmonialis, Lambdoides inferior, occipitis or rona.

16

r.

Cd

ce

of

n.

re

1;

ay

of

are ide

OW

Bafe

rec-

nate

ided

ide,

than

Ad-

be

ures

t are the nally

ra of

to be

tis d

be given them, for they are not always of a triangular Figure; and older Anatomists (a) than Olaus Wormins (b) have described them. The specifick Virtue which these Bones were once thought to have in the Cure of the Epilepsy (c) is not now ascribed to them; and Anatomists generally agree, that their Formation is owing to a greater Number of Points than ordinary of Offification in the Skull, or to the ordinary Bones of the Cramium not extending their Offification far enough or foon enough; in which Case, the unossified Interstice between such Bones, begins a separate Ossification in one or more Points; from which the Offification is extended to form as many distinct Bones as there were Points, that are indented into the large ordinary Bones, and into each other.—Probably those Children who have a large Opening in this Place at their Birth, will have the largest Offa Triquetra. To confirm this Account of the Formation of these little Bones, we may emark, that fuch Bones are fometimes feen in other Sutures, as well as in the Lambdoid (d); nd they are sometimes in one Table of the scull, and not in the other (e).

The

(b) Musæum, lib. 3. cap. 26.

⁽a) Eustach. Ofsum Examen.—Baubin. Theat. Anat. lib.

⁽c) Baubin. & Paaw. ibid.—Bartholin. Anat. reform. lib. 4. ap. 5.—Hildan. Epistol. 65.

⁽d) See Examples in Vefal. lib. 1. cap. 6. fig. 4.—Paaw Hippocrat. de cap. Vuln. Bartholin. Hist. Anat. Cent. 1. ist. 51.—Raysch. Mus. Anat.

⁽c) Hunauld. in Mem. de l'Acad. des Sciences, 1730.

The Sagittal Suture * is placed longitudinally in the Middle of the upper Part of the Scull. and commonly terminates at the Middle of the Coronal, and of the Lambdoid Sutures; between which it is faid to be placed, as an Arrow is between the String and Bow .- However this Suture is frequently continued through the Middle of the Os frontis, down to the Root of the Nose; which, some (a) say, oftener happens in Women than Men; but others (b) alledge, that it is to be met with more frequently in Male Sculls, than in Female: Among the Sculls which I have feen thus divided, the Female are the most numerous. Several (c) have delineated and described the Sagittal Suture, some times dividing the occipital Bone as far down a the great Hole through which the Medulla Spinalis passes. This I never faw.

In some old Sculls that are in my Possession, there is scarce a Vestige of any of the three Sutures which I have now described. In other Heads, one or two of the Sutures only disappear; but I never could discover any Reason for thinking them disposed in such different Manners in Sculls of different Shapes, as some Anti-

he

7

u

OII

ht

on

):

hic e c

t As

(a) mon (b)

(e) V

Th

ents alledge they are (d).

(a) Riolan. Comment. de Offib. cap. 8. (b) Vefal. Lib. 1. cap. 6. et in Epitome.

(d) Hippocrat. de Vulner. Capitis, § 1 .- Galen. de Off

& de usu Part. lib. 9. cap. 17.

Pagoonone, δειασία, επιζευγνύσα, Instar virgae, nervali instar teli, instar veru, secundum capitis longitudinem prost pens, conjungens, columnalis, recta, acualis.

⁽c) Vefal. Lib. 1. cap. 5. fig. 3, 4. et in Text. cap. 6. Paaw in Celf. de re medic. cap. 1.— Laurent. Hist. Anat. 2. cap. 16.

e

n

e.

ulle

he

in

nat

ale

alls

are eli-

men as

Spi-

ion,

Su-

ther

isap.

n for

Man-

Anti

The

ervalis prote

0.6.

at.

e Off

The Squamous Agglutinations, or False Suures +, are one on each Side, a little above the Ear, of a semicircular Figure, formed by the overlopping (like one Scale upon another) of he upper Part of the temporal Bones on the ower Part of the Parietal, where, in both Bones, there are a great many small Risings nd Furrows, which are indented into each oher; though these Inequalities do not appear ill the Bones are separated. In some Sculls ineed the Indentations here are as conspicuous xternally as in other Sutures (a); and what is ommonly called the posterior Part of this squanous Suture, always has the evident ferrated form: and therefore is reckoned by some (b). distinct Suture, under the Name of Additaventum posterius sutura squamosa.—I have en two fourmous Sutures on the same Temple, ith a semicircular Piece of Bone between the Frontal: hem.

We ought here to remark, that the true namous Sort of Suture is not confined to the onjunction of the temporal and parietal Bones, at is made Use of to join all the Edges of the ones on which each temporal Muscle is placed): For the two Parts of the sphenoidal Suture hich are continued from the anterior End of the common squamous Suture just now describle, of which one runs perpendicularly downwards.

[†] Λιπιδοκιδή, προσπολλήματα, προταφίαι, Temporales, Corales, Mendofæ, Harmoniales, Commiffura in unguem.

⁽a) Columb. de te Anat. lib. 1. cap. 4.—Dionit, Anat. 3. monst. des Os.

⁽b) Albin. de Offib. § 54.

⁽c) Vesal. Anat. lib. 1. cap. 6.—Winslow, Mem. de l'Acad.
Sciences, 1720.

wards, and the other horizontally forwards, and also the lower Part of the coronal Suture already taken Notice of, may all be justly faid to pertain to the squamous Suture.—The Manner how I imagine this Sort of Suture is formed at these Places, is, That, by the Action of the strong temporal Muscles on one Side, and by the Pressure of the Brain on the other, the Bones are made fo thin, as to admit of no a ther Suture. The fuamous Form is also more convenient here; because such thin Edges of Bones, when accurately applied one to another, have scarce any rough Surface, to obstruct of hart the Muscle in its Contraction; which is still further provided for, by the Manner of laying these Edges on each other; for, in view ing their Outside, we see the temporal Bona covering the Sphenoidal and Parietal, and this last supporting the Sphenoidal, while both mount on the Frontal: From which Disposition it is evident, that while the temporal Muscle's contracting, which is the only Time it press strongly in its Motion on the Bones, its Fibre flide easily over the external Edges. Another Advantage still in this is, that all this bony Par is made stronger by the Bones thus supporting each other. tare continued from the

t

ſė

be

be do

the

wh

WC

or . baff

ure It

he !

ut t ine

Th

The Bones of the Scull are joined to thok of the Face, by Schyndelesis and Sutures.—The Schyndelesis is in the Partition of the Nose.— Schyndelesis is in the Partition of the Nose.— the Standard Francisco The Sutures said to be common to the Cranium Francisco and Face are five, viz. the Ethmoidal, Sphenoidal, Transverse, and two Zygomatic.—Parts how ever of these Sutures are at the Junction of on the a ifest

ly the Bones of the Scull.

The Ethmoidal and Sphenoidal Sutures furround the Bones of these Names; and in some Places help to make up other Sutures, particularly the Squamous and Transverse; and in other Parts there is but one Suture common to these two Bones.

he

by

the

0.

ore

of

ner,

Of

h is

· of

èw.

ones

this

both

tion

lei

effet

ibre

other

Pan rting

thole-The

-

enium benoi

how

Th

The Transverse Suture runs quite cross the Face, from the external Canthus of one Orbit to the same Place of the other, by sinking from the Canthus down the Out-side of the Orbit to its Bottom; then mounting upon the Inside, it is continued by the Root of the Nose down the internal Part of the other Orbit, and up again to the other Canthus. It may be here remarked, that there are some Interruptions of this Suture in the Course I have described; for the Bones are not contiguous every where, but are separated, to leave Holes and Apertures, to be mentioned hereafter.

The Zygomatic Sutures are one on each Side, being short, and santing from above obliquely downwards and backwards, to join a Process of the Cheek-bone to one of the temporal Bone, which advances towards the Face; so that the two Processes thus united, form a fort of Bridge, or Jugum, under which the temporal Muscle passes; on which account the Processes, and Suure joining them, have been called Zygomatic.

It must be observed, that the Indentations of he Sutures do not appear on the Inside of the Frankium, by much so strong as on the Out-side, ut the Bones seem almost joined in a streight line: nay, in some Sculls, the internal Surace is found entire, while the Sutures are majest without; which may possibly be owing to

the

the less Extent of the concave than of the con. vex Surface of the Cranium, whereby the Fibres of the internal Side would be stretched farther out at the Edges of the Bones, than the exterior ones, if they were not relifted. The Resistances are the Fibres of the opposite Bone, the Parts within the Scull, and the Diploe; of which the last being the weakest, the most advanced Fibres or Servæ run into it, and leave the contiguous Edges equal, and more ready to unite: Whereas the Serræ of the external Table have space enough for their Admission between the Fibres of the opposite Bone; and therefore remain of the indented Form, and are less is able to the Concretion, whereby the Suture are obliterated (a). By this Mechanism there is no Rifk of the sharp Points of the Bone growing inwards, fince the external Serra each of the conjoined Bones rest upon the in ternal smooth edg'd Table of the other; and external Forces applied to these Parts an strongly resisted, because the Sutures cannot yield, unless the serrated Edges of the on Bone, and the plain internal Plate of the other are broken (b).

The Advantages of the Sutures of the Crais um are these: 1. That this Capsula is more easily formed and extended into a spherical surre, than if it had been one continued Bong. That the Bones which are at some Distant from each other at Birth, might then yield and allow to the Head a Change of Shape, a commodated to the Passageit is engaged in

t

e

B

5.

rell

des

(a) Hunauld, Memoires de l'Acad. des Sciences, 1730 (b) Winslow, Memoires de l'Acad. des Sciences, 1720 he

he

ne,

ot

ad.

ave

y to

able

veen

fore

s li

tura

nifm,

3one

ræd

ne in

and

s are

anno

e on

othe

Cran

mon

cal f

Bon

istand

yiel

pe, a ged i Theno

1730

1720

Whence, in hard labour of Child-bed, the Bones of the Cranium, instead of being only brought into Contact, are sometimes made to mount one upon the other. 3. It is alledged, that, thro' the Sutures, there is a Transpiration of Steams from the Brain, which was the old Doctrine; or some Communication of the Vesfels without, and of those within the Scull, larger here than in any other Part of the Cranium, according to some Moderns; and therefore Cucupha, Fomentations, Cataplasms, cephalie Plaisters, Blisters are applied, and Issues are eroded, or cut in the Head, at those Places where the Sutures are longest in forming, and where the Connexion of the Bones is afterwards loofest, for the Cure of a Phrenitis, Mania, inveterate Head-ach, Epileply, Apoplexy, and other Diseases of the Head. The Favourers of the Doctrine of Transpiration, or Communication of Vessels at the Sutures, endeavour to support it by Observations of Persons subject to Head-achs which caused Death, from the Sutures being too closely united (a). 4. That the Dura Mater may be more firmly suspended by its Processes, which infinuate themselves into this Conjunction of the Bones; for doing this equally, and where the greatest Necessity of Adhesion is, the Sutures are disposed at nearly equal Distances, and the large Reservoires of Blood, the Simises, are under or near them. 5. That Fractures might be prevented from G reaching

⁽a) Columb. de re Anat. lib. 1. cap. 5. — Verduc. nouvelle Osteologie, chap. 14. — Dionis, Anat. 3. Demonstr. des Os.

reaching so far as they would in a continued bony Substance: 6. That the Connexion at the Sutures being capable of yielding, the Bones might be allowed to separate; which has given great Relief to Patients from the violent symptoms which they had before this Separation happened (a). And it seems reasonable to believe, that the opening of the Sutures was of great Benefit to several others who were rather judged to have been hurt by it (b). For we must think, that the Consequences of such a Force acting upon the Brain, as was capable of thrusting the Bones asunder, must have been fatal, unless it had been thus yielded to.

Having gone through the general Structure of the Cranium, I now proceed to examine each Bone of which that Brain-case consists, in the Order in which I first named them.

The OS FRONTIS + has its Name from its being the only Bone of that Part of the Face we call the Forehead, though it reaches a good deal further. It has some resemblance in Shape to the Shell of the Concha bivalvis, commonly called the Cockle; for the greatest Part of it is convex externally, and concave internally, with a serrated circular Edge; while the

(a) Ephemerid.) Germanic. Dec. 1. Ann. 4: & 5.06

† Μετώπη, Βρέγμα, Coronale, Inverecundum, Puppis, fet

Ins communis; Sincipitis.

⁽b) Ephemerid. Germ. Dec. 2. Ann. 9. Obs. 230. Ibid. Gent. 10. Obs. 31. — Vander Linden Medicin. Phys. cap. 8. art. 4. §. 16.—Hildan. Observ. Cent. 1. Obs. 1. Cent. 2. Obs. 7.—Bauhin. Theat. Anat. Lib. 3. cap. 6.—Pechlis. Observ. Lib. 2. Observ. 39.

smaller Part has Processes and Depressions, which

make it of an irregular Figure.

ied

at

nes

ven

np-

10n

be-

of

her

we

h a

e of

peen

ture

nine

, in

m its

Face

5.00q

hape

only

rt of

nter-

e the

naller

5. Ob.

. Ibid.

cap. 8.

Pechlin.

The external Surface of the Os frontis is smooth at its upper convex Part; but several Processes and Cavities are observable below: For, at each Angle of each Orbit, the Bone jutts out, to form four Processes, two internal, and as many external; which, from this Situation, may well enough be named angular. Between the internal and external angular Processes of each Side, an arched Ridge is extended, on which the Eye-brows are placed. --- Very little above the internal End of each of these superciliary Ridges, a Protuberance may be remarked, in most Sculls, where there are large Cavities, called Sinuses, within the Bone; of which hereafter. -Betwixt the internal augular Processes, a small Process rifes, which forms some Share of the Nose, and thence is named Nasal- Some observe a protuberant Part on the Edge of the Bone behind each external angular Process, which they call temporal Processes; but their are inconfiderable. - From the under Part of the superciliary Ridges, the frontal Bone runs a great Way backwards; which Parts may justly enough be called orbitar Processes. These, contrary to the rest of this Bone, are concave externally, for receiving the Globes of the Eyes, with their Muscles, Fat, &c.

In each of the orbitar Processes, behind the Middle of the supercitiary Ridges, a considerable Sinuosity is observed, where the Glandula innominata Galeni or lathrymalis is lodged.—Behind each internal angular Process, a small Pit may be remarked, where the cartilaginous Pul-

mand of G.2 and Time Short

ly of the Musculus obliques major of the Eye is fixed. — Betwixt the two orbitar Processes, there is a large Discontinuation of the Bone, into which the cribrisorm Part of the Os ethemoides is incased.—The frontal Bone frequently has little Caverns formed in it here where it is joined to the ethmoid Bone—Behind cach external angular Process, the Surface of the frontal Bone is considerably depressed where

part of the temporal Muscle is placed.

The Foramina, or Holes, observable on the external Surface of the frontal Bone, are three in each Side. — One in each fuperciliary Ridge, a little removed from its Middle towards the Nose: through which a Twig of the Oph. thalmic Branch of the fifth Pair of Nerves palfes out of the Orbit, with a small Artery from the internal Carotid, to be distributed to the Teguments and Muscles of the Forehead. These Vessels in some Sculls make Furrows in the Os frontis, especially in the Bones of Children, as has also been observed of another confiderable Vessel of this Bone near its Middle (a): and therefore we ought to beware of transverle Incisions on either Side of the Os frontis, which might either open these Vessels, while they are yet in Part within the Bone; for, in such a Case, it is difficult to stop an Hæmorrhagy, because the Adhesion of a Part of the Artery to the Bone, hinders its Contraction, and confequently Styptics can have little Effect; the Sides of the Furrow keep off compressing Substances from the Artery; and we would with to thus Cauteries

nc Bra

Vo

⁽a) Ruysch. Mus. Anat. Theca D. Reposit. 4. No. 3.

es,

ne,

eth-

nt-

ere

ind.

of

ere

the

arce

iary

ards

Oph-

pal-

rom

the

78 IN

Chil:

con-

: (a):

verle

which

y are

Case,

caule

o the

uent.

ances

thun

eries

3.

Cauteries or Escharotics, because they make the Bone carious; or might wound the Nerves, fo as to produce violent Symptoms.—But, to return to the superciliary Foranina, we must remark, that often, instead of a Hole, a Notch only is to be feen: Nay, in some Sculls, scarce a Vestige even of this is left; in others, both Hole and Notch are observable, when the Nerve and Artery run feparately. Frequently a Hole is found on one Side, and a Notch on the other: at other Times we fee two Holes; or there is a common Hole without, and two distinct Entries internally. The Reason of this Variety of a Hole, Notch, Depression, or Smoothness in the superciliary Ridge, is the different Length and Tension of the Nerves and Vessels; the thorter they are, the more they are limk into the Bone as it grows.—Near the Middle of the Infide of each Orbit, hard by, or in the transverse Suture, there is a small Hole for the Passage of the nafal Twig of the first Branch of the fifth Pair of Nerves, and of a Branch of the ophthalmic Artery. This Hole is fometimes entirey formed in the Os frontis; in other Sculls, the Sides of it are composed of this last Bone, and of the Os planum. It is commonly known by the Name of Orbitarium internum, though antein should be added, because of the next, which s commonly omitted.—This, which may be alled Orbitarium internum posterius, is such ano her as the former; only smaller, and about an nch deeper in the Orbit: Through it a finall branch of the ocular Artery passes to the Nofe.—Besides these six, there are a great Numer of small Holes observable on the outer Surface

face of this Bone, particularly in the two Protuberances above the Eye-brows. Most of these penetrate no further than the Sinuses, or than the Diploe, if the Sinuses are wanting; though sometimes I have seen this Bone so persorated by a vast Number of these small Holes, that, placed between the Eye and a clear Light, it appeared like a Sieve.—In the Orbit of the Generality of Skeletons, we may observe one, two, or more Holes, which allow a Passage to a Hog's Bristle through the Scull. The Place, Size and Number of these, are however uncertain: They generally serve for the Transmission of small Arteries or Nerves.

The internal Surface of the Os frontis is concave, except at the orbitar Processes, which are convex, to support the anterior Lobes of the Brain. This Surface is not fo smooth as the external; for the larger Branches of the Arteria of the Dura Mater make some Furrows in is Sides and back Parts. The Sinuofities from the huxuriant Rifings of the Brain, mentioned when describing the general Structure of the Cranium, are often very observable on its upper Part; and its lower and fore Parts are marked with the Contorsions of the anterior Lobes of the Brain.—Through the Middle of this internal Surface, where always in Children, and fome times in old People, the Bone is divided, either a Ridge stands out, to which the upper Edge of the Falx is fastened, or a Furrow runs, a which the upper Side of the superior longitud nal Sinus is lodged; on both these Accounts chi rurgical Authors justly discharge the Applicate on of the Trepan here. The Reason of the Difference

2126

Difference in Scalls, is alledged by some Authors to be this, That in thin Sculls the Ridge strengthens the Bones, and in thick ones there is no Occasion for it. To this Way of accounting for this Phænomenon, it may justly be objected, that generally very thick Sculls have a large Spine here, and frequently thin ones have only a Furrow. Perhaps this Variety may be owing to the different Times of a compleat Offification of those Parts in different Subjects : For if the two Sides of this Bone meet before they arrive at their utmost Extent of Growth. they unite very firmly, and all their Fibres endeavour to stretch themselves out where the least Resistance is, that is, between the Hemispheres of the Brain. To support this Reasoning, we may remark, that those Adults, whose frontal Bone is divided by the fagittal Suture, never have a Ridge in this Place.

Immediately at the Root of this Ridge or Furrow there is a small Hole, which sometimes pierces through the sirst Table, and, in other Sculls, opens into the superior Simus of the Ethmoid Bone within the Nose. In it a little Process of the Falx is lodged, and a small Artery, and sometimes a Vein, runs (a); and the superior longitudinal Simus begins here.—This Hole, however, is often not entirely proper to the Os frontis; for in several Sculls, the lower Part of it is formed in the upper Part of the Base of the Crista Galli, which is a Process of the Os Eth-

moides (b).

ro-

ele

nan

igh

ted

lat.

, it

Ge-

WO.

og's

and

mall

con-

the

e ex-

eries

n ils

n the

when

nium,

Part;

with f the

ternal

fomt.

either

Edge

ns, n

ts chi-

licati

f this

rence

The

⁽a) Morgagn. Adversar. 6. Animad. 31.

⁽b) Ingraff. Comment. in Galen. de Ossib. cap. 1. Comment. 8.

The Os frontis is composed of two Tables, and an intermediate Diploe, as the other Bones of the Cranium are, and, in a middle Degree of Thickness between the Os occipitis and parietalia, is pretty equally dense all through, except at the orbitar Processes; where, by the Action of the Eye on one Side, and Pressure of the Lobes of the Brain on the other, it is made extremely thin and diaphanous, and the Meditulian is entirely obliterated. Since in this Place there is so weak a Desence for the Brain, the Reason appears why Fencers esteem a Puth in

the Eye mortal (a).

The Diploe is also exhausted in that Part a bove the Eye-brows, where the two Tables of the Bone separate, by the external being protruded outwards, to form two large Cavilies, called Sinus fromates. These are divided by a middle perpendicular bony Partition. --- Their Capacities in the fame Subject are feldom equal; in some the Right, in others the Lest is largest .-And in different Bones their Size is as incomstant; nay, I have examined some, where they were entirely wanting; which oftener happens in fuch as have a flat Fore-head, and whose fagittal Suture is continued down to the Note, than in others (b).—In some Sculls, besides the large perpendicular Seption, there are feveral bony Pillars, or thort Partitions, found in each Sinus; in others thefe are wanting. For the most Part the Septum is entire; at other Times

F

(0)

(b) Fallop. Exposit. de Offibus; cap. 13.

⁽a) Ruysch. Observ. Anat. Chir. Observ. 54.—Diemertro eck. Anat. lib. 3. cap. 10.—Bonet. Sepulch. Anat. l.b. 4 § 3. Observ. 17.

iles,

gree

arte.

cept

the

ex-

Place

the

th in

rt a.

es of

pro-

Ities,

by a

Their

qual;

cit.-

ncon-

they

ppens

ose sa-

Nofe,

es the

everal

each

or the

Fimes

. 11

mertro-1.b. 4 it is discontinued, and the two Simuses commutationicate—When the Simuses are seen in such Sculls as have the frontal Bone divided by the sagittal Suture, the Partition dividing these Cavities, is evidently composed of two Plates, which easily separate.—Each Simus commonly opens by a roundish small Hole, at the inner and lower Part of the internal angular Processes, into a Simus formed in the Nose, at the upper and back Part of the Os unguis; near to which there are also some other small Simuses of this Bone (a), the greater Part of which open separately nearer he Septum narium, and often they terminate in the same common Canal with the large ones.

In a natural and found State, thele Cavities re of confiderable Advantage; for the Organ f Smelling being thus enlarged, the Effuvia f odorous Bodies more difficultly escape it; nd their Impressions being more mimerous, are herefore itronger, and affect the Organ more. These and the other Cavities which open into he Nofe, increase the Sound of our Voice, and ender it more melodious, by ferving as fo many aults to refound the Notes. Hence People laouring under a Coryza, or Stoppage of the lose from any other Cause, when they are by he Vulgar, though fallely, faid to speak through heir Nose, have such a disagreeable harth oice.—The Liquor separated in the Membrane f these Sinuses, drills down upon the Memrane of the Nose to keep it moist.

From the Description of these Simtses, it is sident, how useless, nay, how pernicious it

must

⁽a) Cowper in Drake's Anthropolog. Book 3. Chap. 10.

.....

must be, to apply a Trepan on this Part of the Scull; for this Instrument, instead of piercing into the Cavity of the Cranium, would reach me further than the Simifes; or, if the inner Table was perforated, any extravalated Blood that happened to be within the Scull, would not ke discharged outwardly, but would fall into the Simples, there to stagnate, corrupt, and stime late the sensible Membranes; from which allo there would be such a constant Flow of glain Mucus, as would retard, if not hinder a Cure and would make the Sore degenerate into a incurable Fiftula: Besides, as it would be a most impossible in this Case to prevent the Air passing through the Nose, from having con stant Access to the Dura Mater, or Brain; such a Corruption would be brought on these Parts as would be attended with great Danger. In ther, in Respiration, the Air rushing violents into these Cavities of the Os frontis, and pal fing through the external Orifice, whenever was not well covered and defended, would no only prevent the cloting up of the external 0 rifice, but might otherwise bring on bad Con sequences (a).—The Membrane lining the Sinufes is fo fensible, that Inflammations of must create violent Torture (b); and Worm or other Infects, crawling there, must give great Uncafinels (c). regal to pal off

0

u

Ö

he

if

ari

1.11

afa.

hd

roc

bme

Th

rt.

con

e G

ptilli

e D

es o

In a

oug

for

Observ. 10.

⁽a) Paaw, de Offbus Parf. 1. cap. 7.—Palfyne Anatos Chir. Traite 4. chap. 15. Nouvelle Offeologie, Partici chap. 3.

⁽c) Fernel. Partholog. lib. 5. cap. 7.—Barthelin. Epilo Medic. Cent. 2. Epilt. 74.—Hift. de l'Acad. des Scient 1708 & 1733.

no

ole

at

he

the

110

alfo

iry

an

al

Air,

còn-

fuch arts,

Fur-

pal

reri

d not

Cod

the

of

giv

naton

The upper circular Part of the Os frontis, is oined to the Offa parietalia, from one Temple to the other, by the coronal Suture. From the Termination of the coronal Suture to the exernal angular Processes, this Bone is connected o the Sphenoid by the Sphenoidal Suture. At he external Canthi of the Eyes, its angular Proeffes are joined by the transverse Suture to the Ma malarum, to which it adheres one Third own the Outside of the Orbits; whence to the lottom of these Cavities, and a little up on heir internal Sides, these Orbitar Processes are onnected to the sphenoidal Bone by that same uture. -In some few Sculls, however, a Difontinuation of these two Bones appears at the pper Part of the long Slit, near the Bottom f the Orbit. On the Inside of each Orbit, he orbitar Process is indented between the criiform Part of the ethnioid Bone, and the Os anum and unquis.—The transverse Suture afrwards joins the frontal Bone to the superior tial Processes of the Ossa maxillaria superiora, nd to the nafal Bones. And, lastly, its nafal ocess is connected to the nasal Lamella of the bmoid Bone. worth a reagh. I talen them

The frontal Bone serves to defend and support the anterior Lobes of the Brain. It forms considerable Part of the Cavities that contain a Globes of the Eyes, helps to make up the plum narium, Organ of Smelling, &c. From a Description of the several Parts, the other as of this Bone are evident.

In a ripe Child, the frontal Bone is divided rough the Middle; the superciliary Holes are t formed; often a small round Piece of each

orbitar

orbitar Process, behind the superciliary Ridge, is not offished, and there is no Sinus to be see

within its Substance.

Each of the two OSSA PARIETALIA, or Bones serving as Walls to the Encephalm, is an irregular Square; its upper and fore Sides being longer than the one behind or below. The inferior Side is a concave Arch; the middle Part receiving the upper round Part of the temporal Bone.—The Angle formed by this upper Side, and the fore One, is so extended, a

to have the Appearance of a Process.

The external Surface of each Os parietale in convex. Upon it, fomewhat below the middle Height of the Bone, there is a transverse and ed Ridge, of a whiter Colour generally tha any other Part of the Bone; from which, i Bones that have frong Prints of Muscles, w fee a great many converging Furrows, like many Radii drawn from a Circumference to wards a Center. From this Ridge of each Bon the temporal Muscle rises; and, by the Presim of its Fibres, occasions the Furrows just no mentioned. Helow thefe, we observe, ne the semcircular Edges, a great many Rising and Depressions, which are joined to like la qualities on the Infide of the temporal Bon to form the squamous Suture. The tempor Bone may therefore ferve here as a Buttres, prevent the lower Side of the Parietal fro starting outwards when its upper Part is pres or struck (a). Net

1

T

11

21

if flu

fo

th

tal

diff Th

lon

its nal

fma In

Hol

Kορυφής, Paria, syncipitis, verticis, arcualia, nervalia, gitationis, rationis, bregmatis, madefactionis.
11.(a) Hunaul! in Mem. de l'Acad. des Sciences, 1730.

Near the upper Sides of these Bones, towards the hind Part, is a small Hole in each, through which a Vein passes from the Teguments of the Head to the longitudinal Simis. Sometimes I have feen a Branch of the temporal Artery pass through this Hole, to be distributed to the upper Part of the Falx, and to the Dura Mater at its Sides, where it had frequent Anastomoles, with the Branches of the Arteries derived from the external Carotids, which commonly have the Name of the Arteries of the Dura Mater, and with the Branches of the internal Carotids which ferve the Falx.—In feveral Sculls, one of the Offa parietalia has not this Hole; in others, there are two in one Bone; and in some not one in either. Most frequently this Hole is thro' both Tables; at other Times the external Table is only perforated .-The Knowledge of the Course of these Veslels, may be of Use to Surgeons, when they make any Incision near this Part of the Head; lest, if the Vessels are rathly cut near the Hole, they farink within the Substance of the Bone, and fo cause an obstinate Hæmorrhagy, which neither Ligatures nor Medicines can stop.

25

2 1

Idle

rch har

, 11

to Bon

elliun

no

1162

ifing

e Ine Bone

por

efs, t

rese

Ne

12:13,

30.

On the inner concave Surface of the parietal Bones, we see a great many deep Furrows, disposed somewhat like the Branches of Trees: The Furrows are largest and deepest at the lower Edge of each Os parietale, especially near its anterior Angle, where sometimes a full Canal is formed: They afterwards divide into small Furrows, in their Progress upwards.—In some Sculls a large Furrow begins at the Hole near the upper Edge, and divides into

Branches,

Branches, which join with those which come upwards, shewing the Communications of the upper and lower Arteries of the Dura Mater.-In these Furrows we frequently see Passages in. to the Diploe; and fometimes I have observed Canals going off, which allowed a fmall Probe to pass some Inches into the bony Substance. Some (a) tell us, that they have observed these Canals piercing the Bone towards the Occiput. On the Infide of the upper Edge of the Offa parietalia, there is a large Sinuolity, frequently larger in the Bone of one Side than of the other, where the upper Part of the Falx is fastened, and the superior longitudinal Sinus is lodged.—Generally Part of the lateral Similes makes a Depression near the Angle, formed by the lower and posterior Sides of these Bones; and the Pits made by the prominent Parts of the Brain are to be feen in no Part of the Scull more frequent, or more confiderable, than in the internal Surface of the parietal Bones.

01

11

th

ne

Cre

and

net

Ope

ent

fluo

fom

ing, felt

nous

parie Th

Chile

tielf

may

he O

sitre

elt t

But if

blerv

· Palp

Chap.

The Osa parietalia are amongst the thinnest Bones of the Cranium; but enjoy the general Structure of two Tables and Diploe the completest, and are the most equal and smooth.

These Bones are joined at their Fore-side to the Os frontis by the coronal Suture; at their long inferior Angles, to the fphenoid Bone, by Part of the Suture of this Name; at their lower Edge, to the Ossa temporum, by the squamous Suture and its posterior Additamentum; behind, to the Ossaccipitis, or Ossa triquetra, by the

⁽a) Comper, Anatom. Explic. of XC. Tab. Fig. 1.

judge

the lambdoid Suture; and above, to one ano-

ther, by the fagittal Suture.

They have no particular Uses besides those mentioned in the Description of their several Parts, except what are included in the Account of the general Structure of the Cranium.

In a Child born at the full-Time, none of the Sides of this Bone are completed; and there never is a Hole in the officed Part of it

near to the fagittal Suture.

.

S

3

y

3;

of 111

in

eft

al

m.

to eir

by

·W(

112.

111;

by

the

The large unoffified ligamentous Part of the Cranium observable between the parietal Bones, and the Middle of the divided Os frontis of new-born Children, called by the Vulgar the Open of the Head, was imagined by the Antients to serve for the Evacuation of the superfluous Moisture of the Brain; and therefore they named it Bregma *, or the Fountain; sometimes adding the Epithet pulfatilis, or beating, on account of the Pulfation of the Brain. felt through this flexible Ligamento-cartilaginous Substance. Hence very frequently the parietal Bones are called Offa Bregmatis.

The upper middle Part of the Head of a Child, in a natural Birth, being what prefents. tielf first at the Os uteri (a), an Accoucheur may reach the Bregma with his Finger, when he Os uteri is a little opened. If the Bregma siftretched, and the Pullation of the Brain is. elt through it, the Child is certainly alive: But if it is shrivelled and flaccid, without any bservable Pulsation in it, there is Reason to H 2

^{*} Palpitans vertex, foliolum, folium, triangularis lacuna. (a) Burton's Midwifery, § 51 .- Smellie's Midwifery, Book . Chap. 1. \$ 5.

judge the Child to be very weak, or dead. Those who practise Midwifery should therefore examine the State of the Bregma accurately.

All the Bregma is generally offified before feven Years of Age. Several Authors (a) fay, they have observed it unossified in Adults; and Physicians, who order the Application of Medicines at the Meeting of the coronal and fagital Sutures, feem yet to think that a Derivation of noxious Humours from the Encephalon is more easily procured at this Part than any other of the Scull; and that Medicines have a greater Essect here, than estewhere, in the in-

t

11

31

CO

ar

ar

fee

an

wa

fin

the

1710

Hei

Its 1

tem

Part

The

oblo

is co

cont

tia, ju

ternal Diforders of the Head.

OSSA TEMPORUM ||, fo named, fay Authors, from the Hair's first becoming gray on the Temples, and thus discovering Peoples Ages, are each of them equal and fmooth a bove, with a very thin semicircular Edge; which, from the Manner of its Connexion with the neighbouring Bones, is distinguished by the Name of Os squamofum.—Behind this, the upper Part of the temporal Bone is thicker, and more unequal, and is sometimes described as a distinct Part, under the Name of Pars manmil laris (b).—Towards the Base of the Scull, the temporal Bone appears very irregular and une qual; and this Part, instead of being broad, and placed perpendicularly, as the others are, is contracted into an oblong very hard Substance, extended

(b) Albin. de Offib. § 26.

⁽a) Bartholin. Anat. Reform. lib. 4. cap. 6.——Diemerbroeck, Anat lib. 9. cap. 6.——Kerkring. Osteogen. cap. 1.

| Κρόταρῶν, κοροῶν, κοβρῶν, λεπιδοειδῆ, πολυειδή, λιθοική temporalia, lapidosa, mendosa, dura, arcualia, tympasum, armalia, saxea, parietalia.

extended horizontally forwards and inwards; which in its Progress becomes smaller, and is

commonly called Os petrofum.

e

d

1.

į.

is

0-

a n.

ay

ay

les

are;

ith

the

up.

and

15 1

mil-

the

me-

oad,

e, 18

nce,

aded

p. 1.

m, ar

Three external Processes of each temporal Bone are generally described.—The first placed at the lower and hind Part of the Bone, from its Resemblance to a Nipple, is called Mastoides, or Mammillaris. It is not folid, but within is composed of Cancelli, or small Cells, which have a Communication with the large Cavity of the Ear, the Drum; and therefore Sounds, being multiplied in this vaulted Labyrinth, are increafed, before they are applied to the immediate Organ of Hearing. Into the mastoid Process, the Sternomastoideus Muscle is inserted; and to its back Part, where the Surface is rough, the Trachelomastoideus, and part of the Splenius are fixed. - About an Inch farther forward, the second Process begins to rise out from the Bone; and having its Origin continued obliquely downwards and forwards for fome Way, it becomes smaller, and is stretched forwards to join with the Os malæ; they together forming the bony jugum, under which the temporal Muscle passes. Hence this Process has been named Zygomatic. + Its upper Edge has the strong Aponeurosis of the temporal Muscle fixed into it; and its lower Part gives rife to a Share of the Maffeter. — The Fore-part of the Base of this Process is an oblong Tubercle, which in a recent Subject is covered with a smooth polished Cartilage, continued from that which lines the Cavity im-H 3 mediately

[†] Rayagos, Paris, ansæ ossium temporum, ossa arcualia, pa-

2

le

tī

CE

M

fu

m Ja

the

tila

cei

par cel

fide

a T

the

Sim

Side

thef

ral S

ilag

hrm Th

lide (

each

betwe

s the

which

(2)

mediately behind this Tubercle. - From the under craggy Part of the Os temporum, the third Process stands out obliquely forwards. The Shape of it is generally faid to refemble the ancient Stylus scriptorius; and therefore it is called the Styloid Process *. Some Authors (a) however contend, that it ought to be named Steloid, from its being more like to a Pillar. Several Muscles have their Origin from this Process, and borrow one half of their name from it; as Stylo-glossis, Stylo-byoideus, Stylopharyngeus; to it a Ligament of the Os hyoides is fometimes fixed; and another is extended from it to the infide of the Angle of the lower Jaw. This Process is often even in Adults not entirely offified, but is ligamentous at its Root, and fometimes is composed of two or three distinct Pieces. - Round the Root of it, especially at the Fore-part, there is a remarkable Rising of the Os petrosium, which some have esteemed a Process; and, from the Appearance it makes with the Styliforan, have named it Vaginalis. - Others again have, under the Name of Auditory Process, reckoned among the external Procelles that femicircular Ridge, which, running between the Root of the mastoid and Zygomatic Processes, forms the under-Part of the external Meatus auditorius.

The Sinuolities or Depressions on the external Surface of each Os temporum are these:

— A long Fossa at the inner and back Part of the

* Τρατοείδη, βελονοείδη, πλημπρον, Os calaminum, fagittale, clavale, acuale, calcar capitis.

(a) Galen, de usu Part. lib. 2. cap. 4. - Fallop. Obsers.

e

.

e

it

1) d

r.

18

e

0-

es

d

V.

ts

ES

or

it,

k-

ne

p-

ve

ler

ng

ge,

120-

er-

er-

le: of

the

ttale,

ferr.

the Root of the mammary Process, where the posterior Head of the digastric Muscle has its Origin. — Immediately before the Root of the zygomatic Process, a considerable Hollow is left, for lodging the crotaphite Muscle. - Between the zygomatic, auditory and vaginal Procelles, a large Cavity is formed; through the Middle of which, from Top to Bottom, a Fiffure is observable, into which Part of the Ligament that fecures the Articulation of the lower law with this Bone is fixed. The Fore-part of the Cavity being lined with the fame Cartilage which covers the Tubercle before it, receives the Condyle of the Jaw; and in the Backpart a small Share of the parotid Gland, and a cellular fatty Substance are lodged. - At the Infide of the Root of the flyloid Apophyse, there is a Thimble-like Cavity, where the beginning of the internal jugular Vein, or End of the lateral Simus is lodged—And as the Sinules of the two Sides are frequently of unequal Size; so one of these Cavities is as often larger than the other (a). -Round the external Meatus auditorius, feveral Sinuofities are formed for receiving the Carilages and Ligaments of the Ear, and for their firm Adhesion.

The Holes that commonly appear on the Outlide of each of these Bones, and are proper to each of them, are five. — The first, situated petween the zygomatic, and mastoid Processes, s the Orifice of a large Funnel-like Canal, which leads to the Organ of Hearing; there-

fore.

⁽a) Hunauld. in Mem. de l'Acad. des Sciences, 1730.

2

0

1

0

to

ir

B

w

er

al.

un

mo Fu

Bo

the Bra

a l

run

than

ing

oth

Rid

ral

fore is called Meatus auditorius externust.-The second, the Aquaduct of Fallopius, or Paffage for the Portio dura of the auditory Nerve. is between the mastoid and styloid Processes. -Some way before; and to the Infide of the styloid Process, is the third Hole, the Canal from which runs first upwards, then forwards, and receives into it the internal carotid Ar. tery, and the Beginning of the intercostal Nerve; where this Canal is about to make the Turn forwards, one, or sometimes two very small Holes go off towards the Cavity of the Ear call. ed Tympanum; through these Valsatva (a) affirms the proper Artery or Arteries of that Cavity are fent. On the Anterior Edge of this Bone, near the former, a fourth Hole is observable, being the Orifice of a Canal, which runs outwards and backwards, in a horizontal Direction, till it terminates in the Tympanum. This, in the recent Subject, is continued for ward and inward, from the Parts which I mentioned just now as its Orifice in the Skeleton, to the Side of the Nostrils; being partly cartilaginous, and partly ligamentous. The whole Canal is named, Iter a palato ad curem, or Eustaebian Tube. — On the external Side of the bony Part of this Canal, and a-top of the Chink in the Cavity that receives the Condyl of the lower law, is the Course of the little Nerve faid commonly to be reflected from the lingual Branch of the fifth Pair, till it enters the Tympanum, to run across this Cavity, and . [0]

[†] Πίρος της αχολς, οπή τῶν ἄτῶν. Fenestra aurium (a) De Aure humana, cap. 2. Ş. 22 et Tab. 7. Fig. L

to have the Name of Chorda Tympani-The fifth Hole is very uncertain, appearing sometimes behind the mastoid Process; sometimes it is common to the temporal and occipital Bones; and in feveral Sculls there is no fuch Hole. The Use of it, when found, is for the Transmission of a Vein from the external Teguments to the lateral Sinus: But, in some Subjects, a Branch of the occipital Artery passes through this Hole, to ferve the back Part of the Dura Mater; in others, I have feen two or three fuch Holes: But they are oftener wanting than found. we may, once for all, in general remark, That the Largeness, Number, Situation and Existence of all fuch Holes, that for the most part allow only a Passage for Veins from without to the internal Receptacles, are very uncertain.

e

s,

r.

al

e

Ill

11.

1)

at

of is

ch

tal

111.

01-

11.

011,

ti-

ole

01

ile

the

dyle ttle

the

ters

and

10

. I.

The internal Surface of the Offa temporum is unequal; the upper circular Edge of the squamous Part having numerous small Ridges and Furrows for its Conjunction with the parietal Bones; and the rest of it is irregularly marked with the Convolutions of the middle Part of the Brain, and with Furrows made by the Branches of the Arteries of the Dura Mater.

From the under Part of this internal Surface, a large transverse hard craggy Protuberance runs horizontally inwards and forwards, with a sharp Edge above, and two states Sides, one facing obliquely forwards and outwards, and the other as much backwards and inwards. To the Ridge between these two Sides, the large lateral Process of the Dura Mater is fixed.

Sometimes

Sometimes a small Bone, a-kin to the Sesamoid, is found between the small End of this petrous Process and the sphenoid Bone (a).

Towards the back Part of the Inside of the Os temporum, a large deep Fossa is conspicuous, where the lateral Simus lies; and frequently on the Top of the petrous Ridge, a Furrow may be observed, where a small Sinus is situated.

ti

B

n

th

pi

tic

Bo

15

is

(0)

ant

111

ma

gall

all

g00

per

here

con

ton.

The internal proper Foramina of each of these Bones are, first, the internal Meatus au. ditorius in the polterior plain Side of the petrous Process. This Hole soon divides into two; one of which is the Beginning of the Aquadust of Fallopius; the other ends in feveral very finall Canals (b) that allow a Passage to the Branches of the Portio mollis of the seventh Pair of Nerves, into the Vestibule and Cochles. Through it also an Artery is sent, to be distributed to the Organ of Hearing. The Jecond Hole, which is on the anterior plain Side of the craggy Process, gives Passage to a Branch of the lateral Artery of the Dura Mater, which joins the Portio dura of the auditory Nerve, while it is in the Aquaduct (c). Frequently, however, a Branch of the Portio dura, accompanied with a Branch of that Artery which went into the Meatus, auditorius along with the acoustick Nerve, enters into the Cranium here, to be distributed to the fost Parts, at the Side of the Sella Turcica (d).—Near to this second Hole

⁽a) Riolan, Comment de Ossib. cap. 32.---Winslow, Exposition Anatomique de corps humain, Trait. des Os Secs. \$. 266.

⁽b) Valsalv. De aure humana, cap. 3. § 11.

⁽c) Fallop, Obs. Anatom.

⁽d) Valfalv. De aure, cap. 3. § 10..

Hole, there are often several small ones, where Arteries or Twigs of Nerves enter. On the posserior Surface, near the auditory Foremen, other small Holes may be taken Notice of: They seem only to convey Vessels to the Substance of the Bone.—The Passage of the cutaneous Vein into the lateral Simus, or of a Branch of the occipital Artery, is seen about the Middle of the large Fossa for that Simus; and the Orifice of the Canal of the carotid Artery, is evident at the under Part of the Point of the petrous Process.

Besides these proper Holes of the temporal Bones which appear on their external and internal Surfaces, there are two others in each Side that are common to this Bone, and to the occipital and sphenoidal Bones; which shall be mentioned afterwards in the Description of these

Bones.

e

LY

of

11.

e-0;

ict.

ry

18

th

ri.

nd

of

ch

ch

e,

y,

m-

ch

he

e,

de

nd

ole

Ex-

ecs,

The upper round Part of the squamous Bones is thin, but equal; while the low petrous Part is thick and strong, but irregular and unequal, having the Distinction of Tables and Diploe confounded, with several Cavities, Processes, and Bones within its Substance, which are Parts of the Organ of Hearing. That a clear Idea may be had of this beautiful; but intricate Organ, Anatomists generally chuse to demonstrate all its Parts together. I think the Method good; and therefore, since it would be improper to insert a compleat Treatise on the Ear here, shall omit the Description of the Parts contained within the Os petrosium of the Skeleton.

The temporal Bones are joined above to the parietal Bones by the squamous Sucures, and their posterior Additamenta: Before, to the sphenoid Bone by the Suture of that Name to the Cheek Bones by the zygomatic Sutures Behind, to the occipital Bone, by the Lamidout Sature and its Additamenta; and they are articlated with the lower Jaw, in the Manner which shall be described when this Bone is examined.

The Purposes which these two Bones serve, are easily collected, from the general Use of the Cranium, and from what has been said in

f

f

li

ot

in

W

Sie In

Su

in

Co

nue

No

the

ces.

the Description of their feveral Parts.

In an Infant, a small Fissure is to be observed between the thin upper Part, and the lower craggy Part of each of these Bones; which points out the recent Union of these Parts .-Neither mastoid nor styloid Processes are yet to be feen—Instead of a bony Funnel-like external Meatus auditorius, there is only a smooth bony Ring, within which the Membrane of the Drum is fastened.—At the Entry of the Es Itachian Tube, the Side of the Tympanum is not completed.—A little more outward than the internal auditory Canal, there is a deep Pit, o ver the upper Part of whose Orifice the interior semicircular Canal of the Ear is stretched; and some Way below this, the posterior semicircu lar Canal also appears manifestly.

OS OCCIPITIS*, fo called from its stuation, is convex on the Outside, and concart internally. Its Figure is an irregular Square or rather Rhomboid; of which the Angle about

^{*} Ivov, Basilare, Proræ, Memoriæ, Pixidis, Fibrosum, Mavosum, Lambde.

e

ve,

of

111

1977

nich

S.-

t to

ter-

ooth

the

Eu.

s not

1 the

t, 0

erio

and

ircu

its S.

ncave

quare

about

m, No

Angles are more finished, but obtuse; and the lower one is stretched forward in Form of a Wedge, and thence is called by some the cuneiform Process.—If one would, however, be very nice in observing the several Turns which the Edges of the Os occipitis make, sive or seven Sides, and as many Angles of this Bone, might be described.

The external Surface is convex, except at the cuneiform Apophyse, where it is flatted. At the Base of this triangular Process, on each Side of the great Hole, but more advanced forwards than the Middle of it, the large oblong Protuberances, named the Condyles, appear, to ferve for the Articulation of this Bone with the first Vertebra of the Neck. The smooth Surface of each of these Condyloid Processes is longest from behind forwards, where, by their oblique Situation, they come much nearer to each other than they are at their Back-part. inner Sides are lower than the external, by which they are prevented from sliding to either Side out of the Cavities of the first Vertebra (a). In some Subjects each of these plain smooth Surfaces feems to be divided by a small Rising in its Middle; and the lower Edge of each Condyle, next the great Foramen, is discontinued about the Middle, by an interveening Notch: whence fome (b) alledge, that each of these Apophyses is made up of two Protuberances.—Round their Root a small Depression and fpongy

⁽a) Galen. de usu Part. lib. 12. cap. 7.

⁽³⁾ Diemerbroeck, Anat. lib. 9. cap. 6.

fpongy Roughness is observable, where the Ligaments for furrounding and fecuring their Articulations adhere.-Though the Motion of the Head is performed on the Condyles, yet the Center of Gravity of that Globe does not fall between them, but is a good Way further forward: from which Mechanism it is evident, that the Muscles which pull the Head back, must be in a constant State of Contraction: which is stronger than the natural Contraction of the proper Flexors, else the Head would always fall forwards, as it does when a Man is afleep, or labours under a Palfy, as well as in Infants, where the Weight of the Head far exceeds the proportional Strength of these Muscles. This feeming difadvantageous Situation of the Condyles is however of good Use to us, by allowing fufficient Space for the Cavities of the Mouth and Fauces, and for lodging a fufficient Number of Muscles, which commonly ferve for other Uses; but may at Pleasure be directed to act on the Head, and then have an advantageous Lever to act with, fo as to be able to fustain a confiderable Weight appended, or other Force applied, to pull the Head back.

Somewhat more externally than the Condyles, there is a small Rising and semilunated Hollow in each Side, which make up Part of the Holes, common to the occipital and petrous Bones. Immediately behind this, on each Side, a scabrons Ridge is extended from the Middle of the Condyle, towards the Root of the Mastoid Process. Into this Ridge the Musculus lateralis, commonly ascribed to Fallopius, is inserted.—About the Middle of the external convex Surface, a large

Arch

-

e

le.

r-

1;

n

11.

1.

in

X.

ıf-

on

ıs,

of

ffi-

ly

be

an

ole

01

les,

OW

es,

m-

OUS

011-

ess.

on-

the

rge

rch

Arch runs cross the Bone; from the upper lateral Parts of which the occipital Muscles have their Rife; to its Middle the Trapezii are attached; and half Way between this and the great Hole, a lesser Arch is extended.—In the Hollows between the Middle of these Arches. the Complexi are inferted; and in the Depressions more external and further forward than these, the Splenii are inserted.—Between the Middle of the lesser Arch and the great Hole, the little hollow Marks of the recti minores appear; and on each Side of these the fleshy Insertions of the obliqui superiores and recti majores make Depressions.—Through the Middle of the two Arches a small tharp Spine is placed, which ferves as fome Sort of Partition between the Muscles of different Sides, or rather is owing to the Action of the Muscles depressing the Bone on each Side of it, while this Part is free from their Compression.—These Prints of the Muscles on this Bone are very Arong and plain. in some Subjects, but are not so distinct in others.—All round the great Foramen the Edges are unequal, for the firmer Adhesion of the frong circular Ligament which goes thence to the first Vertebra .- One End of each lateral or moderator Ligament of the Head, is fixed to a rough Surface at the Fore-part of each Condyle, and the perpendicular one is connected to a rough Part of the Edge of the great Hole between the two Condyles. Immediately before the Condyles, two little Depressions are made in the external Surface of the cuneiform Process, for the Infertion of the recti anteriores minores Muscles, which are unjustly ascribed to Cowper: And still. further

further forward, nearer the sphenoid Bone, are two other fuch Depressions, for the Reception of the recti anteriores majores.—When we confider the Size of the Prints of Muscles on the occipital Bone, before and behind its Condyles. and, at the same Time, compare their Distances from these Centers of Motion of the Head, we must see how much stronger the Muscles are which pull the Head backwards, than those are which bend it forward; and how much greater Force the former acquire by the long Lever they act with, than the latter which are infert. ed so near the Condyles. This great Force in the extensor Muscles is altogether necessary, that they might not only keep the Head from falling forward in an erect Posture, but that they might support it when we bow forward in the most necessary Offices of social Life, when the Weight of the Head comes to act at right Angles on the Vertebræ of the Neck, and obtains a long Lever to act with.

fee two Ridges; one standing perpendicular, the other running horizontally across the sirst. The upper Part of the perpendicular Limb of the Cross, to which the Falx is fixed, is hollowed in the Middle, or often in one Side, for the Reception of the superior longitudinal Sinus, and the lower Part of it has the small or third Process of the Dura Mater sastened to it, and is sometimes hollowed by the occipital Sinus. Each Side of the horizontal Limb is made hollow by the lateral Sinuses inclosed in the transverse Process of the Dura Mater; the Fossa in the right Side being generally a Continuation

are

lon

on.

the

les,

ces

We

are

are

ter

ver

ert-

e in

ry,

om.

hat

d in

hen

ght

ob-

we

ılar,

irst.

to of

OW-

the

mus,

hird

and

inus.

holranf-

tion

of

of the one made by the longitudinal Sinus in: the perpendicular Limb, and therefore is larger than the left one (a).—Round the Middle of the Cross there are four large Depressions separated by its Limbs; the two upper ones being formed by the Back-part of the Brain, and the two lower ones by the Cerebellum. Farther forward than the last mentioned Depressions, is the lower Part of the Fossa for the lateral Sinus on each Side. The inner Surface of the cuneiform Apophyse is made concave for the Reception of the Medulla oblongata, and of the Basilar Artery.—A Furrow is made on each Side, near the Edges of this Process, by a Simus of the Dura Mater, which empties itself into the lateral Simis (b).

The Holes of this Bone are commonly five proper, and two common to it and to the tentporal Bones.—The first of the proper Holes, called Foramen magnum * from its Size, is immediately behind the Wedge-like Process, and allows a Passage to the Medulla oblongata, Nervi accessorii, to the vertebral Arteries, fometimes to the vertebral Veins.—At each Side of this great Hole, near its Fore-part, and immediately above the Condyles, we always find a Hole; fometimes two, which foon unite again into one that opens externally; thro' thefe the ninth Pair of Nerves go out of the Scull .-The fourth and fifth Holes pierce from behind the Condyle of each Side into the Fossie of the lateral Sinuses; they serve for the Passage of

⁽a) Morgagn. Advers. Anat. 6. Animad. 1.

⁽b) Albin. de Ossib. § 65. Rachitidis, Medullæ spinalis.

the cervical Veins to these Simuses. Often one of these Holes is wanting, sometimes both, when the Veins pass thro' the great Foramen. Besides these five, we frequently meet with a ther Holes near the Edges of this Bone, for the Transmission of Veins; but their Number and Diameter are very uncertain.—The two common Foramina are the large irregular Holes, one in each Side, between the Sides of the cuneiform Process, and the Edges of the petrous Bones. In a recent Subject, a strong Membrane runs cross from one Side to the other of each of these Holes; in some Heads I have seen this Membrane offified, or a bony Partition dividing each Hole; and, in the greater Number of a dult Sculls, there is a finall sharp-pointed Pro cess stands out from the Os petrofum, and a more obtuse Rising in the occipital Bone, between which the Partition is stretched. Behind this Partition, where the largest Space is left, the lateral Simus has its Passage; and before it the eight Pair of Nerves and Accessorius make their Exit out of the Scull; and some Authors fay, an Artery passes through this Hole, to be bestowed on the Dura Mater.

The occipital Bone is among the thickest of the Cranium, though unequally so; for it is stronger above, where it has no other Desence than the common Teguments, than it is below, where, being pressed by the Lobes of the Brain and Cerebellum on one Side, and by the Action of the Muscles on the other, it is so very thin, as to be diaphanous in many Sculls: But then these Muscles ward off Injuries, and the Ridge and Spines, which are frequent here, make it

fuffici-

Bo

Pa

ot.

gre

the

fufficiently strong to resist ordinary Forces. The Tables and Diploe are tolerably distinct in this Bone, except where it is so thin as to be-

come diaphanous.

one

oth,

1.-

0

the

and

om.

les,

cu-

rous

ane

ach

this ling

f a.

Pro.

nd a

hind

left.

e it

nake

hors

o be

At of

it is

ence

elow,

Brain Ction

thin,

then

idges

ke it

The occipital Bone is joined above to the Offa parietalia and Triquetra when present, by the Lambdoid Suture; -- laterally to the temporal Bones, by the Additamenta of the Lambdoid Suture; below to the sphenoid Bone, by the End of its cuneiform Process, in the same Way that Epiphyses and their Bones are joined: For. in Children, a ligamentous Cartilage is interposed between the occipital and sphenoid Bones, which gradually turns thinner, as each of the Bones advances, till their Fibres at last run into each other; and, about fixteen or eighteen Years of Age, the Union of these two Bones becomes fo intimate, that a Separation cannot be made without Violence.—The Os occipitis is joined by a double Articulation to the first Vertebra of the Neck, each Condyle being received into a superior oblique Process of that Vertebra. What Motion is allowed here, we shall consider afterwards, where the Vertebræ are described.

The Uses of this Bone appear from the preceeding Description, and therefore need not be

repeated.

An Infant born at the full Time, has this Bone divided, by unoffified Cartilages, into four Parts.—The first of these is larger than the other three, is of a triangular Shape, and constitutes all the Part of the Bone above the great Foramen. Generally Fishers appear in the upper Part and Sides of this triangular Bone, when

when all the Cartilage is separated by Maceration; and sometimes little distinct Bones are seen towards the Edges of it.—The second and third Pieces of this Bone are exactly alike, and situated on each Side of the great Foramen; from which very near the whole Condyles are produced; and they are extended forwards almost to the Fore-part of the Hole for the ninth Pair of Nerves.—The fourth Piece is the cuneiform Process, which forms a small Share of the great Hole, and of these for the ninth Pair of Nerves, and of the Condyles: Betwixt it and the sphenoid Bone, a Cartilage is interposed.

Of the eight Bones which belong to the Cranium, there are only two which are not yet defcribed, viz. the Ethnoid and Sphenoid. These we already mentioned, in Complaifance to the Generality of Writers on this Subject, as Bones common to the Granium and Face, because the enter into the Composition of both: But the same Reason might equally be used for calling the frontal Bone a common one too. I shall, however, pass any idle Dispute about the Propriety of ranging them, and proceed to examine the Structure of the Bones themselves.

OS ETHMOIDES*, or the Sieve-like Bone, has got its Name from the great Number of small Holes with which that Part of it first taken Notice of is pierced. When this Bone is entire, the Figure of it is not easily described; but, by a Detail of its several Parts some Idea may be afforded of the whole; and therefore I shall distinguish it into the Cribit form

[•] Cribriforme, σπογραθλές, Spongiforme.

form Lamella with its Process, the Nasal Lamel-

la. Cellulæ, and Offa spongiofa.

7-

ire

nd

nd

71;

are

al.

nth

cu-

e of

air

t it

fed.

Cra-

de.

hefe

the

ones

thev

the

lling

hall,

Pro.

exa-

S.

-like

Vum

of it

1 this

v de

Parts,

: and

Cribit

form

The thin horizontal Lamella, is all (except its Back-part) pierced obliquely by a great Number of small Holes, through which the Filaments of the olfactory Nerves pass. In a recent Subject, these Holes are so closely lined by the Dura Mater, that they are much less conspicuous than in the Skeleton. From the Middle of the internal Side of this Plate, a thick Procels rifes upwards, and, being highest at the Fore-part, gradually becomes lower, as it is extended backwards. From some Resemblance which this Process was imagined to have to a Cock's Comb, it has been called Crista galli +. The Falx is connected to its Ridge, and to the unperforated Part of the cribriform Plate.-When the Crista is broke, its Base is sometimes found to be hollow, with its Cavity opening into the Nose (a),—Immediately before the highest Part of this Process, is the blind Hole of the frontal Bone, which, as was formerly emarked, is often in a good Measure formed by a Notch in the Fore-part of the Root of the Crista.

From the Middle of the outer Surface of the Cribriform Lamella, a thin folid Plate is extended downwards and forwards, having the same common Base with the Crista galli. Generally it is not exactly perpendicular, but is inclined to me Side or other, and therefore divides the Caity of the Nose unequally. Its Inclination to

one

[†] Verruea prædura, septum ossis spongiosi. (4) Palfyn. Auat. Chir. Tr. 4. Chap. 15.

one Side, and Flexure in the Middle, is some times so great, that it fills up a large Share of one of the Nostrils, and has been mistook for a Polypus there.—It is thin at its Rise, and rather still thinner in its Middle; yet afterwards, towards its lower Edge, it becomes thicken, that its Conjunction with the Bones and middle

Cartilage of the Nose might be firmer.

At a little Distance from each Side of this external Process, a cellular and spongy bom Substance depends from the cribriform Plate The Number and Figure of the Cells in this regular Process of each Side, are very unce tain, and not to be represented in Words; of ly the Cells open into each other, and into the Cavity of the Nose: The uppermost, which are below the Aperture of the frontal Simifes, an formed like Funnels.—The outer Surface these Cells is smooth and plain, where this Bow assists in composing the Orbit; at which Place on each Side, it has got the Name of Os pl num; on the upper Edge of which, a final Notch or two may sometimes be observed which go to the Formation of the internal of bitar Holes; as was remarked in the Description of the frontal Bone.

Below the Cells of each Side, a thin Plates extended inwards, and then bending down, becomes thick and of a spongy Texture.—
This spongy Part is triangular, with a streight upper Edge placed horizontally, an anterior of slanting from above, downwards and forwards and with a pendulous convex one below.—It upper and lower Edges terminate in a shar Point behind.—The Side of this pendulous spot

gy Part next to the Septum Narium is convex, and its external Side is concave.—These two Processes of the ethnoid Bone have got the Name of Ossa spongiosa, or Turbinata superiora, from their Substance, Figure, and Situation.

e of

for

d ra.

ard,

cker.

iddle

this

bony

Plate

is is

nce.

; 01.

to the

h are

s, an

ce of

Both

Place

s pli

fmal

erved

nal or

ription

Platel

wn,

e.—

treigh

ior of

ty ard

.—Th

a shan

All the Prominencies, Cavities and Meanders of this ethmoid Bone, are covered with a Continuation of the Membrane of the Nostrils, in a recent Subject.—Its horizontal cribriform Plate is lodged between the orbitar Processes of the frontal Bone, to which it is joined by the athmoid Suture, except at the Back-part, where it is connected with the Cuneiform Bone, by a Suture common to both these Bones, though it is generally esteemed Part of the sphenoidal.—Where the Offa plana are contiguous to the frontal Bone within the Orbit, their Conjunction is reckoned Part of the transverse Suture.—Farther forward than the Offa plana, the Cells are covered by the Offa unguis, which are not only contiguous to these Cells, but cannot be separated from them, without breaking the bony Substance; and therefore, in Justice, those Bones ought to be demonstrated as Part of the ethmoid Bone. Below the Ossa unguis and plana, these Cells and Ossa spongiosa are overlopped by the maxillary Bones.—The celular Part of each palate Bone is contiguous o each Os planum and Cells backwards.—The ower Edge of the Nafal perpendicular Plate is received into a Furrow of the Vomer. -- Its ofterior Edge is joined to the Fore-part of he Processus azygos of the Sphenoid Bone.ts upper Edge joins the Nasal Process of the

frontal and nasal Bones, and its anterior one is connected to the middle Cartilage of the Nose.

From all which, the Uses of this Bone are e. vident, viz. to fustain the anterior Lobes of the Brain; to give Passage to the olfactory Nerves, and Attachment to the Falx; to enlarge the Organ of Smelling, by allowing the Membrane of the Nose a great Extent; to straiten the Passage of the Air through the Nose, by leav. ing only a narrow winding Canal, on the fenf. ble membranous Sides of which the Substances conveyed along with the Air must strike; to form Part of the Orbit of the Eyes and Septum narium; while all its Parts are so light, as not to be in Hazard of separating by their Weight; and they are fo thin, as to form a large Surface, without occupying much Space. This brittle Substance, however, is sufficiently protected from external Injuries by the firm Bones which cover it.

If this Bone is seized on by any corroding Matter, we may easily conceive what Destruction may ensue. Hence it is, that an Ozana is dissicult to cure; and that, in violent Scurvies, or in the Lues Venerea, the Fabrick of the Nose, the Eyes, and Life itself are in Danger.—The Situation of the Nasal Plate may shew us, how dangerous a Fracture of the Bones of the Nose may be, when made by a Force applied to their middle Fore-part, of a Person in whom this nasal Plate is perpendicular.

The Ethmoid Bone of ripe Children is divided into two, by a perpendicular Cartilage, which, when offified, is the Crista galli, and Na

pal

F

tl

W

th

ofe.

e.

the

res, the

ane

the av-

nsi.

ces

to

tun

not ght;

ace,

ittle

hich

ding

truc

72a 15

Tole,

-The

how

Nose their

this

dividilage,

d Na-

fal Plate: But its other Parts are offified and

compleat.

OS SPHENOIDES*, or Wedge-like
Bone, so called because of its Situation in the
Middle of the Bones of the Cranium and Face,
is of such an irregular Figure, that I know not
any Thing to which it may be likened, unless,
perhaps, it bear some faint Resemblance to a
But with its Wings extended.

When we view the Outside of the Os sphenoides, five remarkable Processes may be obferved, which are all of them again fubdivided.-The first and second are the two large lateral Processes or Wings; the upper Part of each of which is called the temporal Process, because they join with the temporal Bones in forming the Temples, and the Seat for some Share of the crotaphite Muscles. That Part of the Wings which jutts out towards the Infide, somewhat lower than the temporal Apophyses, smooth and hollowed, where it makes up Part of the Orbit, is thence named orbitar Processes. Behind the Edge, separating these two Processes, there is often a small Groove, made by a Branch of the superior maxillary Nerve, in its Passage to the temporal Muscle. The lowest and back Part of each Wing, which runs out sharp to meet the Offa petrofa, has been styled the spinous Process: From near the Point of which a sharp-pointed Process is frequently produced downwards, which some call Styliform, that affords Origin to the Ptery-staphylinus exter-K mus

^{*} Cuneiforme, πολύμορφον, multiforme, paxillum, cribratum, palati, colatorii, cavilla, basilare.

mus Muscle. From this styloid Process-a very fmall Groove is extended along the Edge of the Bone to the Hollow at the Root of the in. ternal Plate of the following Processes, which forms Part of the Eustachian Tube (a).—The third and fourth external Processes of the cu. neiform Bone, are the two which stand out al. most perpendicular to the Base of the Scull, Each of them has two Plates, and a middle Fossa facing backwards, and should, to carry on our Comparison, be likened to the Bat's Legs, but are commonly faid to refemble the Wings of that Creature; and therefore are named Pterygoid or Aliform * Processes. The external Plates are broadest, and the internal are longest. From each Side of the external Plates the pte. rygoid Muscles take their Rise. At the Root of each internal Plate, a finall Hollow may be remarked, where the Musculus ptery-staphylimus in termus, or circumflexus palati rifes, and some Share of the cartilaginous End of the Eustach. an Tube rests; and, at the lower End of the same Plate, is a Hook-like Rising or Proces, round which the Tendon of the last named Muscle plays, as on a Pulley. From the Edge of the external Plates some small sharp Spikes stand out; but their Number and Bulk are uncertain.—The fifth external Process of the Sphenoid Bone is that sharp middle Ridge which stands out from its Base: Because it wants a Fellow, it may be called Processus azygos. The lower Part of this Process, where it is received into

CE

20

21

ar

ec

its

* Naviculares.

⁽a) Winstow, Exposition Anatomique du corps humain Traité des Os secs, § 233.

in.

ich

The

cu-

ull.

dle

On

gs,

ngs

ned

est.

pte-

re-

ome chi-

the

es,

ned dge

kes

unthe

nich

ts a The

ved into

nain

into the Vomer, is thick, and often not quite perpendicular, but inclining more to one Side than the other. The Fore-part of this Process, where it joins the Nasal Plate of the Os Ethmoides, is thin and streight. These two Parts have been described as two distinct Processes by some.

The Depressions, Sinuosities and Fossa, on the external Surface of this sphenoid Bone, may be reckoned up to a great Number, viz. two on the temporal Apophyses where the crotaphite Muscles lodge - Two on the orbitar Processes, to make way for the Globes of the Eyes .-Two between the temporal and spinous Processes for receiving the temporal Bones. - Two berween the Plates of the pterygoid Processes, where the Musculi pterygoidei interni and Ptery-staphylini interni are placed. - Two between the pterygoid and orbitar Processes, for forming the Holes, common to this, and to the Cheek and maxillary Bones. Two on the lower Ends of the aliform Processes, which the palate Bones enter into. - Two at the Roots of the temporal and pterygoid Processes, where the largest Share of the external pterygoid Muscles have their Rife. - Two at the Sides of the Processius azygos, for forming Part of the Nose, &c.

What I described under the Name of temporal and spinous Processes on the Out-side of the Scull, are likewise seen on its Inside, where they are concave, for receiving Part of the Brain; and commonly three Apophyses on the internal Surface of the sphenoid Bone are only mentioned—Two rising broad from the Fore-part of its Body, become smaller as they are extended

K 2 obliquely

obliquely backwards. — The third standing on a long transverse Base, near the Back-part of the Body of this Bone, rifes nearly erect, and of an equal Breadth, terminating often in a little Knob on each Side. The three are called Clinoid, from some resemblance which they were thought to have to the Supporters of a Bed. Sometimes one or both the anterior clinoid Processes are joined to the Sides of the posterior one, or to the Body of the Bone itself .-From the Roots of the anterior clinoid Processes the Bone is extended on each Side outwards and forwards, till it ends in a sharp Point, which may have the Name of the Transverse fpinous Processes. — Between, but a little far. ther back than the two anterior clinoid Processes, we see a Protuberance considerably fmaller than the posterior clinoid Process, but of its Shape. — Another Process often forces itself forwards into the Os Ethmoides, from between the transverse Processes.

to

co

ed

be

T

the

Pa

nal

im

Within the Scull, there are two Sinuosities in the internal Part of each Wing of the sphenoid Bone, for receiving the middle Part of the Brain.—One between the transverse spinous Processes, for lodging the Part of the Brain where the Crura medullæ oblongatæ are.—Immediately before the third or middle clinoid Process, a single Pit generally may be remarked, from which a Fossa goes out on each Side to the Holes through which the optic Nerves pass. The Pit is formed by the conjoined optic Nerves; and in the Fossæ these Nerves are lodged, as they run divided within the Scull.—Between that third Protuberance and the posse.

d

li-

0-

les

ds

ıt,

rfe

ar.

0-

oly

out

ces

m

ies

he-

the

ous

ain

m-.

noid

irk-

side

ves

op-

are

1.—
ofterior

rior clinoid Process, the large Pit for the Glandula pituitara may be remarked. This Cavity, because of its Resemblance to a Turkish Saddle, is always described under the Name of Sella Turcica, or Ephippium.—On the Sides of the posterior clinoid Process a Fossa may be remarked, that stretches upwards, then is continued forwards along the Sides of the Sella Turcica, near to the anterior clinoid Processes, where a Pit on each Side is made. These Fossa point out the Course of the two internal carotid Arteries after they have entered the Scull.—Besides all these, several other Fossa may be observed, leading to the several Holes, and imprinted by the Nerves and Blood-vessels.

The Holes on each Side of the Os sphenoides are fix proper, and three common. The first is the round one immediately below the anterior clinoid Processes, for the Passage of the optic Nerve, and of the Branch of the internal carotid Artery that is fent to the Eye.---The fecond is the Foramen lacerum, or large Slit between the transverse spinous and orbitar Processes: The interior End of which Slit is large; and, as it is extended outwards, it becomes narrower. The outer End of it is formed in the Os frontis; and therefore this might be reckoned among the common Foramina. Through it the third, fourth, the first Branch of the fifth, and the greater Share of the fixth Pair of Nerves, and an Artery from the internal Carotid, go into the Orbit. Sometimes a imall Branch of the external Carotid enters near

K 3

its End, to be distributed to the Dura Mater (a), and a Vein, some call it the Venous Duct, or, Nuck's Aquæduct, returns through it to the ca. vernous Sinus. — The third Hole, fituated a little behind the one just now described, is called Rotundum, from its Shape. It allows passage to the second Branch of the fifth Pair of Nerves. or fuperior maxillary Nerve, into the Bottom of the Orbit ___ The fourth is the Foramen o. vale, about half an Inch behind the round Hole. Throughit the third Branch of the fifth Pair, or inferior maxillary Nerve, goes out; and sometimes a Vein from the Dura Mater passes out here (b). — Very near the Point of the spinous Process is the fifth Hole of this Bone: It is fmall and round, for a Passage to the largest Artery of the Dura Mater, which often is accompanied with a Vein. The fixth proper Hole (c) cannot be well feen, till the cuneiform Bone is separated from all the other Bones of the Cranium; for one End of it is hid by a small Protuberance of the internal Plate of the pterygold Process, and by the Point of the Processiv petrofus of the temporal Bone. Its Canal is extended above the inner Plate of the pterygoid Process; and where it opens into the Cavity of the Nose, it is concealed by the thin laminous Part

A

fa

An

(b) Ingraff. Commentar. in Galen. de Offib. lib. 1. comment. 8

⁽a) Winstow, Exposition Anatomique de corps humain, Traité des Arteres, §. 60. & de la Tete, §. 26.

⁽c) Vefal. Anat. lib. 1. cap. 12.—Eustach. Tab. 46. Fig. 33. & 16. —Vidus Vidius, Anat. lib. 2. cap. 2. Explicat Tab. 5. & Tab. 5. Fig. 8. 9. 10. lit. O.

r,

2.

2

ed

es,

m

0-

le.

10

ne-

out

ous

- is

eft

ac-

per

rm

of

77.

fis

ex-

roid

of

ous

nain,

com-

Fig.

Part of the palate Bone. Through it an Artery from the external Carotid runs, to be distributed to the Nostrils: So that as this Hole does not open within the Cranium, it differs from all the other Foramina of this Bone already described. In several Subjects, soon after the Entry of this Artery, a small Branch goes backwards from it to the Fossa pterygoidea, and a second enters the Scull to ferve the Dura Mater. — Often in the Middle of the Sella Turcica a small Hole or two pierce as far as the cellular Substance of the Bone: and fometimes at the Sides of this Sella, one or more small Holes penetrate intothe sphenoidal Sinuses. These Observations afforded some Anatomists (a) an Argument of Weight in their Days in defence of Galen (b), who afferted the Descent of the Pituita that Way into the Simises below.

The first of the common Holes is that unequal. Fisher at the Side of the Sella Turcica, between the extreme Point of the Os petrosum and the spinous Process of the cuneiform Bone. This Hole only appears after the Bones are boiled; for, in a recent Subject its Back-part is covered by a thin bony Plate that lies over the internal carotid Artery, and further forward it is filled with a cartilaginous Ligament, under which the cartilaginous Part of the Eustachian Tube is placed: Sometimes, however, a Branch of that Artery which enters the fixth Hole of the sphenoid Bone, is sent through this Substance to the Dura Mater. It was by this Pasage that the Ancients believed the slimy Matter

was

(b) Galen. De usu Part. lib. 9. cap. 1.

⁽a) Jac. Sylv. Calumniæ secundæ amolitio. — Laurent. Hist. Anat. lib. 2. quæst. 11.

was conveyed from the Emunctory of the Brain. the Glandula pituitaria, to the Fauces. The fecond common Hole is the large Discon. tinuation of the external Side of the Orbit, left between the orbitar Processes of the cuneiform Bone, the Os maxillare, male, and pa. lati. In this large Hole the Fat for lubricating the Globe of the Eye and temporal Muscle is lodged, and Branches of the fuperior maxillary Nerve, with small Arteries from the Carotid and Veins pass. — The third Hole is formed between the Base of this Bone and the Root of the Orbitar Process of the Palate-bone of each Side. Through this a Branch of the external carotid Artery, and of the second Branch of the fifth Pair of Nerves, are allowed a Passage to the Nostrils, and a returning Vein accompanies them. Sometimes, however, this Hole is proper to the Palate-bone, being entirely formed out of its Substance.

Under the Sella Turcioa, and some Way farther forward, but within the Substance of the sphenoid Bone, are two Simises, separated by a bony Plate. Each of them is fined with a Membrane, and opens into the upper and Backpart of each Nostril by a round Hole, which is at their upper Fore-part. This Hole is not formed only by the Os sphenoides, which has an Aperture near as large as any transverse Section of the Sinus, but also by the Palate Bones which are applied to the Forepart of these Simises, and close them up, that Hole only excepted, which was already mentioned. Frequently the two Sinuses are of unequal Dimensions, and sometimes there is only

m

le

th

m

the

exi

Na

only one large Cavity, with an Opening into one Nostril. These Cavities are likewise said (a) to be extended sometimes as far back as the great Foramen of the occipital Bone. In other Subjects they are not to be found, when the Bone is composed of large Cells (b). Some (c) mention a Cavity within the Partition of the Sinuses; but it is small.—The sphenoidal Sinuses serve the same Uses as the frontal do.

As this Bone is extremely ragged and unequal, so its Substance is of very different Thickness, being in some Places diaphanous; in others it is of a middle Thickness, and its middle Back-part surpasses the greatest Share of the

Cranium in Thickness.

18

id

ed

of.

ch.

al

of

)a-

19

m-

ar-

the

ya

1 4

ck-

ich

not

ich

anf-

the

ore-

that

en-

e is

only

The Os sphenoides is joined, by its Wings, to the parietal Bones above, to the Os frontis and Offa malarum before, to the temporal Bones behind; -- by the Fore-part of its Body and spinous Processes, to the frontal and Ethmoid Bones; -by its Back-part, behind the two Simifes, to the occipital, where it looks like a Bone with the Epiphyses taken off, and, as was formerly observed in the Description of the occipital Bone, it cannot be separated without Violence in Adults;—to the Palate Bones, by the Ends of the pterygoid Processes, and still more by the Fore-part of the internal Plates of the pterygoid Processes, and of the Simsses;—to the maxillary Bones, by the Fore-part of the external pterygoid Plates; - to the Vomer and Nasal Plate of the Os Ethmoides, by the Pro-

(c) Id. ibid.

⁽a) Albin. de Ossib. § 39. (b) Vesal. lib. 1. cap. 6.

cessis azygos. All these Conjunctions, except the last, which is a Schindylesis, are said to be by the Suture proper to this Bone; though it is at first Sight evident, that several other Sutures, as the transverse, ethnoidal, &c. are consounded with it.

We fee now how this Bone is joined to all the Bones of the Cranium, and to most of the upper Jaw; and therefore obtained the Name of the Wedge, like Bone.

The Uses are so blended with the Description, as to leave nothing new to be added concerning

them.

The *sphenoid* Bone is almost compleat in a Fætus of nine Months; only the great Alæ separate after Maceration from the Body of the Bone.—The *Processus azygos* is very large and hollow;—the internal Surface of the Body is unequal and porous;—the Sinuses do not ap-

pear.

Whoever is acquainted with each Bone of the Cranium, can, without Difficulty, examine them as they stand united, so as to know the Shapes, Sizes, Distances, &c. of their several Parts, and the Forms, Capacities, &c. of the Cavities formed by them, which is of great Use towards understanding the Anatomy of the Parts contiguous to, contained within, or connected to them. Such a Review is necessary, after considering each Class of Bones. Thus the Orbits, Nostrils, Mouth, Face, Head, Spine, Thorax, Pelvis, Trunk, Extremities, and Skeleton, ought likeways to be examined.

The FACE is the irregular Pile of Bones, composing the fore and under Part of the Head,

which

te

fo

177

W

au

M

pa

Spo

which is divided, by Authors, into the Upper

and Lower Maxillæ or Jaws.

is

S,

d-

Ill

he

110

n,

ng

fe-

the

ind is

ap-

the

em

oes,

rts,

ards-

nti-

con-

oits,

rax,

ton,

nes,

ead,

hich

The fuperior Maxilla + is the common Defignation given to the upper immoveable Share of the Face; though, if we would follow Celfus (a), we should apply the Word Maxilla to the Lower Jaw only, and use the Name Mala for this Upper Jaw. In Complaisance to prevailing Custom, I shall, however, follow the Terms now commonly employed.—The Shape of the superior Jaw cannot easily be expressed; nor is it necessary, provided the Shape and Situation of all the Bones which compose it are described. It is bounded above by the transverse Suture, behind by the Fore-part of the sphenoid Bone, and below by the Mouth.

The upper Jaw consists of six Bones on each Side, of a thirteenth Bone which has no Fellow, placed in the Middle, and of sixteen Teeth. The thirteen Bones are, two Ossa nasi, two Ossa unguis, two Ossa malarum, two Ossa maxillaria, two Ossa palati, two Ossa spongiosa

inferiora, and the Vomer.

The Ossa nasi are placed at the upper Part of the Nose;—the Ossa unguis are at the internal Canthi of the Orbits;—Ossa malarum form the Prominence of the Cheeks;—Ossa maxillaria form the Side of the Nose, with the whole Lower and Fore-part of the Upper Jaw, and the greatest Share of the Roof of the Mouth;—Ossa palati are situated at the Backpart of the Palate, Nostrils, and Orbit;—Ossa spongiosa are seen in the lower Part of the Na-

res ;-

[†] Σίαγων, γένος, Mandibula.
(a) Lib. 8. cap. .

res; — and the Vomer helps to separate these two Cavities.

The Bones of the Upper Jaw are joined to the Bones of the Scull by the Schindylesis and Sutures already described as common to the Cranium and Face, and they are connected to each other by Gomphosis and fifteen Sutures.

The Gomphosis only is where the Teeth are fixed in their Sockets, and the Schindyless is only where the Edges of the Vomer are joined

to other Bones.

The Sutures are generally distinguished by Numbers, which have been differently applied; and therefore I join those (a) who prefer the giving. Names to each, which may be easily contrived from their Situation, or from the Bones which they connect.

The first is the anterior Nasal +, which is streight, and placed longitudinally in the mid-

dle Fore-part of the Nofe.

The second and third are the lateral Nasal*, which are at each Side of the Nose, and almost parallel to the first Suture.

Each of the two Lacrymal is almost semicircular, and is placed round the lacrymal Groove

The fixth and seventh are the internal Orbitar; each of which is extended obliquely from the Middle of the lower Side of an Orbit to the Edge of its Base.

The

ti

h la fin

⁽a) Vander Linden, Medicin. physiolog. cap. 13. art. is \$ 10-Rolfinc. Anat. lib. 2. cap. 25.—Schenk. Schol. part. § ult. par. 2. cap. 5.

[†] Nafalis Recta. * Nafalis Obliqua.

The two external Orbitar are continued, each from the End of the internal Orbitar, to the Under and Fore-part of the Cheek.

ele

to

and

the

d to

are

5 18

ined

l by

ied;

the

afily

the

ch is

mid-

fal *,

most

nicir.

Orbi-

from o the

The

art. 4

The tenth is the mystachial, which reaches only from the lower Part of the Septum narium to between the two middle Dentes Incifores.

The longitudinal Palate + Suture stretches from the Middle of the foremost Teeth thro' the Middle of all the Palate.

The transverse Palate one ‡ runs across the Palate, nearer the Back than the Fore-part of it.

Each of the two *Palato-maxillary* is at the Back-part of the Side of each Nostril.

The fifteenth is the Spinous, which is in the Middle of the lower Part of the Nostrils. This may, perhaps, be rather thought a double Schindylesis.

The Connexion of the Ossa spongiosa to the Side of each Nostril, is so much by a Membrane in young Subjects, and by Concretion or Union of Substance in Adults, that I did not know well how to rank it: But if any chuses to call it a Suture, the Addition of two transverse Nasal Sutures may be made to those above named.

These Sutures of the Face (formerly called Harmoniæ) have not such conspicuous Indentations as those of the Scull have; the Bones here not having Substance enough for forming large Indentations, and there being less Necelsity for Security against external Injuries, or any internal protruding Force, than in the Cranium

[†] Laquearis, Palataria recta.

Arcuata, Palatina postica.

nium.—These Sutures often disappear in old People, by the Bones running into each other; which can do little Prejudice, because the principal Use of the Bones being so numerous here, is to allow them to be extended into a proper Form.

It is evident, from the Manner of the Conjunction of these Bones, that they can have no Motion, except in common with the Crenium.

The Purposes which this Pile of Bones serves, will be shewn in the Description which I am

to give of each.

OSSA NASI, so named from their Situation at the Root of the Nose, are each of an irregular oblong square Figure, being broadest at their lower End, narrowest a little higher than their Middle, and becoming somewhat larger at the Top, where they are ragged and thickest, and have a Curvature forwards, that their Connexion with the frontal Bone might be stronger.—These Bones are convex externally, and thereby better resist any Violence from without; and they are concave internally, for enlarging the Cavity of the Nose.

The lower Edge of these Bones is unequal, and is stretched outwards and backwards, to join the Cartilages of the Nostrils.—Their anterior Side is thick, especially above, and unequal, that their Conjunction to each other might be stronger; and a small Rising may be remarked on their inner Edge, where they are sustained by the Septum narium.—Their posterior Side, at its upper Half, has externally a Depression, where it is overlopped some Way by the maxillary Bones, while its lower Half covers these

Bones:

Bones: By which Contrivance, they do not yield easily to Pressure applied to their Fore-

part or Sides.

ble

er;

re,

011-

no

es,

am

ua.

an

left

her

hat

und

hat

ht

er-

nce

lly,

and

oin-

eri-

ial,

be

ked

ned

de,

on,

xil-

lele

es:

A finall Hole is frequently to be observed on their external Surface, into which two, three, or four Holes, which appear internally, terminate for the Transmission of small Veins, sometimes the Holes go no further than the Cancelli of the Bones.

The Nafal Bones are firm and folid, with very few Cells or Cancelli in them; the thin Subtlance, of which they confist, not requiring

much Marrow.

They are joined above to the frontal Bone, by the Middle of the transverse Suture;—behind, to the maxillary Bones, by the lateral Nafal Sutures;—below, to the Cartilages of the Nose;—before, to one another, by the anterior Nasal Suture;—internally, to the Septum narium.

These Bones serve to cover and defend the

Root of the Nofe.

In an Infant the Nafal Bones are proportionally shorter, and less thick at their upper Part, than in an Adult, but are otherwise compleat.

OSSA UNGUIS, or LACRYMALIA, are so named, because their Figure and Magnitude are something near to those of a Nail of one's Finger, and because the Tears pass upon them into the Nose.

Their external Surface is composed of two smooth Concavities and a middle Ridge.—
The Depression behind forms a small Share of the Orbit for the Eye-ball to move on, and the one before is a deep perpendicular Canal, or

L 2 Fossai,

Fossa, larger above than below, containing Part of the lacrymal Sac and Duct. This is the Part that ought to be pierced in the great Operati. on for the Fiftula lacrymalis .- This Fossa of the Bone is cribriform, or has a great Number of small Holes through it, that the Filaments from the Membrane which lines it, infinuating themselves into these Holes, might prevent a Separation of the Membrane, and fecure the Bone in its natural Situation. - The Ridge between these two Cavities of the Os unguis, is the proper Boundary of the Orbit at its internal Canthus; and beyond which Surgeons should not proceed backwards in performing Operations here.—The internal or posterior Surface of this Bone confifts of a Furrow in the Middle of two Convexities.

The Substance of the Os unguis is as thin as Paper, and very brittle; which is the Reason that those Bones are often wanting in Skeletons, and need little Force to pierce them in

living Subjects.

Each of these Bones is joined, above, to the frontal Bone, by Part of the transverse Suture;—behind, to the Os planum of the Ethmoid Bone, by the same Suture;—before, and below, to the maxillary Bone, by the lacrymal Suture.—Internally, the Ossa unguis cover some of the Sinus ethmoidales; nay, are really continuous with the bony Lamella, which make up the Sides of these Cells; so that they are as much Part of the Ethmoid Bone, as the Ossa plana.

These unguisorm Bones compose the anterior internal Parts of the Orbits, lodge a Share of the lacrymal Sac and Duct, and cover the eth-

flance, make a rash Operator in Danger of destroying a considerable Share of the Organ of Smelling, when he is performing the Operation of the Fistula lacrymalis; but when these Bones are hurt, they cast off without much Dissiculty, and consequently the Wound is soon cured, unless the Patient labours under a general Cacoethes, or there is a Predisposition in the Bones to Caries; in which Case, a large Train of bad Symptoms sollows, or, at best, the Care proves tedious.

These Bones are fully formed in a new born

Child.

Part

Part

rati-

a of

nber

ents

ting

nt a

the

idge

5, 18

iter-

ould

rati-

face

Mid-

n as

ason

ele-

n in

the

ure:

noid

OW,

ure.

the

10US

the.

uch

erior eth-

12014

a.

OSSA MALARUM* was the Name given by Celsus, as was already remarked, to all the Upper Jaw; but is now appropriated to the prominent square Bones, which form the Cheek on each Side.—Before, their Surface is convex and smooth; backward, it is unequal and concave, for lodging Part of the Crotaphyte Muscles.

The four Angles of each of these Bones have been reckoned Processes by some.—The one at the external Canthus of the Orbit, called the superior orbitar Process, is the longest and thickest.—The second terminates near the Middle of the lower Edge of the Orbit in a sharp Point, and is named the inserior orbitar Process.—The third, placed near the lower Part of the Cheek, and thence called maxillary, is the shortest, and nearest to a right Angle.—The fourth, which is called Zygomatic, because it is extended

^{*} Jugalia vel Zygomatica, hypopia, subocularia,

extended backwards to the Zygonia of the tem. poral Bone, ends in a Point, and has one Side streight, and the other sloping.—Between the two orbitar Angles there is a concave Arch, which makes about a Third of the external Circumference of the Orbit, from which a fifth Process is extended backwards within the Orbit, to form near one Third of that Cavity; and hence it may be called the internal orbitar Process.—From the lower Edge of each of the Offa malarum, which is between the maxillary and zygomatic Processes, the Masset Muscle takes its Origin; and from the exterior Part of the Zygomatic Process, the Musculus distortor oris rifes; in both which Places the Surface of the Bone is rough.

On the external Surface of each Cheek-bone, one or more small Holes are commonly sound, for the Transmission of small Nerves or Blood-vessels from, and sometimes into the Orbit.—On the internal Surface are the Holes for the Passage of the nutritious Vessels of these Bones.—A Notch on the Outside of the internal orbitar Process of each of these Bones, assists to form the great Slit common to this Bone, and to the Sphenoid, Maxillary, and Palate Bones.

The Substance of these Bones is, in Proportion to their Bulk, thick, hard, and solid, with

fome Cancelli.

Each of the Offa malarum is joined, by its fuperior and internal orbitar Processes, to the Os frontis, and to the orbitar Process of the sphenoid Bone, by the transverse Suture.—By the Edge between the internal and inferior orbitar Processes to the maxillary Bone, by the internal

ternal orbitar Suture.—By the Side between the maxillary and inferior orbitar Process, again to the maxillary Bone, by the external orbitar Suture.—By the zygomatic Process to the Os temporum, by the zygomatic Suture.

The Cheek-bones are entire, and fully offifi-

ed in all their Parts in Infants.

m-

de

he

ch,

nal fth

oit,

ro-

the

ary

of

of of

me,

md,

od-

the

nes.

s to

and

nes.

ortiwith

v its

the

the By

r or-

21714

ossa Maxiellaria superior A, are the largest Bones, and constitute the far greater Part of the Upper Jaw, which has appropriated the Name of Maxislaria to them. The Figure of one of them, or of the two when joined, is so irregular, that Words can

scarce give an Idea of it.

The Processes of each Os maxillare may be reckoned feven .- The first is the long Nasal one at its upper and fore Part, which is broad below, and turns smaller, as it rises upwards, to make the Side of the Nofe.—At the Root of this, a transverse Ridge may be observed within the Nostrils, which supports the Forepart of the upper Edge of the Os fpongiofum inferius. The fecond is produced backwards and outwards, from the Root of the Nafal Procels, to form the lower Side of the Orbit; and therefore may be called orbitar. The Edge of this orbitar Process, and the Ridge of the nafal one, which is continued from it, make a considerable Portion of the external Circumference of the Orbit.—From the proper orbitar Process, a very rough triangular Surface is extended downwards and outwards, to be connected to the Cheek-bone; and therefore may be called the malar Process, from the lowest protuberant Part of which some Share of the maffeter

masseter Muscle takes its Rife. - Behind the orbitar Process, a large Tuberosity, or Bulge of the Bone appears, which is esteemed the fourth Process.—On the internal Part of this we often meet with a Ridge, almost of the fame Height with that in the Nafal Process, which runs transversely, and is covered by a similar Ridge of the palate Bone, on which the Back-part of the upper Edge of the Os spongi. ofum inferius rests. The convex Back-part of this Tuberosity is rough, for the Origin of Part of the external pterygoid Muscle (a), and more internally is feabrous, where the Palate and sphenoid Bones are joined to it. - That fpongy Protuberance * at the lower Circumference of this Bone, where the Sockets for the Teeth are formed, is reckoned the fifth.—The fixth is the horizontal Plate, which forms the greater Part of the Base of the Nostrils, and Roof of the Mouth: Its upper Surface, which belongs to the Nostrils, is very smooth, but the other below is arched and rough, for the stronger Adhesion of the Membrane of the Mouth, which is stretched upon it, and, in chewing, speaking, &c. is liable to be separate ed.—The feventh rifes like a Spine from the inner Edge of the last, and forms a small Part of the Partition of the Nostrils.

The Depressions in each maxillary Bone are, 1. A Sinuosity behind the orbitar Process, made by the temporal Musele. 2. A Pit immediately before the same Process, where the Origin of

C

n

11

la

01

ce

ly

ce

⁽a) Albin, de Offib. §. 79.

ne

he

he

efs,

fi-

the

igi-

art of

and

late hat

fer-

the

The

the

and

hich

but

the

the

, in

arat.

the

Part

are,

made

ately

zin of

the

the Musculus elevator labiorum communis, and Elevator labii superioris, with a Branch of the fifth Pair of Nerves, are lodged fecurely. hollow Arch of the Palate. 4. The femicircular great Notch, or Entry to the lower Part of the Nostrils; betwixt the Root of the Nasal Process and Spine of the palate Plate-Below this, the Fore-part of the Bone is flatted, or fometimes hollowed by the Musculus depressor labii superioris. 5. Sockets for the Teeth +: The Number of these Sockets is uncertain; for the same Number of Teeth is not in all People, and the four backmost Teeth of each Jaw vary greatly in their Number of Roots; and when the Teeth of a living Person fall out, or are taken away, the Sockets fill up with an offeous Net-work, which becomes folid afterwards.— 6. The lacrymal Fossa in the Nasal Process, which assists the Os unguis to form a Passage for the lacrymal Duct. — Immediately on the Out-side of this, there is a small Depression, from which the inferior or leffer oblique Mufcle of the Eye has its Origin (a). 7. The Canal on the upper Part of the great Tuberofity within the Orbit, which is almost a compleat Hole; in this a Branch of the superior maxillary Nerve paffes. - Befides thefe, the fuperior Surface of the great Bulge is concave, to receive the under Part of the Eye. - Immediately above the transverse Ridge in the Nasal Procels, a small Hollow is formed by the Os spongiofum --- In some Subjects, the Nasal Process, has

[†] Βόδρια, ὁλμίζαοι, Alveoli, Fossulæ, mortariola, Fræna, Locelli, Cavæ, Præsepiola, Loculamenta.
(a) Winstow, Exposition Anatomique des Os Secs, §. 276.

has a small round Pit above the lacrymal Duck, where the little Tendon or Ligament of the orbicular Muscle of the Eye-lids is inserted. It is this Tendon, and not the Tendon of the larger oblique Muscle of the Eye, which there is a Probability of cutting in the Operation of

the Fiftula lacrymalis.

The Holes of this Bone are two proper and two common, which are always to be found, besides several others, whose Magnitude, Num. ber, &c. are uncertain. The first of the proper is the external Orbitar, immediately be low the Orbit, by which the infra-orbitar Branch of the fecond Branch of the fifth Pair of Nerves, and a finall Artery, come out, after having passed in the Canal, at the Bottom of the Orbit, described Numb. 7. of the Depreifions - This Hole is often double, and that when the Nerve has happened to split before it has escaped from the Bone. - The second is the Foramen incifivum, just behind the Fore-teeth which, at its under Part, is one irregular Hole common to both the maxillary Bones when they are joined; but, as it ascends, soon divides into two, three, or fometimes more Holes; fome of which open into each Nostril. Through them small Arteries and Veins, and a Twig of the fecond Branch of the fifth Pair of Nerves pass, and make a Communication between, or join the lining Coats of the Nose and Mouth-In some Subjects, Steno's Duct may be traced fome way on the Side of these Passages next to the Nose, and small Orifices may be observed opening into the Mouth. The

1

t

i

p:

fc

th

tw

in

The first common Hole is that which appears at the inner Side of the Back-part of the Tuberosity and of the Sockets of the Teeth, and is formed by a Fossa in this Bone, and a corresponding one in the Os palati: Through it a Nerve, which is a Branch of the second Branch of the fifth Pair, runs to the Palate.—The other common Hole is the great Slit in the Out side of the Orbit described already, as the second common Hole of the sphenoid Bone.

On the Nafal Process often Holes may be obferved for the Passage of Vessels to the Substance of the Bones; and, at the Back-part of each Tuberosity, several Foramina are placed, for the Transmission of Nerves to the Cavity

within: But those are uncertain.

All the Body of the maxillary Bone is hollow, and leaves a large Simus a-kin to the frontal and sphenoid, which is commonly, but unjustly, called Antrum Highmorianum*. When the Os maxillare is fingle or feparated from all the other Bones of a Skeleton, its Antrum appears to have a large Aperture into the Nostrils; but, in a recent Subject, it is so covered at its Backpart, by the palate Bone; in the Middle, by the Os spongiosium inferius; before, by a strong Membrane, that one, or fometimes two Holes scarce larger than a Crow-quill, are only left at the upper Part; which, after a short winding Progress, open into the Nostrils between the two Offa spongiofa. — At the Bottom of this Cavity, we may often observe some Protuberances, in which the small Points of the Roots of the Teeth

ud,

the

It

the

n of

and

and,

um.

the

be

oitar

11. 0

after

n of

prel-

that

re it

the

eeth

Hole

they

vides

oles;

ough

g of

erves

1, Of

th.-

raced xt to

erved

The

^{*} Genæ.

Teeth are contained (a).—This Cavern, and the Sockets of the Teeth are often divided by the Interpolition only of a very thin bony Plate, which is liable to be eroded by acrid Matter collected in the Antrum, or to be broke in drawing a Tooth (b). The Symptoms of a Collection of Matter here naturally lead us to the Practice of pulling out the Teeth, and piercing through this Plate into the Antrum, to procure an Evacuation of the collected Matter; by which considerable Service is frequently done (c).

The maxillary Sinuses have the same Uses as the frontal and sphenoidal; and the Situation of the Sinuses is such, that the Liquor drilling from them, from the Cells of the Ethmoid and Palate Bones, and from the lacrymal Duck, may always moisten all the Parts of the Membrane of the Naves in the different Situations

which the Head is in.

Though the Membranes, which line the frontal, sphenoidal, and maxillary Sinuses, are Continuations of the one which covers the Bones within the Nose; yet they are much thinner than it is, and have so much smaller Vessels, that the Injection which makes the Membrane of the Nose red all over, fills only some few Vessels of the maxillary Sinuses, and scarce is observed in the frontal and sphenoidal. Are not the larger Vessels intended for a more plentiful Secretion of a viscid Liquor to defend the Membrane from the Effects of the Persa.

S

pa

th

720

hv

Pa

12.2

of

Can ftri

ture

⁽a) Highmore, Disquis. Anat. lib. 3. part 2. cap. 1.

⁽b) Highmore, ibid.
(c) Cowper in Drake's Anthropol. Book 3. chap. 10.—
Medical Eslays and Observ. Vol. V. Art. 30.

tus, which is constantly through the Nose? Are not the Membranes which have the smallest Vessels, cæteris paribus, the most sensible? Are not many Phænomena of Smelling, Inslammations of these Parts, Megrim, Polypi, &c. depending on this Structure of these Membranes?

The Substance of the Ossa maxillaria is compact and firm, except at the inferior Process, in which the Teeth are lodged, where it is very

loongy.

nd

by

te,

ter

W.

0.

the

by

(c).

s as tion

ling

and ucts,

em-

ions

the

are

the

nuch

the :

only

, and

roidal.

more

efend

ertla.

1.

10,--

till

The maxillary Bones are joined above by the upper Ends of their nafal Processes to the Os frontis, by the transverse Suture;—at the Sides of these Processes, to the Osfa unguis, by the lacrymal Sutures;—to the nafal Bones, by the lateral Nasal Sutures; - by their orbitar Processes, to the Cheek Bones, by the external orbitar Sutures :- by the internal Sides of the internal orbitar Processes, to the Ossa plana, by Part of the ethmoidal Suture; --- by the Back-part of the Tuberosities, to the palate Bones, by the Suturæ palato-maxillares; — by the posterior Edges of their palatine Lamella, to the Offa palati, by the transverse Palate Suture; by their Nafal Spines, to the Vomer, by the spinous Suture; --- by their Sockets, to the Teeth, by Goniphosis; — by the internal Edge of the Palate Plate, to one another, by the longitudinal Palate Suture; on the Upper and Fore-part of which a Furrow is left for receiving the Cartilage which torms the Partition of the Nostrils; -between the Fore-part of the Nostrils and Mouth, to each other, by the mystachial Suture;—fometimes they are connected to the

Offa spongiosa inferiora, by a plain Concretion

or Union of Substance.

These Bones form the greater Part of the Nose and of the Roof of the Mouth, and a confiderable Share of the Orbit. They contain fixteen Teeth, give Rise to Muscles, Transmis. fion to Nerves, &c. as mentioned in the De.

fcription of their feveral Parts.

In each of the maxillary Bones, of a new. born Child, the external orbitar Process is hol. low with remarkable Holes in it; -- there are five Sockets for the Teeth, of which the two posterior are very large, and, when divided by a second cross Partition, make the Number of Sockets fix (a). The Palate Plate is cribii. form about the Middle. The great Tubero. fity is not formed; —instead of the Antrum, there is only an oblong Depression at the Side of the Nostrils.

8

T

fo

of

co

dle

cal

thi

Co

gre que

the

OSSA PALATI are commonly described as two small square Bones, at the Back-part of the Palate or Roof of the Mouth, though they are of much greater Extent, being continued up the Back-part of the Nostrils to the Orbit (b). Each Palate-bone may therefore be divided into four Parts, the Palate Square-bone, the pterygoid Process, nasal Lamella, and orbitar Pro-

ceis.

The Square-bone is unequally concave, for enlarging both the Month and Cavity of the Note.

(a) Albin. Ofteogen. Tab. 5. Fig. 45. - Ungebav. de Dentit. secund. jun, § 1.

⁽b) Eustach. Tab. 47. Fig. 1. 3, 6, 7, 8. - Vidus Vidus de Anat. lib. 2. cap. 2. Explicat. Tab. 6. Fig. 19. - Winstew Memoires de l'Acad. des Sciences, 1720.

on

he

in if-

)e.

W-

ol-

are

OW

by

of

ori.

ero.

um,

Side

d as

the

are

up

(b).

in-

pte-

Pro.

for

the

Jole.

av. de

Vidius,

inflow,

commu-

Nofe. The upper Part of its internal Edge rifes into a Spine, after the same Manner as the Palate Plate of the maxillary Bone does, to be joined with the Vomer .- Its anterior Edge is unequally ragged, for its firmer Connexion with the Palate Process of the Os maxillare .-The internal Edge is thicker than the rest, and of an unequal Surface, for its Conjunction with its Fellow of the other Side. Behind, this Bone is somewhat in Form of a Crescent, and thick, for the firm Connexion of the Velum pendulum palati; the internal Point being produced backwards, to afford Origin to the palato-staphylimis or azygos Muscle.—This Square-bone is well distinguished from the pterygoid Process by a perpendicular Fossa, which, applied to such another in the maxillary Bone, forms a Passage for the palatine Branch of the fifth Pair of Nerves; and by another small Hole behind this, through which a Twig of the same Nerve passes.

The pterygoid Process is somewhat triangular, having a broad Buse, and ending smaller above. The Back-part of this Process has three Fossia formed in it; the two lateral receive the Ends of the two Plates of the sphenoid Bone, that are commonly compared to a Bat's Wing; the middle Fossia makes up Part of what is commonly called the Fossia pterygoidea; the Fore-side of this palatine pterygoid Process is an irregular Concave, where it receives the Back-part of the great Tuberosity of the maxillary Bone.—Frequently several small Holes may be observed in this triangular Process, particularly one near the Middle of its Base, which a little above

M 2

communicates with the common and proper Holes of this Bone already taken Notice of.

The Nafal Lamella of this Bone is extremely thin and brittle, and rifes upwards from the up. per Side of the external Edge of the Square. bone, and from the narrow Extremity of the pterygoid Process; where it is so weak, and, at the same Time, so firmly fixed to the maxillary Bone, as to be very liable to be broken, in feparating the Bones .- From the Part where this Plate rifes, it runs up broad on the Infide of the Tuberosity of the maxillary Bone, to form a confiderable Share of the Sides of the maxillary Sinus, and to close up the Space between the Sphenoid and the great Bulge of the maxillary Bone, where there would otherwife be a large Slit opening into the Nostrils (a).-From the middle internal Side of this thin Plate, a cross Ridge placed on such another of the maxillary Bone is extended; on it, the Back-part of the Os spongiosum inferius rests:-Along the Outside of this Plate, the perpendicular Fossa made by the Palate Nerve is observable.

At the upper Part of this Nasal Plate, the Palate Bone divides into two Processes, which I already named orbitar;—between which and the Body of the sphenoid Bone, that Hole is formed, which I mentioned as the last of the Holes common to the sphenoid Bone.—Sometimes this Hole is wholly formed in the Os palati, by a cross Plate going from the one orbitar Process to the other. A Nerve, Artery and Vein

⁽a) Albin. de Offib. § 88.

per

lely

up.

are-

the

, at

llary

i le-

here

ifide

to

the

be-

the

rwife

a).-

thin

er of

the

ts:-

endi-

serv.

, the

which

h and

ole 18

f the

Some.

Os pa

orbi-

y and

Vein

Vein belonging to the Nostrils pass here.-The anterior of the two orbitar Processes is the largeft, and has its Fore-part contiguous to the Back-part of the maxillary Simus, and its upper Surface appears in the Bottom of the Orbit, bchind the Back-part of the Os maxillare and planum.—It has Cells behind resembling those of the ethmoid Bone, to which it is contiguous; it is placed on the Aperture of the Simis Sphenoidalis, fo as to leave only a round Hole at its upper Fore-part.—The other Part of the orbitar Process is extended along the internal Side of the upper Back-part of the maxillary Tuber ofity, to the Base of the sphenoid Bone, between the Root of the Processis azygos and the pterygoid Process.

The Palate square Part of this Palate Bone, and its pterygoid Process, are firm and strong, with some Cancelli; but the Nasal Plate and or-

bitar Processes are very thin and brittle.

The Palate Bones are joined to the maxillary, by the Fore-edge of the Palate Square-bone, by the transverse Palate Suture :- By their thin-Nafal Plates, and Part of their orbitar Processes, to the same Bones, by the Palato-maxillares Sutures: - By their pterygoid Processes, and Backpart of the Nasal Plates, to the Ala vespertiliomm, by the Sphenoid Suture: -- By the transverse Ridges of the Nasal Plates, to the Offa spongiosa inseriora, by Contact; hence frequently there is an intimate Union of the Substance of these Bones in old Sculls: By the orbitar Processes, to the Offa plana and Cellula ethmoidea, by the ethmoid Suture:---To the Body of the Sphenoid Bone, by the Sphenoid Suture:-M 3 By

By the internal Edge of the Square-bones to each other, by the longitudinal Palate Suture; and by their Nasal Spines to the Vomer, by the spinous Suture.

The Palate Bones form Part of the Palate, Nostrils, Orbits, and Fossæ pterygoideæ, and they cover Part of the Sinus maxillares, sphe.

noidales and ethmoidei.

These Bones are very complete in a newborn Infant, the Nasal Plates being then thicker and stronger than in Adults; but the orbitar Processes have not the Cells which appear in the Bones of Adults.

When we are acquainted with the History of these Bones, the Reason is evident, why the Eyes are so much affected in Ulcers of the Palate, as to be often attended with Blindness, which frequently happens in an ill-managed Lues Venerea; or why, on the other Hand, the

Palate fuffers from an Ægylops (a).

OSSA TURBINATA, or spongiosa inseriora, resemble the superior Ossa spongiosa in Shape and Substance, but have their anterior and upper Edges contiguous to the transverse Ridges of the Nasal Processes of the maxillary and Palate Bones.—From their upper streight Edge, two small Processes stand out: The posterior, which is the broadest, descends to cover some of the Antrum Highmorianum; the anterior rises up to join the Os unguis, and to make Part of the lacrymal Duct.

Below the fpongy Bones already mentioned, there are fometimes two others, one in each

Nostril,

⁽a) Hoffman. in Ephemerid, German. Cent. 1. and 2. 0b. ferv. 135.

Nostril, which seem to be a Production of the Sides of the maxillary Simus turned downwards (a). When this third Sort of spongy Bones is found, the middle one of the three in each Nostril is the largest, and the lowest is the smallest.—Besides all these, there are often several other small Bones standing out into the Nostrils, that, from their Shape, might also deserve the Name of Turbinata, but are uncertain in their Bulk, Situation, and Number (b).

The Names of these Bones sufficiently declare their spongy Substance, which has no firm ex-

ternal Plate covering it.

to.

re;

the

te.

ind

he-

ew.

ck-

tar

ın

of

the

Pa-

efs,

red

the

eri-

ape

up-

ges Pa-

ge,

or,

of ifes

t of

ach ril,

06-

They are joined to the Ossa maxillaria, palati and unguis in old Subjects, by a firm Union of Substance; and as this happens also frequently in People of no great Age, some (c) are of Opinion, that they should be esteemed Part of the Palate-bones; others (d) think, that since their upper Edge is continued by a Plate to a Part of the Os ethmoides, they ought to be esteemed to be a Part of this Bone.

Their Use is, to straiten the Nostrils, to afford a large Surface for extending the Organ of Smelling, to cover Part of the Antra maxillaria, and to assist in forming the Under-part of the lacrymal Ducts, the Orifices of which into

the Nose are concealed by these Bones.

The Offa turbinata are nearly complete in a new-born Infant.

VOMER,

(b) Santorin, Observat. Anatomic. cap. 5. § 9.

(c) Id. ibid. cap. 5. § 7.

⁽a) Cowper in Drake's Anthropolog. Book 3. Chap. 10.

⁽d) Hunauld. in Memoires de l'Acad. des Sciences, 1730.

VOMER, or Bone refembling a Plough. Share, is the thirteenth of the Upper Jaw, with out a Fellow, forming the Lower and Back. parts of the Partition of the Nose (a).

The Figure of this Bone is an irregular Rhomboid.—Its Sides are flat and fmooth.— Its posterior Edge appears in an oblique Direction at the Back-part of the Nostrils. The upper one is firmly united to the Base of the Sphenoid Bone, and to the Nasal Plate of the ethmoid; and, when it can be got separated, is hollow, for receiving the Processus azygos of the Sphenoid.—The anterior Edge has a long Furrow in it, where the middle Cartilage of the Nose enters. The lower Edge is firmly united to the Nasal Spines of the maxillary and Palate Bones. These Edges of this Bone are much thicker than its Middle, which is as thin as the finest Paper; by which, and the firm Union or Connexion this Bone has above and below, it can very feldom be separated entire in Adults: But, in a Child, it is much more easily separated entire; wherefore I shall examine all its Parts of such a Subject.

Its Situation is not always perpendicular, but often inclined and bended to one Side, as well as the Nasal Plate of the ethmoid Bone.

The Vomer is convex at its upper Part, and then is streight, as it is extended downwards and forwards where it is composed of two Plates; the Edges of which have a great Number of small Processes, disposed somewhat like the Teeth of a Saw, but more irregularly, and several

b

B

by Pa

⁽a) Columb. de re Anat. lib. 1. cap. 8. - Fallop. Observat.

rh-

th.

ck-

lar

ec-

he

he

he

is

of

ng

of

nly

nd

ne

as

rm

ind

ire

ore

xa.

but

rell

ind

rds

es;

of

the

fe-

ral

vat-

veral of them are reflected back. Between these Plates a deep Fossa is left, which, so far as the Top of the Curvature, is wide, and has strong Sides, for receiving the Processus azygos of the fphenoid Bone. Beyond the Arch forwards, the Fossa is narrower and shallower gradually to the Point of the Bone, receiving for some Way the Nasal Lamella ethmoidea; which. after the Offification is compleat, is fo closely united to the Vomer by the little Processes piercing into its Substance, as to prevent any Separation: on which account it has been efteemed by some (a) a Part of the ethmoid Bone. The middle Cartilage of the Nofe fills up what remains of the Fossa at its Fore-part. The posterior Edge of the Vomer, which appears above the Back-part of the Palate-bones, is broader above; but as it descends forwards, becomes thinner, though it is still folid and firm. -The lower Edge of this Bone, which refts on the Nafal Spine of the Palate and maxillary Bones, has a little Furrow on each Side of a small middle Ridge, answering to the Spines of the Bones of different Sides, and the Interstice between them. This Edge, and the upper one meet in the pointed Fore-end of this Bone.

The Body of the Vomer has a smooth Surface, and solid, but thin Substance; and towards its Sides, where it is thickest, some Cancelli may

be observed, when the Bone is broken.

It is joined above to the *fphenoid* and *ethmoid* Bones, and to the middle Cartilage of the Nose, by *Schindylesis*;—below, to the maxillary and Palate Bones, by the *fpinous* Suture.

The

⁽a) Lieutaud. Essais Anatomiques 1. fest. l'Os ethmoide.

The Vomer divides the Nostrils, enlarges the Organ of Smelling, by allowing Place for expanding the Membrane of the Nose on its Sides, and sustains the Palate Plates of the maxillary and palate Bones, which otherwise might be in Hazard of being pressed into the Nostrils; while the Vomer is secured from shuffling to one Side or other by the double Schindyiesis, by which it is joined to the Bones above and below.

These then are all the Bones which compose the Upper Jaw, except the Teeth, which are so much a-kin to those of the lower Jaw, that I chuse to make one Description serve for both, in which the Disserences observable in them shall be remarked, after the second Part of the Face, the Lower Jaw, is examined: because the Structure of the Teeth cannot be well understood, until the Case in which they are set is explained.

MAXILLA INFERIOR *, the Lower law, confifts only of one moveable Bone, and

fixteen Teeth incased into it.

This Bone, which is somewhat of the Figure of the Greek Letter v, is situated at the lower Part of the Face, so as its convex middle Part is forwards, and its Legs are stretched back. It is commonly divided into the Chin, Sides, and Processes.—The Chin is the middle Forepart, the Extent of which to each Side is marked on the external Surface by the Holes observable there, and internally by the Beginning of an oblique Ridge.—Beyond these the Sides appears.

^{*} Tivos, ordyw. Mandibula, facies.

pear, and are continued till the Bone, by bending upwards, begins to form the Processes.

On the Fore-part of the Chin, a transverse Ridge appears in the Middle, on each Side of which the Musculi quadrati, or Depressores labii inferioris, and the levatores labii inferioris, depress the Bone: And below these Prints, a small Rising may be observed, where the Depressores commence. — On the Back-part of the Chin, sometimes three, always two, small Protuberances appear in the Middle. To the uppermost, when it is seen, the Franum of the Tongue is connected. From the middle one, the Musculi genioglossi rise; and from the lowest, the Geniobyoidei have their Origin. Below the last, we see two rough Sinuosities formed by the digastric Muscles.

At the Lower and Fore-part of the external Surface of each Side of the Lower Jaw, a small Eminence may be observed, where the Depressor labiorum communis rises. Near the upper Edge of the Side a Ridge runs Length-ways, to which the under Part of the Musculus Buccinator is connected.—Internally, towards the upper Edge of each Side, another Ridge appears from which the Mylolyoidei have their Origin, and to which the internal Membrane of the

Gums adheres.

the

ex.

its

na-

oht

ils;

one

by

be-

ole

are

hat

oth,

lem

the

ule

un-

set

wer

and

gure

wer

Part

ack.

des,

ore.

ark-

erv-

g of

ap.

eat,

In the upper Edge of both Chin and Sides are a great many deep Pits or Sockets, for receiving the Roots of the Teeth. The Number and Magnitude of these Sockets are various, because of the different Number, as well of the Teeth themselves, as of their Roots, in different People. These Sockets in this Lower Jaw,

as well as in the Upper one, are less deep as old Age comes on; when freed from the Teeth by any Means, they are some time after filled up with an offeous Net-work, which at last becomes entirely folid, and as smooth as any other Part of the Bone; fo that in a great many old Jaws one cannot observe a Vestige of the Sockets: But then the Taw becomes less, and much nar. rower (a).—Hence we may know why the Chin and Nose of edentulous People are much near. er than before the Teeth were loft; while their Lips either fall in towards the Mouth, or stand prominent forwards. - When new Teeth are protruded, new Sockets are formed (b). — The lower Edge of the Chin and Sides is smooth and equal, and is commonly called the Base of the Lower Jaw .- The Ends of the Base, where the Jaw turns upwards, are called its Angles; the external Surface of each of which has feveral Inequalities upon it, where the Maffeter Muscle is inferted; as the internal Surface also has, where the Pterygoideus internus is inferted, and a Ligament extended from the styloid Process of the temporal Bone is fixed.

The Processes are two on each Side.—The anterior sharp thin coronoid ones have the crotaphite Muscles inserted into them—The posterior Processes or Condyles * terminate in an oblong smooth Head, supported by a Cervix. The Heads whose greatest length is transverse, and whose Convexity is turned forwards, are tipped with a Cartilage, as the articulated Parts

· (a) Vefal. Anat. lib. 1. cap. 10.

⁽b) Fallop. Observ. Anat. * Articulatorii.

of all other moved Bones are. - The Fore-part of the Root and Neck of these condyloid Processes are a little hollow and rough, where the external Pterygoid Muscles are inserted.

old

by

up

mes art

aws ets:

nar-

hin

ear.

heir

and

are

The

and

the

the

the

eral

ascle

has,

and

fs of

The

cro-

The

e 111

TUX.

erie,

are

Parts

Of

The Holes of the Lower Jaw are two on each Side; one at the Root of the Processes internally, where a large Branch of the third Branch of the fifth Pair of Nerves enters with an Artery, and a Vein returns. small sharp Process frequently jutts out backwards from the Edge at the Fore-part of this Hole, to which a Ligament extended from the temporal Bone is fixed (a), which faves the Nerve and Vessels from being too much pressed by the pterygoid Muscles. - From the lower Side of this Hole, either a small superficial Canal or a Furrow descends, where a Branch of the Nerve is lodged, in its Way to the Mylo-hyoideus Muscle and sublingual Gland (b). -The other Hole is external, at the Confines of the Chin, where Branches of the Nerve and Vessels come out.—The Canal betwixt these two Holes is formed in the middle of the Substance of the Bone, and is pierced by a great Number of small Holes by which the Nerves and Blood-vessels of the Cancelli and Teeth pass. This Canal is continued a little further than the external Hole at the Chin. — On account of the Veffels and Nerves in the Lower Jaw, Fractures of it may be attended with dangerous Symptoms.

The Surface of the lower Jaw is hard and firm, except at the spongy Sockets, where however

⁽¹⁾ Weitbrecht, Syndesmolog. Fig. 32. 1. (b) Palfyn. Anat. Chirur, traite 5. chap. 6.

however it is stronger than the Upper Jaw.—
Its internal Substance is cellular, without any solid Partition between the Cancelli in its Middle.—— At the Base, especially of the Chin, where this Bone is most exposed to Injuries, the solid Sides of it are thick, compact, and hard.

The lower Jaw generally receives the Roots of fixteen Teeth into its Sockets, by Gompholis: and its condyloid Processes, covered with Cartilage, are articulated with the temporal Bones, in a Manner that is not commonly described right: For, as was already mentioned in the Description of the temporal Bones, not only the Fore-part of the Cavity between the zygomatic, auditory and vaginal Processes, but also the adjoining Tubercle at the Root of the zygomatic Process of each Os temporum is covered with a smooth Cartilage, for this Articulation .-Here also an intermediate moveable Cartilage is placed, which being thin in the Middle, and thick at the Edges, is concave on both Sides; and is connected fo firmly by Ligaments to each Condyle, as to follow the Motions of the Condyle; and so loosely to the temporal Bone, as readily to change its Situation from the Cavity to the Tubercle, and to return again; while the common Ligament of the Articulation affords Space enough for such a Change of Place backwards and forwards; but, like other Ligaments of the Joints by Ginglimus, is strong and short at the Sides, to confine the lateral Motions.

When therefore the Teeth of both Jaws coincide, the Condyles are lodged fecurely in

he

ny

id-

in,

he

1.

ots

is:

ti-

es.

ed

he

he

tic,

ad.

tic

1 1

ige

le,

oth

ga-

10-

the

ua-

to

. of

uch

ds;

by

to

IWS

in

the

the temporal Cavities, but their Motions to either Side must be confined both by the Firmnels of the Ligaments, and the rifing Brims which are on each Side of the Cavities. -When the Jaw is brought directly forwards, the Condyles and intermediate Cartilages defeend and advance forwards upon the Tubercles. — In this Situation the lateral Motions are a little more free than in the former one, from the want of rifing Brims to stop the Condyles. - When the Fore-teeth of the Lower Jaw are moved forwards, and to a Side, the Condyle of the opposite Side is either advanced from the Cavity to the Tubercle, while the Condyle of the fame Side remains in the Cavity; or if both Condyles are on the Tubercles, when the Jaw is moved obliquely to a Side, the Condyle of the Side to which the Motion is made, flides back from the Tubercle to the Cavity.— When the Mouth is opened by the Descent of the Lower Jaw, the Fore-part of it, where the depressing Muscles are fixed, is drawn backwards, as well as downwards, while Resistance is made to the Angles moving backwards by the Maffeter and internal pterygoid Muscles, and, at the same time, the external Pterygoid draw the Condyles and their moveable Cartilages forwards; and therefore, when the Mouth is opened, the Condyles are carried forwards upon the Tubercles, and the Axis of Motion of the Bone is a little above its Angles. But in this Situation there is less Resistance, than in any other, to the Condyles luxating forwards; a Difease which seldom happens, except when People are gaping too wide; and therefore the N 2 common

Jaw of Infants when they are yawning, is reafonable.—In chewing there is a Succession of

the Motions above described. (a).

Here a general Remark may be made, That where-ever moveable Cartilages are found in Joints, either the articulated Bones are of fuch a Figure, or so joined and fixed by their Ligaments, that little Motion would be allowed without such Cartilages; or else some Motions are necessary to the right Use of the Member, which the Form of the Articulation would not otherwise admit of. This will more fully appear after the other Joints with such Cartilages are described.

In a Child born to the full Time the lower Jaw is composed of two Bones, connected by a thin Cartilage in the Middle of the Chin, which gradually offises, and the two Bones intimately unite.——In each of these Bones there are five or six Sockets for Teeth as in the Upper

Taw.

After I have thus described the Incasement of the Teeth, the Insertion of so many Muscles of the Tongue, and of the Os hyoides, the Connexion of the Membrane of the Tongue to the maxillary Bone, and the Motions of this Bone; it is easy to see, that the Lower Jaw must be a principal Instrument in Munducation, Degluttion, and Speech.

The

⁽a) For a more full Account of this Articulation, vid. Edinburgh Medical Essays and Observ. Vol. 1. Art. 11. & Vol. 3. Art. 13.—Memoires de l'Acad. des Sciences, 1744.

The TEETH are the hard white Bodies placed in the Sockets of both Jaws. Their Number is generally fixteen above, and as many below; though some People have more, others have fewer.

The broad thick Part of each Tooth which appears without the Socket, is the Base or Body *.

The smaller Processes sunk into the Maxilto, are the Roots or Fangs, which become gradually smaller towards the End farthest from the Base, or are nearly conical, by which the Surface of their Sides divide the Pressure made on the Bases, to prevent the soft Parts, which are at the small Points of the Sockets, to be hurt by such Pressure.—At the Place where the Base ends, and the Roots begin, there is generally a small circular Depression, which some call the Neck or Collar.

Without the Gums the Teeth are covered with no Membrane, and they are faid to have no proper *Periosteum* within the Sockets; but that is supplied by the reflected Membrane of the Gums; which, after a good Injection, may be evidently seen in a young Subject, with the Vessels from it penetrating into the Substance of the Teeth; and it may be discovered in any Tooth recently pulled, by macerating it in Water (a). The Adhesion of this Membrane to these Roots is strengthned by the small Furrows observable on them.

Each Tooth is composed of its Cortex, or Enamel, and an internal bony Substance. The N 3 Cortex

the

ea.

of

hat

111

uch

iga-

wed

ions

ber,

not

ap-

ages.

19WC

by a

hich

late.

e are

pper

ment

uscles

Con-

o the

Bone;

t be a

gluti

The

vid. E.

1744

^{*} Corona.

⁽a) Cowper, Anatom. Explicat. Tab. 92. Fig. 7. lit. E.

Cortex has no Cavity, or Place for Marrow; and is fo folid and hard, that Saws or Files can with Difficulty make Impression on it. thickest upon the Base, and gradually, as the Roots turn smaller, becomes thinner, but not proportionally to the Difference of the Size of the Base and Roots.—The Fibres of this Ena. mel are all perpendicular to the internal Sub. stance, and are streight on the Base, but at the Sides are arched with the convex Part towards the Roots (a); which makes the Teeth refift the Compression of any hard Body between the Jaws, with less Danger of breaking these Fibres, than if they had been fituated transversely. The fpongy Sockets in which the Teeth are placed, likewise serve better to prevent such an Injury, than a more folid Base would have done.-Notwithstanding the great Hardness of this Cor. tex, it is wasted by Manducation. Hence the sharp Edges of some Teeth are blunted, and made broad, while the rough Surfaces of others are made smooth and flat, as People advance in Life.

The bony Part of the Teeth has its Fibres running streight, according to the Length of the Teeth. When it is exposed to the Air, by the breaking or falling off of the hard Cortex, it soon corrupts. And thence carious Teeth are often all hollow within, when a very small Hole appears only externally.

The Teeth have Canals formed in their Middle, wherein their Nerves and Blood-veffels are lodged; which they certainly need, being constanty

⁽a) Havers's Osteolog. nov. Dift. 1,

stantly wasted by the Attrition they are subjected to in Manducation, and for their further Growth, not only after they first appear, but even in Adults; as is evident when a Tooth is taken out: For then the opposite one becomes longer, and these on each Side of the empty Socket turn broader; so that when the Jaws are brought together, it is scarce observable

where the Tooth is wanting (a).

W;

can

t is

the

not

e of

na-

ub-

the

ards

efift

the

res,

The

ced.

ury,

Cor.

the

and

hers

e in

bres

h of by

rtex,

1 are Hole

Mid.

s are

con-

antly

The Vessels are easily traced so long as they are in the large Canal, but can scarce be obferved in their Distribution from that to the Substance of the Teeth of Adults. Ruysch (b) however affirms, that after Injection he could trace the Arteries into the hardest Part of the Teeth: And Lewenhoek (c) suspected the Fibres of the Cortex to be Vessels. In Children I have frequently injected the Vessels of the Teeth as far as their Base: And in such as are not entirely offified, one can with a lucky Injection fill fo many Vessels, as to make both the Outside and Infide of the cortical Part appear perfectly red. -This plentiful Supply of Vessels must expose the Teeth to the same Disorders that attack other vascular Parts; and such Teeth as have the greatest Number of Vessels, must have the most numerous Chances of being seized with these Diseases.

Every Root of each Tooth has such a distinct Canal, with Vessels and Nerves in it. These Canals in the Teeth with more than one Root, come nearer each other, as they approach the

Bafe

⁽a) Ingraf. de Tumor, cap. 1. p. 24. 25, 26.

⁽b) Thefaur. 10. num 27.

⁽c) Arcan. Natur. Continuat. Epist. p. 3.

Base of the Tooth; and at last are only separated by very thin Plates, which being generally incompleat, allow a Communication of all the Canals; and frequently one common Cavaty only appears within the Base, in which a pulpy Substance composed of Nerves and Vessels is lodged. The Condition therefore of the Nerves here, bears a strong Analogy to that of the cutaneous Nerves which serve for the Sensation of Touching.

The Entry of the Canals for these Vessels is a small Hole placed a little to a Side of the extreme Point of each Root; sometimes, especially in old People, this Hole is entirely closed up, and consequently the Nerves and Blood

vellels are destroyed (a).

The Teeth are seen for a considerable Time in Form of Mucus contained in a Membrane; afterwards a thin cortical Plate, and some sew offeous Layers appear within the Membrane, with a large Cavity silled with Mucus in the Middle; and gradually this exterior Shell turns thicker, the Cavity decreases, the Quantity of Mucus is lessened, and this Induration proceeds, till all the Body is formed; from which the Roots are afterwards produced.

In young Subjects, different Stamina or Rudiments of Teeth are to be observed. Those next the Gums hinder ordinarily the deeperseated ones from making their Way out, while these prevent the former from sending out Roots, or from entering deep into the bony Sockets of the Jaws; by which they come to

be less fixed.

Children

⁽a) De la Hire Histoire de l'Acad des Sciences, 1699.

Children are feldom born with Teeth; but at two Years of Age they have twenty; and their Number does not increase till they are about feven Years old, when the Teeth that first made their Way through the Gums, are thrust out by others that have been formed deeper in the law, and some more of the Teeth begin to discover themselves farther back in the Mouth. About fourteen Years of Age, some more of the first Crop are shed, and the Number is increafed.—This Shedding of the Teeth is of good Use; for if the first had remained, they would have stood at a great Distance one from another; because the Teeth are too hard in their outer Crust, to increase so fast as the Jaws do. Whereas both the fecond Layer, and the Teeth that come out late, meeting, while they are foft, with a confiderable Resistance to their Growth in Length, from those situated upon them, necessarily come out broad, and fit to make that close Guard to the Mouth +, which they now form.

The Teeth are joined to the Sockets by Gomphosis, and the Gums contribute to fix them
there; as is evident by the Teeth falling out
when the Gums are any Way destroyed, or
made too spongy; as in the Scurvy or Salivations: Whence some (a) class this Articulation

with the Syffarcosis.

epa.

ral.

all

avi-

pul.

is 13

rves

CU-

n of

ls is

ex-

eci-

ofed

ood-

ime

ne;

few

ane,

the

urns

y of

eds,

the

udi-

nole

per-

hile

out

Ony

e to

Iren

The Uses of the Teeth are to masticate our Aliment, and to assist us in the Pronunciation of several Letters.

Though

[†] φραγμός.
(a) Drake's Anthropolog. Book 4. Chap. 3.

Though the Teeth fo far agree in their Struc. ture, yet, because of some Things wherein they differ, they are generally divided into three Classes, viz. Incifores, Canini, and Molares.

The Incifores * are the four Fore-teeth in each Jaw, receiving their Name from their Of. fice of cutting our Aliment; for which they are excellently adapted, being each formed in. to a sharp-cutting Edge at their Base, by their Fore-fide turning inwards there, while they are floped down and hollowed behind +; fo that they have the Form of Wedges; and therefore their Power of acting must be considerably in. ereased. Seeing in the Action of the Incife. res, a perpendicular Compression is only neces. fary, without any lateral Motion, they are not fo firmly fixed in their Sockets as the other Teeth are, each having only one short Root, but that is broader from before backwards, than to either Side, to have the greatest Strength where it is exposed to the strongest Force applied to it (a).

11

t

F

tl

21

th

m

Ja

pl

ha

Fo

Ja

mo

mo

men

The Incifores of the Upper Jaw, especially the two middle ones ‡, are broader and longer generally than those of the Under Jaw.

In a new-born Infant, the outer Shell of the Body of these Teeth is only hardned.—Afterwards, when the Stamina of two Sets are formed, each has its own Socket, those nearest to the Gums being placed more forward, and the others

1 Duales.

Γελασίνοι, τομίκοι, διχασήρες, κτίνες, τομείς, πρωσδικ itas, Riforii, quaterii, primi, primores, anteriores, acuti, t 'Oxuioxos.

⁽a) Lettre sur l'Osteologie ascribed to DuVerney.

others are lodged farther back within the Jawbones.

Canini ||, from the Resemblance to Dogs Tulks, are one on each Side of the Incifores in each Jaw. The two in the Upper Jaw are called Eye-teeth, from the Communication of Nerves which is faid to be betwixt them and the Eyes.—The two in the Lower Jaw are named angular or Wike-teeth, because they support the Angles of the Mouth.

.

II

1.

ot

er

ls,

th

li.

lly

he

er-

m-

to

the

ers

8104

The Canini are broader, longer, and stronger, than the Incifores. - Their Bases are formed into a sharp Edge, as the Incifores are; only that the Edge rises into a Point in the Middle .-Each of them has generally but one long Root, tho' fometimes they have two (a). The Roots are crooked towards the End. — The Canini of the Upper Jaw are larger, longer, and with more crooked Roots, than those of the Under Jaw. — The Form of their Base is fit both for piercing and cutting, and the long crooked Root of each, makes it secure in the Socket.

The Canini of a Child are in much the same

Condition as the Incifores are.

The Dentes molares, or Grinders +, which have got their Name because they grind our food, are generally five in each Side of each Jaw; in all twenty. Their Bases are broader. more scabrous, and with a thinner cortical Substance, than the other Teeth. They have also more Roots, and as these Roots generally divaricate

Kurdovrec, Riforii, fractorii, collaterales, columellares. (a) Fauchard, Chirurgien Dentiste, chap. 1.

[†] Μυλίται, γόμιροι, μύλος, πλατάς, φρασήρες, maxillares, mensales, clavales, buccarum.

ricate from each other, the Partitions of the Sockets between them bear a large share of the great Pressure they suffer, and hinder it to all

on their Points (a).

The Base of the first Grinder has an Edge pointed in the Middle, on its Outside, resembling the Canini; from which it slopes inwards, till it rises again into a Point.——It has generally but one Root, which sometimes is long and crooked at its Point.

The fecond Dens molaris has two Points on its Base, rising near equally on its Out and Inside.—It has two Roots, either separate or run together, but shorter than the Root of the first.—These two anterior Grinders are much smaller than the three that are placed farther back in the Mouth.

The third and fourth are very broad in their Bases, with four or five Points standing out;

and they have three or more Roots.

The fifth, called commonly Dens fapientia*, from its coming through the Gums later than the other Grinders, has four Points on its Base, which is not so large as the Base of the third and fourth, and its Roots are less numerous.

The Incifores of the Upper Jaw being broader than those of the Lower Jaw, make the superior Grinders to be placed so much farther back than the Lower ones, that when they are brought together, by shutting the Mouth, the Points of the Grinders of the one Jaw enter into the De-

pressions

C

H

de

01

te

Čl:

Bo

⁽a) Lettre sur l'Osteologie.

Zuppovienpec, npaurnpec, oblivovoi, sensus, intellectus, se rotini, ætatem complentes, genuini, moderatores.

pressions of the opposite Grinders, and they are all equally applied to each other, notwithstand-

ing the Inequality of their Surface.

10

er

eck

ht

of

)e.

115

fe-

The numerous Roots of the Dentes molares prevent their loofening by the lateral Pressure they fuffer in grinding; and as the Sockets in the Upper Jaw are more spongy, and the Teeth are more liable, by their Situation, to fall out (a), the Grinders there have more numerous and more separated Roots than in the Lower Jaw (b). The Number however of the Roots of the Teeth of both Jaws is very uncertain; fometimes they are more, fometimes fewer; frequently feveral Roots are joined together; at other times they are all distinct. The Disposition of fuch as are distinct, is also various; for in some the Roots stand out streight, in others they separate, and in others again they are crooked inwards. When the Roots are united, we can still distinguish them, by remarking the Number of small Holes at their Points, which determine the Number of Roots each Tooth ought to be reckoned to have.

At the Time of Birth, only two Dentes molares in each Jaw have begun to offify, and that at little more than the Base, which has several sharp Points standing out from it.— The temporaneous Grinders are placed more directly upon the internal Set than the other two Classes are; sometimes there is a Piece of the Bone of the Jaws between the two Sets; in o-

ther

⁽a) Galen. de Offib. cap. 5.

⁽b) Fauchard. Chirurg. Dent. chap. 1.

ther Children, the two Sets have no Bone in terposed between them.

From what has been faid, the Answers to the

fil

Co

an

fta

co

gro

Wil

the

1

(b)

tha

Cla

Ope

17701

of '

dray

(5) ferv. Hi(l. (c)

man.

Dentil

V

I

following Queries may be given.

Why are Children subject to Salivation, Fe. vers, Convulsion, Vomiting, Purging, &c. when their Teeth are breeding or cutting the Gums.

Why in Children do the Dentes incifores first cut the Gums, the Canini next, and Molares

last?

Why do Children shed their Teeth?

Wherefore have these temporaneous Teeth

generally no Roots, or very small ones?

Why have these first Teeth sometimes Roots, and that more frequently in Teeth pulled by Art than in those which are shedded by Nature (a)?

Why do these Roots frequently come out-

wards through the Gums?

Whence come Butter or Buck Teeth?

How do these Teeth sometimes go into the natural Row with the others, after pulling a rotten Tooth near them?

How have some People got two Rows of

Teeth in one or both Jaws? (b)

Why do the Teeth of old People loofen, and

then drop out entire?

Whence arise the new Sets of Teeth which

feveral old People obtain (c)?

Why are not the Gums of toothless old People torn by the hard Sockets in chewing?

(a) Fauchard, Chirurgien Dentiste p. 7.

(c) Hoffman. in van. Horn. Microcosin.p. 38.

⁽b) Blaf. Comment. ad Vesling. syntagm. cap. 13.

Why are the Teeth insensible when slightly

filed or rasped?

How come they to be fensible of Heat or Cold, to be set on Edge by Acids, or to give an uneasy Sensation when gritty or sandy Substances are rubbed between them?

Why does a Person who has a pain'd Tooth

imagine it longer than any other?

What is the Reason of some Persons dying convulsed, upon rasping or filing down an overgrown Tooth (a)?

How do the Teeth break and moulder away without any Pain in some People and not in o-

thers.

.

t.

10

of

ch

e0-

hy

What Parts are affected in the Tooth-ach? What are the causes of the Tooth-ach?

May Worms be reckoned among these Causes (b)?

Why are the Dentes molares most subject to-

that Disease?

In what different Manners ought the feveral Classes of Teeth to be extracted when such an Operation is necessary?

Whence proceeds the violent obstinate Hemorrhagy which sometimes attends the drawing

of Teeth (c)?

Why is it more difficult and dangerous todraw the Eye-teeth than any other?

0 2

What

(a) Bartholin. Anat. reformat. lib. iv. cap. 12.

(b) Jacob. in Act. Hafn. Vol. 5. Obs. 108 — Pechlin. Obferv. Medic. lib. 2. Obs. 36.—Bartholin. Hist. medic. Cent. 3.

(c) Pare, livre 6. chap. 2. — Rolfinc. lib. 2. cap. 27. & 30. — Moebii fundam. Medicin. cap. 9. Ephemerid. German. Dec. 1. ann. 3. obf. 316. — Fauchard. Chirurg. Dentifte, Tom. 1. chap. 23. observ. 7.

What makes it impossible frequently to draw Grinders, without bringing away Part of the Jaw-bone with them, or breaking the Fangs?

Why do Teeth foon replaced after being extracted, become again fixed in the Sockets. (a)?

According to the Division made of the Skeleton, we should now proceed to the Description of the Trunk of the Body: But must first consider a Bone, which cannot well be said to belong to either the Head or the Trunk; nor is it immediately joined to any other, and therefore is very seldom preserved with Skeletons. However, it is generally described by Authors, after the Bones of the Face: In obedience therefore to the prevailing Method, I shall next examine the Structure of

The OS HYOIDES*, which is fituated horizontally between the Root of the Tongue, and the Larynx. It is properly enough named named Hyoides, from the Refemblance it bears to the Greek Letter u, and may, for a clearer Demonstration of its Structure, be distinguished

I

fe

lo

N

25

ro

la

pe

ea

into its Body, Cornua, and Appendices.

The Body is the middle broad Part, convex before, and hollow behind. — The convex Fore-part is divided into two, by a Ridge, into the Middle of which the Mylo-hyoidei, and into the Sides the Stylo-hyoidei Muscles are inferted. — Above the Ridge, the Bone is horizontal, but pitted in the Middle by the Infertion

(a) De la Motte Chirurgie, tome 1. chap. 4. obs. 2.-

^{*} Hypsyloides, Lambdoides, παρασάτη, φαρυγετερον, 0s gutturis, os linguæ, os morsus Adami, assessor, os laude, bicorne.

fertion of the two Genio-hyoidei Muscles, and a little hollowed more laterally by the Bassio-glossis.—Below the Ridge, it is convex, but a little flatted in the Middle by the Sterno-hyoidei, and pitted more externally by the Coraco-hyoidei.—The Concavity behind faces backwards and downwards to receive the Thyroid Cartilage, when the Larynx and the Os hyoides are pulled towards each other by the Action of the Sterno-hyoidei and Hyothyroidei Muscles; and to its upper Edge, the ligamentous Membranes of the Epiglottis, Tongue, and Thyroid Cartilage, are fixed.

e.

1-

re

e,

ed

rs

er

n-

d

11-

0-

n.

on

de,

The Cormua of the + Os byoides are stretched backwards from each Side of its Body, where often a small Furrow points out the former Separation; for in young Subjects, the Body and Cornuta are not one continued Substance, as they come afterwards to be in Adults, -These Cormua are not always streight, nor of equal Length; their two plain Surfaces stand obliquely, floping from above outwards and downwards. --- Into the external, the Cerato-glossus is interted above, and the Thyro-byoideus Muscle below; and to the one behind, the ligamentous Membrane of the Tongue and Larynx adheres. Each of the Cornua becomes gradually smaller, as it is extended from the Bale; but ends in a round Tubercle, from which a moveable Cartilage stands out, which is connected to the urper Process of the Cartilago Thyroidea.

Where the Body of the Os byoides joins on each Side with its Cornua, a finall Styliform

O 3 Proceis,

[†] Crura, Latera inferiora.

Process, called Appendix ‡, rises upwards and backwards, into which the Musculi Stylo-hyoidei alteri, and Part of the Hyo-glossi Muscles are fixed. From each of them a Ligament is sometimes extended to the Styloid Processes of the temporal Bones, to keep the Os hyoides from being drawn too much forwards or downwards. The Part of this Ligament next to these Processes sometimes forms into several Cartilages, which afterwards ossify in old People. Russia (a) says that he has seen this Ossification continued as far up as the styloid Processes, which were therefore joined to the Os hyoides by Auchylosis.

te

uI

is

be

gr

10

tu

th

Wa

It

m

fo

fai

loi

er

fer

are

dy

min

brat

The Substance of the Os byoides is cellular, but covered with a firm external Plate, which is of sufficient Strength to bear the Actions of

fo many Muscles as are inserted into it.

It is not articulated with any Bone of the Body, except by Means of the Muscles and Li-

gaments mentioned.

The Use of the Os hyoides, is to serve as a solid Lever for the Muscles to act with, in raising or depressing the Tongue and Larynx, or in enlarging and diminishing the Capacity of the Fauces.

At Birth, this Bone is in a cartilaginous State; excepting a small Point of Bone in the Middle of its Body, and in each of the Cormua.—
The Appendixes frequently remain cartilaginous many Years.

[†] Crura superiora, Latera superiora, Ossa granisormia.
(a) Advers. Anat. Dec. 3. § 9.

Of the TRUNK.

THE TRUNK confifts of the Spine, Pel-

1 vis, and Thorax.

The SPINE * is the long Pile of Bones extended from the Condyles of the Occiput, to the End of the Rump. It fomewhat refembles two unequal Pyramids joined in a common Base. It is not, however, streight; for its upper Part being drawn backwards by strong Muscles, it gradually advances forwards, to support the OE-sophagus, Vessels of the Head, &c. Then it turns backwards, to make Place enough for the Heart and Lungs. It is next bended forwards, to support the Viscera of the Abdomen. It afterwards turns backwards, for the Enlargement of the Pelvis. And, lastly, it is resected forwards, for sustaining the lowest great Gut.

The Spine is commonly divided into true and false Vertebræ; the former constituting the long upper Pyramid, which has its Base below, while the salse Vertebræ make the shorter low-

er Pyramid, whose Base is above.

The TRUE VERTEBRAE+ are the twenty four upper Bones of the Spine, on which the several Motions of the Trunk of our Bodies are performed; from which Use they have justly got their Name.

Each of these Vertebræ is composed of its Bo-

dy and Processes.

Ŝ

The

^{&#}x27;Paχις, νωτον, ακανθα, ίερα σύριγξ, σωκην, Tergum, Hominis Carina.

[†] Στροφάς, ερόφιγες, Spondyli, ossa orbiculata; ossa verte-brata, verticula.

The Body is the thick fpongy Fore-part, which is convex before, concave backwards, horizontal and plain in most of them above and below.—Numerous finall Holes, especially on the Fore and Back-part of their Surface, give Passage to their Vessels, and allow the Li. gaments to enter their Substance .- The Edges of the Body of each Vertebra are covered, e. specially at the Fore-part, with a Ring of Bone firmer and more folid than the Substance of the Body any where else. These Rings seem to be joined to the Vertebræ in the Form of Epiphy. fes, but are alledged by some (a) to be the Ligaments offified. They are of great Use in preventing the spongy Bodies from being broken in the Motions of the Trunk.

I

W

g

31

tl

of

bo

ár

tw

ga

th

er

cu

lo

to

Between the Bodies of each two adjoining Vertebræ, a Substance between the Nature of Ligament and Cartilages is interposed; which feems to confift of concentrical curve Fibres, when it is cut horizontally; but when it is divided perpendicularly, the Fibres appear oblique and decustating each other (b).—The outer Part of the intervertebral Ligaments is the most folid and hard; and they gradually become foster, till they are almost in the Form of a glairy Liquor in the Center; and therefore these Substances were not improperly called mucous Ligaments by the Ancients (c). external fibrous Part of each is capable of being greatly extended, and of being compressed into a very small Space, while the middle fluid Part

(a) Fallop. Observat. Anatom.

(c) Galen, de usu Part, lib. 12. cap. 16.

⁽b) Blancard, Anat. reform. cap. 32.—Weithrecht. Syndefmolog. Sect. 4. § 15.

ld

1-

es

e.

n

ot

h

1-

t.

e

2

9

e

1.

Part is incompressible, or nearly so; and the Parts of this Ligament between the Circumference and Center approach in their Properties to either, in Proportion to their more folid or The middle Point is more fluid Texture. therefore a Fulcrum, or Pivot, on which the Motion of a Ball and Socket may be made, with fuch a gradual yielding of the Substance of the Ligament, in which ever Direction our Spines are moved, as faves the Body from violent Shocks, and their dangerous Consequences (c). -This Ligamento-cartilaginous Substance is firmly fixed to the horizontal Surfaces of the Bodies of the Vertebræ, to connect them, in which it is affisted by a strong membranous Ligament, which lines all their concave Surface, and by still a stronger Ligament that covers all their anterior convex Surface.

We may lay it down as a general Rule, notwithstanding some Exceptions, That the Bodies of the Vertebræ are smaller, and more solid above, but as they descend, they appear larger and more spongy; and that the Cartilages between them are thick, and the surrounding Ligaments strong in Proportion to the Largeness of the Vertebræ, and to the Quantity of Motion they perform: By which Disposition, the greater Weight is supported on the broadest best secured Base, and the Middle of our Body is al-

lowed a large and fecure Motion.

From each Side of the Body of each Vertebra, a bony Bridge is produced backwards, and to a Side; from the posterior End of which,

one

⁽c) Medical Essays and Observ. Vol. V. Art. 18.

one flanting Process rises, and another descends; the smooth, and what is generally the flattest Side of each of these four Processes, which are called the oblique *, is covered with a smooth Cartilage; and the two lower ones of each Vertebra are sitted to, and articulated with the two upper or ascending oblique Processes of the Vertebra below, having their articular Ligaments sixed into the rough Line round their Edges.

From between the oblique Processes of each Side, the Vertebra is stretched out laterally in-

to a Process that is named Transverse.

From the Back-part of the Roots of the two oblique, and of the transverse Process of each Side, a broad oblique bony Plate is extended backwards, where these meet, the seventh Process of the Vertebræ takes its Rise, and stands out backwards: This being generally sharp-pointed and narrow-edged, has therefore been called Spinal Process; from which this whole Chain of Bones has got its Name.

Besides the common Ligament which lines all the internal Surface of the spinal Processes, as well as of the Bodies, there are particular Ligaments that connect the bony Bridges and Processes of the contiguous Vertebræ together.

The Substance of the Processes is considerably stronger and firmer, and has a thicker external Plate than the Bodies of the Vertebræ have.

The feven Processes form a Concavity at their Fore-part, which, joined to the one at the Back-part of the Bodies, makes a great Foramen, and the Foramina of all the Vertebra form

^{*} Articularorii, minimi.

form a long large Conduit *, for containing the fpinal Marrow.—In the upper and lower Edge of each lateral Bridge, there is a Notch. These are so adapted to each other in the contiguous Vertebræ, as to form a round Hole in each Side between each two Vertebræ, through which the Nerves that proceed from the spinal Marrow,

and its Blood-veffels, pass.

re

th

1.

OV

r.

its

ch

n-

vo. ch

led

ro:

rp-

een ole

all

as Li-

10-

era-

ter-

ve.

at

at

Fo-

bre

rm

The Articulations then of these true Vertebræ are plainly double; for their Bodies are joined by the interveening Cartilage above described, and their oblique Processes being tipped with Cartilages, are so connected by their Ligaments, as to allow a fmall Degree of Motion to all Sides. Hence it is evident, that their Center of Motion is altered in different Politions of the Trunk: For when we bow forwards, the upper moved Part bears entirely on the Bodies of the Vertebræ: If we bend back, the oblique Processes support the Weight: If we recline to one Side, we rest upon the oblique Processes of that Side and Part of the Bodies: If we stand erect, all the Bodies and oblique Processes have their Share in our Support.

Hence it follows, 1. That because the Joints, of which the Spine is composed, are so numerous, the spinal Marrow, Nerves, Blood-vessels, &c. are not liable to such Compression, and over-stretching in the Motion of the Trunk of the Body, as they would otherwise be, since several Vertebræ must be concerned in every Motion of the Spine; and therefore a very small Curvature is made at the Conjunction of any

two

^{*} Ίερα σύριγξ, σωλήν, Canalis.

two Vertebræ (a). 2. That an erect Posture is the furest and firmest, because the Surface of Contact of the Fulcra is largest, and the Weight is most perpendicular to them (b). 3. That the Muscles which move the Spine, act with great. er Force in bringing the Trunk to an erect Po. sture, than in drawing it to any other: For in bending forwards, backwards, or to a Side, the Muscles which perform any of these Actions are nearer the Center of Motion; confequently the Lever with which they act is shorter, than when the Center of Motion is on the Part of the Vertebra, opposite to that where these Mus. cles are inferted; which is the Case in raising the Trunk. This is extremely necessary; fince, in the Deflections of the Spine from a perpendicular Bearing, the Weight of the Body foor inclines it which Way we defign; whereas, in raising us erect, this great Weight must be more than counteracted. 4. In calculating the Force exerted by the Muscles which move the Spine, we should always make Allowance for the Action of the Cartilages between the Ver. tebræ, which, in every Motion from an ered Posture, must be stretched on one Side, and compressed on the other, to both which they resist; whereas, in raising the Trunk, these Cartilages affift by their springy Force (c). 5. We are hence naturally led into the Reason of our Height of Stature increasing in the Mornmg

10

C

11

al

(0

to

an

(a) Galen de usu Part. lib. 12. cap. 12.

⁽b) Paaw de Offib. Pars. 2. cap. 3.

(c) Borelli, de motu Animal. pars. 1. schol. ad Propos. 38.

—Parent. Histoire de l'Acad. des Sciences, 1702.

ing, and diminishing at Night (a): For the intermediate Cartilages of the Vertebræ being pressed all Day long by the Weight of our Body, become more compact and thin in the Evening; but when they are relieved from this Pressure in the Night, they again expand themselves to their former Thickness; and seeing the Bulk of any Part must vary according to the different Distension or Repletion of the Vesfels composing it, we may understand how we become taller after a plentiful Meal, and decrease after Fasting or Evacuations (b). From the different Articulations of the Bodies, and oblique Processes of the Vertebræ, and the different Strength of the Ligaments, it is plain, that they are formed fo as to allow much larger Motion forwards than backwards; this last being of much less Use, and might be dangerous, by overstretching the large Blood-vessels that are contiguous to the Bodies of the Vertebræ (c). 7. The intervertebral Cartilages shrivelling as they become more folid by Age, is the Cause why old People generally bow forwards, and cannot raise their Bodies to such an erect Posture as they had in their Youth.

The true Vertebræ serve to give us an erect Posture; to allow sufficient and secure Motion to the Head, Neck, and Trunk of the Body, and to support and defend the Bowels, and o-

ther foft Parts.

ht

It.

0.

ns

ly

an

of

g

11.

110

in.

he

he

10

7.

nd

ey

ele

of

n.

ng

58.

At the ordinary Time of Birth, each Vertebra confists of three bony Pieces, connected by P Carti-

⁽a) Waffe, Philosoph. Transact. Num. 383. Art. 1.

⁽b) Abbe Fontenu, Histoire de l'Acad. des Sciences, 1725.

⁽c) Galen. de usu Part. lib. 1. cap. 16.

Cartilages; to wit, the Body, which is not fully offified, and a long crooked Bone on each Side; on which we see a small Share of the bony Bridge, the oblique Processes compleat, the beginning transverse Processes, and the oblique Plate, but no spinal Process; so that the Teguments are in no Danger of being hurt by the sharp Ends of these spinal Processes, while a Child is in its bended Posture in the Womb,

nor while it is fqueezed in the Birth.

From this general Mechanism of the Spine, an Account is eafily deduced of all the different preternatural Curvatures which the Spine is capable of: For if one or more Vertebra, or their Cartilages, are of unequal Thickness in opposite Sides, the Spine must be reclined over to the thinner Side; which now fustaining the greatest Share of the Weight, must still be more compressed, consequently hindered from extending itself in Proportion to the other Side, which, being too much freed of its Burden, has Liberty to enjoy a luxuriant Growth. The Caufes on which fuch an Inequality of Thick ness in different Sides of the Vertebræ depends, may vary: For either it may be owing to an Over-diffension of the Vessels of one Side, and from thence a preternatural Increase of the Thickness of that Part; or, which more commonly is the Case, it may proceed from an Obstruction of the Vessels, by which the Application of proper Nourishment to the bony Substance is hindered, whether that Obstruction depends on the faulty Disposition of the Vessels or Fluids; or if it is produced by an unequal mechanical Pressure, occasioned by a paralytic Weak.

ul.

ich

the

at,

ob.

the

by

ile

ab,

ne,

ent

ca-

10

in

ver

the

be

om

de,

en,

he

ck-

ids,

and

the

m-

an

pli-

ub-

1011

Tels

jual

ytic

ak.

Weakness of the Muscles and Ligaments, or by a spasmodic Over-action of the Muscles on any Side of the Spine, or by People continuing long, or putting themselves frequently into any Posture declining from the erect one: In all these Cases one common Essect follows, to wit, the Vertebræ, or their Cartilages, or both, turn thick on that Side where the Vessels are free, and remain thin on the other Side where the Vessels are straitned or obstructed. — Whenever any morbid Curvature is thus made, a fecond Turn, but in an opposite Direction to the former, must be formed; both because the Muscles on the convex Side of the Spine being ftretched, must have a stronger natural Contraction to draw the Parts to which their Ends are fixed, and because the Patient makes Efforts to keep the Center of Gravity of the Body perpendicular to its Base, that the Muscles may be relieved from a constant violent contractile State, which always creates Uneafiness and Pain.

When once we understand how these crooked Spines are produced, there is little Dissiculty in forming a just Prognosis; and a proper Method of Cure may be easily contrived, which must vary as to the internal Medicines, according to the different Causes on which the Disease depends: But one general Indication must be pursued by Surgeons; which is, to counterast the bending Force, by increasing the Compression on the convex Part of the Curvature, and diminishing it on the concave Side. The Manner of executing which in particular Cases must be different, and requires a very accurate Examination

mination of the Circumstances both of the Difease and Patient. In many such Cases, I have found some simple Directions, as to Postures in which the Patient's Body should be kept, of

very great Advantage.

Though the true Vertebræ agree in the general Structure which I have hitherto described, yet because of several Specialities proper to a particular Number, they are commonly divided into three Classes, viz. Cervical, Dorsal, and Lumbar.

The Cervical + are the feven uppermost Ver. tebræ; which are distinguished from the rest by these Marks .- Their Bodies are smaller and more folid than any others, and flatted on the Fore-part, to make Way for the OE fophagu; or rather this flat Figure is owing to the Pref. fure of that Pipe, and to the Action of the longi Colli and anterior recti Muscles,are also flat behind, where small Processes rife, to which the internal Ligaments are fixed.— The upper Surface of the Body of each Vertebra is made hollow, by a flanting thin Process which is raifed on each Side: - The lower Surface is also excavated, but in a different Manner; for here the posterior Edge is raised a little, and the one before is produced a confide rable Way. Hence we fee how the Cartile ges between those Bones are firmly connected, and their Articulations are fecure.

The Cartilages between these Vertebra at thick, especially at their Fore-part; which is

QD¢

n

ta

[†] Tpaxnau, auxis, Colli.

ive

in

of

ge.

ed,

2

led

ind

er.

by

ind

the

U;

el-

he

ley

le,

te-

ver

n-

lit-

de.

la-

ed,

are

is

DC

one Reason why the Vertebræ advance forward as they descend, and have larger Motion.

The oblique Processes of these Bones of the Neck more justly deserve that Name, than those of any other Vertebræ. They are situated slanting; the upper ones having their smooth and almost slat Surfaces facing obliquely backwards and upwards, while the inserior oblique Processes have these Surfaces facing obliquely forwards and downwards.

The transverse Processes of these Vertebræ are framed in a different Manner from those of any other Bones of the Spine: For besides the common transverse Process rising from between the oblique Processes of each Side, there is a second one that comes out from the Side of the Body of each Vertebra; and these two Processes, after leaving a circular Hole for the Passage of the cervical Artery and Vein, unite, and are considerably hollowed at their apper Part, with rising Sides, to protect the Nerves that pass in the Hollow; and at last each Side terminates in an obtuse Point, for the Insertion of Muscles.

The spinal Processes of these cervical Bones stand nearly streight backwards, are shorter than those of any other Vertebræ, and are forked or double at their Ends; and hence allow a more convenient Insertion to Muscles.

The thick Cartilages between the Bodies of these cervical Vertebræ, the Obliquity of their oblique Processes, and the Shortness and horizontal Situation of their spinal Processes, all conspire to allow them large Motion.

P 3

The

The Holes between the bony cross Bridges, for the Passage of the Nerves from the Medula In Spinalis, have their largest Share formed in the lowest of the two Vertebræ, to which they are common.

So far most of the cervical Vertebræ agree; but they have some particular Differences, which oblige us to consider them separately.

The first, from its Use of supporting the Head, has the Name of Atlas +; and is also called Epistrophea, from the Motion it performs

on the fecond.

The Atlas, contrary to all the other Verte. bræ of the Spine, has no Body; but, instead of it, there is a hony Arch.—In the convex Fore-part of which, a small Rising appears, where the Mufculi longi colli are inferted; and, on each Side of this Protuberance, a small Cavity may be observed, where the Recti interni minores take their Rife.—The upper and lower Parts of the Arch are rough and unequal, where the Ligaments that connect this Vertebra to the Os occipitis, and to the second Vertebra are fixed. The Back-part of the Arch is concave, smooth, and covered with a Cartilage, in a recent Subject, to receive the Tooth-like Process of the fecond Vertebra. In a first Vertebra from which the second has been separated, this Hollow makes the Passage for the spinal Marrow to feem much larger than it really is: On each Side of it a small rough Sinuolity may be remarked, where the Ligaments going to the Sides of the Tooth-like Process of the following Vertebra

Vertebra are fastened; and on each Side, a small rough Protuberance and Depression is observable, where the transverse Ligament, which secures the Tooth-like Process in the Sinuosity, is fixed, and hinders that Process from injuring the Medulla Spinalis in the Flexions of the Head.

in

ey

e

10

e.

ad

S,

d,

ty

0.

er

re

10

¥-

e,

e-

19

W

n

The Atlas has as little spinal Process as Body; but, instead thereof, there is a large bony Arch, that the Muscles which pass over this Vertebra at that Place, might not be hurt in extending the Head. On the Back and Upper Part of this Arch there are two Depressions, where the Recti postici minores take their Rise; and at the lower Part are two other Sinuosities, into which the Ligaments which connect this

Bone to the following one are fixed. The superior oblique Processes of this Atlas are large, oblong, hollow, and more horizontal than in any other Vertebra. They rife more in their external than internal Brim; by which their Articulations with the Condyloid Processes of the Os occipitis are firmer.—Under the external Edge of each of these oblique Processes, is the Fossa, or deep open Channel, in which the vertebral Arteries make the circular Turn, as they are about to enter the great Foramen of the occipital Bone, and where the tenth Pair of Nerves goes out.—In several Bodies, I have feen this Fossa covered with Bone. -The inferior oblique Processes extending from within outwards and downwards, are large, concave, and circular. So that this Vertebra, contrary to the other fix, receives the Bones with

with which it is articulated both above and be. low.

The transverse Processes here are not much hollowed or forked, but are longer and larger than those of any other Vertebra colli, for the Origin and Insertion of several Muscles; of which those that serve to move this Vertebra on the second have a considerable Lever to ad with, because of the Distance of their Insertion from the Axis of Revolution.

The Hole for the *spinal Marrow* is larger in this than in any other *Vertebra*, not only on account of the *Marrow* being largest here, but also to prevent its being hurt by the Motions of this *Vertebra* on the second one.—This large Hole, and the long transverse Processes, make this the broadest *Vertebra* of the Neck.

The Condyles of the Os occipitis move forwards and backwards in the superior oblique Processes of this Vertebra; but from the Figure of the Bones forming these Joints, it appears, that very little Motion can here be allowed to either Side; and there must be still less circular Motion.

In new-born Children this Vertebra has only the two lateral Pieces offified; the Arch, which it has at its Fore-part instead of a Body, being

cartilaginous.

The fecond Vertebra co'li is called Dentata, from the Tooth-like Process on the upper Part of its Body. Some Authors call it Epistrophea, but improperly, since this Designation is only applicable to the first, which moves on this as on an Axis.

The

be.

uch

the

of

OH

act

ion

in

On

but

of

ge

or-

ue

311

rs,

to

lar

ly

ch

ng

ut

ea,

ly

he

The Body of this Vertebra is somewhat of a pyramidal Figure, being large, and produced downwards, especially at its Fore-fide, to enter into a Hollow of the Vertebra below; while the upper Part has a square Process with a small Point standing out from it. This it is that is imagined to refemble a Tooth *, and has given Name to the Vertebra. The Side of this Process, which plays in the Hollow of the anterior Arch of the first Vertebra, is convex, smooth, and covered with a Cartilage; and it is of the fame Form behind, for moving on the Ligament which is extended transversely from one rough Protuberance of the first Vertebra to the other, and is cartilaginous in the Middle, where this Tooth-like Process plays on it.—A Ligament likewise goes out in an oblique transverse Direction, from each Side of the Processus dentatus, to be fixed at its other End to the first Vertebra, and to the occipital Bone; and another Ligament rises up from near the Point of the Process to the Os occipitis.

The superior oblique Processes of this Vertebra dentata are large, circular, very nearly in an horizontal Position, and slightly convex, to be adapted to the inferior oblique Processes of the first Vertebra.—A moveable Cartilage is said by some Authors to be interposed between these oblique Processes of the first and second Vertebra; but I could never find it.— The inferior oblique Processes of this Vertebra dentata answer exactly to the Description given of those common to all the cervical Vertebra.

The

[·] Conoides, Pyrenoides, Odontoides.

The transverse Processes of the Vertebra den. tata are short, very little hollowed at their upper Part, and not forked at their Ends; and the Canals through which the cervical Arteries pass, are reflected outwards about the middle Substance of each Process; so that the Course of these Vessels may be directed towards the transverse Processes of the first Vertebra. — Had this Curvature of the Arteries been made in a Part so moveable as the Neck is, while they were not defended by a Bone, and fixed to that Bone, scarce a Motion could have been performed without the utmost Hazard of Compresfion, and a stop put to the Course of the Liquids, with all its Train of bad Confequences. we observe this same Mechanism several times made use of, when there is any Occasion for a fudden Curvature of a large Artery. This is the third remarkable Instance of it we have feen. The first was the Passage of the Carotides through the temporal Bones; and the fecond was that lately described in the vertebral Arteries, turning round the oblique Processes of the first Vertebra, to come at the great Foramen offw occipitis.

The spinal Process of this Vertebra dentata is thick, strong, and short, to give sufficient Origin to the Musculi recti majores, and obliqui interiores, and to prevent the Contusion of these and other Muscles in pulling the Head back.

This fecond Vertebra confifts at the Birth, of four Bony pieces: For, besides the three which I already mentioned as common to all the Vertebræ, the Tooth-like Process of this Bone is begun at this Time to be offssied in its Middle.

Middle, and is joined as an Appendix to the Body of the Bone.—Lest this Appendix be bended or displaced, Nurses ought to keep the Heads of new-born Children from falling too far backwards, by Stay-bands, or some such Means, till the Muscles attain Strength sufficient

to prevent that dangerous Motion.

le

ld

at

ce

es

2

ve

les

725

e3,

rft

is

ri-

in-

elè

th,

ree

all

his

its.

When we are acquainted with the Structure and Articulations of the first and second Vertebre, and know exactly the Strength and Connexion of their Ligaments, there is no Difficulty in understanding the Motions that are performed upon or by the first; though this Subject was formerly Matter of hot Dispute among fome of the greatest Anatomists (a). none of my Purpose at present to enter into a Detail of the Reasons advanced by either Part; but to explain the Fact, as any one may fee it, who will remove the Muscles, which, in a recent Subject, hinder the View of these two Joints, and then will turn the Head into all the different Positions it is capable of. The Head may then be feen to move forwards, and backwards on the first Vertebra, as was already said, while the Atlas performs the Circumgyratio upon the second Vertebra; the inferior oblique Processes of the first Vertebra shuffling easily in a circular Way on the superior oblique Processes of the second, and its Body or anterior Arch having a Rotation on the Tooth-like Process, by which the perpendicular Ligament that is ient from the Point of the Tooth-like Process to the occipital Bone is twisted, while the lateral Ligaments that fix the Processus dentatus

to

⁽a) See Eustach. de motu Capitis.

to the Sides of the first Vertebra, and to the Or occipitis, are very differently affected; for the one upon the Side towards which the Face is turned by the Circumgyratio, is much shortned and lax, while the opposite one is stretched and made tense, and, yielding at last no more, pre vents the Head from turning any further round on this Axis. So that these lateral Ligaments are the proper Moderators of the Circumgyratio of the Head here, which must be larger or fmaller, as these Ligaments are weaker or stronger, longer or shorter, and more or less capable of being stretched. Besides the Revolution on this Axis, the first Vertebra can move a finall Way to either Side; but is prevented from moving backwards and forwards, by its anterior Arch, and by the cross Ligament, which are both closely applied to the Tooth-like Motion forwards here would have Process. been of very bad Consequence, as it would have brought the Beginning of the Spinal Marrow upon the Point of the Tooth-like Process.

The rotatory Motion of the Head is of great Use to us on many Accounts, by allowing us to apply quickly our Organs of the Senses to Objects: And the Axis of Rotation was altogether proper to be at this Place; for if it had been at a greater Distance from the Head, the Weight of the Head, if it had at any time been removed from a perpendicular Bearing to the small very moveable Joint, and thereby had acquired a long Lever, would have broke the Ligaments at every Turn inconsiderately performed; or these Ligaments must have been somed much stronger than could have been connect-

W

m

de

ed to such small Bones. Neither could this circular Motion be performed on the first Vertebra without Danger, because the immoveable Part of the Medulla oblongata is so near, as, at each large Turn, the Beginning of the spinal Marrow would have been in danger of being twisted, and of suffering by the Compression this would have made on its tender Fibrils.

It is necessary to observe, that the lateral or moderator Ligaments confine so much the Motion of the first Vertebra upon the second, that, though this Joint may ferve us on feveral Oc. casions, yet we are often obliged to turn our Faces farther round, than could be done by this Joint alone, without the greatest danger of twisting the spinal Marrow too much, and also of luxating the oblique Processes; therefore, in large Turns of this kind, the Rotation is affifted by all the Vertebræ of the Neck and Loins: and if this is not sufficient, we employ most of the Joints of the lower Extremities. — This Combination of a great many Joints towards the Performance of one Motion, is also to be observed in several other Parts of the Body: notwithstanding such Motions being generally faid to be performed by some single Joint only.

The third Vertebra of the Neck is by some called Axis; but this Name is applied to it with much less Reason than to the second.

This third, and the three below, have nothing particular in their Structure; but all their Parts come under the general Description formerly given, each of them being larger as they

descend.

d

0

n

e-

t,

e

W

us b-

er

at ht

vall

ir-

d;

ed et-

ed

Q

The

The seventh * Vertebra of the Neck is near to the Form of those of the Back, having the Upper and Lower Surfaces of its Body less hollow than the others:—The oblique Processes are more perpendicular;—neither spinal nor transverse Processes are forked.—This seventh, and the sixth Vertebra of the Neck, have the Hole in each of their transverse Processes more frequently divided by a small cross Bridge, that goes between the cervical Vein and Artery, than any of the other Vertebræ.

The twelve Dorfal + may be distinguished from the other Vertebrae of the Spine by the

following Marks.

Their Bodies are of a middle Size, betwixt those of the Neck and Loins;—they are more convex before than either of the other two Sorts; and are flatted laterally by the Pressure of the Ribs, which are inserted into small Cavities formed in their Sides. This flatting on their Sides, which makes the Figure of these Vertebræ almost an half Oval, is of good Use; as it affords a firm Articulation to the Ribs, allows the Trachea arteria to divide at a small Angle, and the other large Vessels to run secure from the Action of the vital Organs.—These Bodies are more concave behind than any of the other two Classes.—Their Upper and Lower Surfaces are horizontal.

The Cartilages interposed between the Bodies of these Vertebrae are thinner than in any other of

^{*} Atlas quibusdam, maxi...., magna Vertebra, prominens. + Θωρακος, μεταρρένε, νώτε, ύποτραχήλιε, Antisterni, εξ. Ctoris, tergi.

of the true Vertebræ; and contribute to the Concavity of the Spine in the Thorax, by being

thinnest at their Fore-part.

10

n

d

e

0

15

3

1

9

The oblique Processes are placed almost perpendicular; the upper ones slanting but a little forwards, and the lower ones slanting as much backwards.—They have not as much Convexity or Concavity as is worth remarking.—Between the oblique Processes of opposite Sides, several sharp Processes stand out from the upper and lower Parts of the Plates, which join to form the spinal Process; into these sharp Processes strong Ligaments are fixed, for connecting the Vertebræ.

The transverse Processes of the dorsal Vertebræ are long, thicker at their Ends than in the Middle, and turned obliquely backwards; which may be owing to the Pressure of the Ribs, the Tubercles of which are inserted into a Depres-

fion near the Eud of these Processes.

The Spinal Processes are long, small-pointed, and sloping downwards and backwards; from their Upper and Back-part a Ridge rises, which is received by a small Channel in the Fore-part of the Spinal Process immediately above, which

is here connected to it by a Ligament.

The Conduit of the spinal Marrow is here more circular, but, corresponding to the Size of that Cord, is smaller than in any of the other Vertebræ; and a larger Share of the Holes in the bony Bridges, for the Transmission of the Nerves, is formed in the Vertebra above, than in the one below.

The Connexion of the dorfal Vertebræ to the Ribs, the Thinness of their Cartilages, the Q 2 erect

erect Situation of the oblique Processes, the Length, Sloping, and Connexion of the Spinal Processes, all contribute to restrain these Verte. bræ from much Motion, which might diffurb the Actions of the Heart and Lungs; and, in Consequence of the little Motion allowed here. the intervertebral Cartilages fooner shrivel, by becoming more folid: And therefore the first remarkable Curvature of the Spine observed, as People advance to old Age, is in the least stretched Vertebræ of the Back; or old People first become round-shouldered.

The Bodies of the four uppermost dorfal Vertebræ deviate from the Rule of the Vertebræ becoming larger as they descend; for the first of the four is the largest, and the other three below gradually become fmaller, to allow the Trachea and large Vessels to divide at smaller

Angles.

The two uppermost Vertebræ of the Back, instead of being very prominent forwards, are flatted by the Action of the Musculi longi colli

and recti majores.

The proportional Size of the two little Depressions in the Body of each Vertebra for receiving the Heads of the Ribs, feems to vary in the following Manner: The Depression on the Upper Edge of each Vertebra decreases as far down as the fourth, and after that increases.

The transverse Processes are longer in each lower Vertebra to the seventh or eighth, with their smooth Surfaces, for the Tubercles of the Ribs, facing gradually more downwards; but afterwards as they descend they become

thorter

d

horter, and the smooth Surfaces are directed

more upwards.

al

ln

V

ft

The spinous Processes of the Vertebræ of the Back become gradually longer and more slanting from the first, as far down as the eighth or ninth Vertebra; from which they manifestly turn shorter and more erect.

The first * Vertebra, besides an oblong Hollow in its lower Edge, that assists in forming the Cavity, wherein the second Rib is received, has the whole Cavity for the Head of the first

Rib formed in it.

The fecond has the Name of Axillary +, without any Thing particular in its Structure.

The eleventh ‡ often has the whole Cavity for the eleventh Rib in its Body, and wants the smooth Surface on each transverse Process.

The twelfth || always receives the whole Head of the last Rib, and has no smooth Surface on its transverse Processes, which are very short.—The smooth Surfaces of its inserior oblique Processes face outwards as the Lumbar do.—And we may say, in general, that the upper Vertebræ of the Back lose gradually their Resemblance to those of the Neck, and the lower ones come nearer to the Figure of the Lumbar.

The Articulation of the Vertebræ of the Back with the Ribs, shall be more particularly considered after the Ribs are described. Only it

Q3 may

^{*} Λοφία, Gutturalis.

⁺ Mar xalishp.

[†] Αρβεπής, in neutram partem inclinans.

[|] Διασφεήρ, præcingens.

may be proper now to remark, that the Ligaments which ferve that Articulation, affift in

connecting the Vertebræ.

The lowest Order of the true Vertebræ is the Lumbar *, which are five Bones, that may be distinguished from any others by these Marks: 1. Their Bodies, though of a circular Form at their Fore-part, are somewhat oblong from one Side to the other; which may be occasion. ed by the Pressure of the large Vessels, the Aor. ta and Cava, and of the Viscera. The Epiphy. fes on their Edges are larger, and therefore the upper and lower Surfaces of their Bodies are more concave, than in the Vertebræ of the Back. 2. The Cartilages between these Ver. tebræ are much the thickest of any, and render the Spine convex within the Abdomen, by their greatest Thickness being at their Fore: part. 3. The oblique Processes are strong and deep; those in opposite Sides being almost placed in parallel Planes; the superior, which are concave, facing inwards, and the convex inferior ones facing outwards: and therefore each of these Vertebræ receives the one above it, and is received by the one below; which is not so evident in the other two classes already described. 4. Their transverse Processes are fmall, long, and almost erect, for allowing large Motion to each Bone, and sufficient lafertion to Muscles, and for supporting and defending the internal Parts. 5. Betwixt the Roots of the fuperior oblique and transverse Processes, a small Protuberance may be observed, where

^{*} Coquas, iguas, Joiev, Renum, Lumborum.

where some of the Muscles that raise the Trunk of the Body are inferted. 6. Their spinal Processes are strong, streight, and horizontal, with broad flat Sides, and a narrow Edge above and below; this last being depressed on each Side by Muscles. And at the Root of these Edges, we fee rough Surfaces for fixing the Ligaments. 7. The Canal for the numerous Cords, called Cauda equina, into which the spinal Marrow divides, is rather larger in these Bones, than what contains that Marrow in the Vertebra of the Back. 8. The Holes for the Passage of the Nerves are more equally formed out of both the contiguous Vertebræ than in the other Classes; the upper one furnishes however the larger share of each Hole.

The thick Cartilages between these humbar Vertebræ, their deep oblique Processes, and their erect spinal Processes, are all fit for allowing large Motion; though it is not so great as what is performed in the Neck; which appears from comparing the Arches which the Head describes when moving on the Neck, or the Loins

only.

16

De

at

m

n-

1.

1-

18

11-

V

e-

1d

97

re

n-

he

rle d, re The lumbar Vertebræ as they descend, have their oblique Processes at a greater Distance from each other, and facing more backwards and forwards.

Both transverse and spinal Processes of the middlemost Vertebra of the Loins are longest and thickest; in the Vertebra above and below they are less: So that these Processes of the first *

and

[&]quot; Neggirns, tenalis,

and fifth † are the least, to prevent their striking on the Rios or Osfa Iliûm, or their bruising the Muscles in the Motions of the Spine.

The Epiphyses round the Edges of the Bodies of the lumbar Vertebræ are most raised in the two lowest, which consequently make them appear hollower in the middle than the others are.

The Body of the fifth Vertebra is rather thinner than that of the fourth. — The spinal Process of this fifth is smaller, and the oblique Processes face more backwards and forwards

than in any other humbar Vertebra.

After confidering the Structure of the particular Vertebrae, and their mutual Connexion. we may observe a solicitous Care taken that they shall not be disjoined, but with great Difficulty; for, besides being connected by strong Ligaments proportioned to the Forces which are to be refifted; their Bodies either enter fo into each other, as to prevent their being displaced any Way, as in the Vertebrae of the Neck; or they are proped on all Sides, as these of the Back are by the Ribs; or their Surfaces of Contact are so broad, as to render the Separation almost impracticable, as in the Loins; while the Depth and Articulation of the oblique Processes are exactly proportioned to the Quantity of Motion, which the other Parts of the Bones allow, or the Muscles can perform: Yet, as these oblique Processes are small, and therefore not capable of fo secure a Conjunction as the larger Bodies, they may fooner yield to a disjoining Force; but then their diflocation

2

be

gu

Ve

th

ba

the

Li

Bo

Ma

wh

⁺ Aoxanims, fulciens.

is not of near fo bad Consequence as the Separation of the Bodies would be: For, by the oblique Processes being dislocated, the Muscles, Ligaments, and *spinal* Marrow are indeed stretched; but this Marrow must be compressed, or entirely destroyed when the Body of the

Vertebra is removed out of its Place.

e

al

n,

1.

to

ny

ev

ick

ct

al-

he

les

of

al.

25

re-

on

to

on

is

The FALSE VERTEBRAE compose the under Pyramid of the Spine. They are diffinguished from the Bones already described justly enough by this Epithet of false; because, tho each Bone into which they can be divided in young People, resembles the true Vertebræ in Figure, yet none of them contribute to the Motion of the Trunk of the Body; they being intimately united to each other in Adults, except at their lower Part, where they are moveable; whence they are commonly divided into two Bones, Os sacrum, and Coccygis.

OS SACRUM*, is so called, from being offered in Sacrifice by the Ancients, or rather because of its Largeness in Respect of the other Vertebræ. — This Bone is of an irregular triangular Shape, broad above, narrow below, convex behind, for the advantageous Origin of the Muscles that move the Spine and Thigh backwards; and concave before, for enlarging the Cavity of the Pelvis. — Four transverse Lines of a Colour different from the rest of the Bone which are seen on its Fore-part, are the Marks of Division of the five different Bones of which it consists in young Persons.

The

^{* &}quot;Ιερον, επονδυλος μεγας Hippocrat. ὖποςπονδυλον, Oribas. Πλατυ, Latum, Os clunium, Clavium.

The Fore-part of the Os facrum, analogous to the Bodies of the true Vertebræ, is smooth and flat, to allow a larger Space for the contained Bowels, without any Danger of hurting them; or this flat Figure may be owing to the equal Pressure of these Bowels, particularly of the last Gut. - The back Part of it is almost streight, without so large a Cavity as the Ver. tebræ have; because the spinal Marrow, now se: parated into the Cauda equina, is small.-The Bridges between the Bodies and Processes of this Bone, are much thicker, and in Proportion shorter, than in the former Class of Bones. The Strength of these cross Bridges is very remarkable in the three upper Bones, and is well proportioned to the incumbent Weight of the Trunk of the Body, which thele Bridges sustain in a transverse, consequently at unfavourable, Situation, when the Body is ered

There are only two oblique Processes of the Officerum; one standing out on each Side from the upper Part of the first Bone.—Their plain erect Surfaces face backwards, and are articulated with the inferior oblique Processes of the last Vertebra of the Loins, to which each of these Processes is connected by a strong Ligament, which rises from a scabrous Cavity round their Roots, where mucilaginous Glands are also lodged.—Instead of the other oblique Processes of this Bone, four rough Tubercles are to be seen on each Side of its Surface behind, from which the Musculus sacer has its

0

11

0

n

fp

Origin.
The transverse Processes here are all grown together into one large strong oblong Process

on each Side; which, so far as it answers to the first three Bones, is very thick, and divided into two irregular Cavities, by a long perpendicular Ridge. — The foremost of the two Cavities has commonly a thin cartilaginous Skin covering it in the recent subject, and is adapted to the unequal Protuberance of the Os Iliúm, and a strong Ligament connects the Circumference of these Surfaces of the two Bones.—
The Cavity behind is divided by a transverse Ridge into two, where strong ligamentous Strings that go from this Bone to the Os Iliúm with a cellular Substance containing Mucus, are lodged.

The transverse Processes of the two last Bones of the Os facrum are much smaller than the former.—At their Back-part near their Edge, a Knob and oblong slat Surface give rise to two strong Ligaments which are extended to the Os Ischium; and are therefore called

facrosciatic.

1115

oth

ing

the

of

101

Zer.

fe:

1.0:

of

lges

nes,

nent

acle

y an

ect.

e Oi

the

lain

ticu-

the

h of

iga.

VITY

ands

ique

rcles

e be-

s its

[W0

oces

OR

The spinal Processes of the three uppermost Bones of the Os sacrum, appear short, sharp, and almost erect, while the two lower ones are open behind; and sometimes a little Knob is to be seen on the fourth, though generally it is bisurcated, without the two Legs meeting into a Spine; in which Condition also the first is often to be seen; and sometimes none of them meet, but leave a Sinus, or rather Fossa, instead of a Canal (a)—The Musculus latissimus and longissimus Dorsi, Sacrolumbalis, and Glutaeus maximus, have Part of their Origins from these spinal Processes.

The

⁽a) Verbeyen, Anat. Tract. 5. cap. 9.

The Canal between the Bodies and Processes of this Bone, for the Cauda equina, is triangu. lar; and becomes fmaller as it descends, as the Cauda also does .- Below the third Bone, this Passage is no more a compleat bony Canal, but is open behind; and is only there defended by a strong ligamentous Membrane stretched over it, which, with the Muscles that cover it, and are very prominent on each Side, is a fufficient Defence for the Bundle of Nerves within.

At the Root of each oblique Process of this Bone, the Notch is conspicuous by which, and fuch another in the last Vertebra of the Loins. a Passage is left for the twenty fourth spinal Nerve; and, in viewing the Os facrum, either before or behind, four large Holes appear in each Side, in much the fame Height, as where the Marks of the Union of its feveral Bones remain. Some of the largest Nerves of the Body, pass through the anterior Holes; and superficial Grooves running outwards from them in different Directions, shew the Course of these Nerves - From the Intervals of these Grooves, the Pyriformis Muscle chiefly rises. — The Holes in the back Part of the Bone are covered by Membranes which allow fmall Nerves to pais through them .- The two uppermost of these Holes, especially on the Fore-side, are the largest; and as the Bone descends, the Holes turn smaller. Sometimes a Notch is only formed at the lower Part in each Side of this Bone; and in other Subjects there is a Hole common to it and the Os Coccygis, through which the twenty ninth Pair of spinal Nerves passes; and frequently a bony Bridge is formed on the back Part

0

tl

to

111

fac

ad:

Lig

ru.

he his

but

by

ver

ind

ent

his

and

ns,

nal her

in

ere

re-

dy,

ici-

dif-

refe

ves,

The

red

pals

refe

the

oles

rmne;

non

the

and

ack

Part

Part of each Side by a Process sent up from the back Part of the Os Coccygis, and joined to the little Knobs which the last Bone of the Os sacrum has instead of a Spinal Process. Under this Bridge, or Jugum, the twenty ninth Pair of spinal Nerves runs in its Course to the common Holes just now described:

The upper Part of the Body of the first Bone resembles the Vertebrae of the Loins; but the small fifth Bone is oblong tran versely, and hollow in the Middle of its lower Surface.

The Substance of the Os sacrum is very spongy, without any considerable solid external Plates, and is lighter proportionally to its Bulk than any other Bone in the Body; but is secured from Injuries by the thick Muscles that cover it behind, and by the strong ligamentous Membranes that closely adhere to it. —— As this is one of the most remarkable Instances of this Sort of Defence afforded a soft weak Bone, we may make the general Observation, That, where-ever we meet with such a Bone, one or other, or both these Defences are made use of; the sirst to ward off Injuries, and the second to keep the Substance of the Bone from yielding too easily.

This Bone is articulated above to the last Vertebra of the Loins, in the Manner that the Lumbar Vertebræ are joined; and therefore the same Motions may be performed here.—
The Articulation of the lower Part of the Os facrum to the Os Coccygis seems well enough adapted for allowing considerable Motion to this last Bone, was it not much confined by Ligaments.—Laterally, the Os facrum is joined

R

to the Ossa Iliam by an immoveable Synchondrosis, or what almost deserves the Name of a Suture; for the cartilaginous Crust on the Surface of the Bones is very thin, and both their Surfaces are so scabrous and unequal, as to be indented into each other; which makes such a strong Connexion, that great Force is required to separate them, after all the Muscles and Ligaments are cut. — Frequently the two Bones grow together in old Subjects.

The Uses of the Os facrum are, to serve as the common Base and support of the Trunk of the Body, to guard the Nerves proceeding from the End of the spinal Marrow, to defend the back Part of the Pelvis, and to afford sufficient Origin to the Muscles which move the Trunk

and Thigh.

The Bones that compose the Os facrum of Infants, have their Bodies separated from each other by a thick Cartilage; and, in the same Manner as the true Vertebræ, each of them consists of a Body and two lateral Plates, connected together by Cartilages; the Ends of the Plates seldom being contiguous behind.

OS COCCYGIS*, or Rump-bone, is that triangular Chain of Bones depending from the Os facrum; each Bone becoming smaller as they descend, till the last ends almost in a Point. The Os Coccygis is convex behind, and concave before; from which crooked pyramidal Figure which was thought to resemble a Cuckow's Beak, it has got its Name.

This

Ci

e

th

fo

fc Sl

ea

Pa

No

ead

nal Bo oft

^{* &#}x27;Offonuyiov, bffos, Caudæ os, Spondylium, Os Cuculi.

This Bone confifts of four Pieces in People of middle Age: - In Children, very near the whole of it is Cartilage: In old Subjects, all the Bones are united, and become frequently

one continued Bone with the Os facrum.

ľ

e

.

IS

of

n

e

It. k

0-

n-

10

·1-

10

as

2

nd lal

IC-

is

The highest of the four Bones is the largest. with Shoulders extended farther to each Side than the End of the Os facrum; which Enlargement should, in my Opinion, serve as a distinguishing Mark to fix the Limits of either Bone; and therefore should take away all Dispute about reckoning the Number of Bones, of which one or other of these two Parts of the falle Vertebræ, is composed; which Dispute must still be kept up, so long as the numbering five or fix Bones in the Os facition depends upon the uncertain Accident of this broad-shouldered little Bone being united to, or separated from it. The upper Surface of this Bone is a little hollow. — From the Back of that bulbous Part called its Shoulders, a Process often rises up on each Side, to join with the bifurcated Spine of the fourth and fifth Bones of the Os faction, to form the bony Bridge mentioned in the Description of the Os sacrum. - Sometimes these Shoulders are joined to the Sides of the fifth Bone of the Os sacrum, to form the Hole in each Side common to these two Bones, for the Paffage of the twenty ninth Pair of spinal Nerves. — Immediately below the Shoulders of the Os Coccygis, a Notch may be remarked in each Side, where the thirtieth Pair of the spinal Nerves passes. — The lower End of this Bone is formed into a small Head, which very often is hollow in the Middle.

R 2

The

The three lower Bones gradually become smaller, and are very spongy; but are strengthened by a strong Ligament which covers and connects them.— Their Ends, by which they are articulated, are formed in the same Man-

ner as those of the first Bone are.

Between each of these sour Bones of young Subjects a Cartilage is interpoled; therefore their Articulation is analogous to that of the Bodies of the Vertebræ of the Neck: For, as has been above remarked, the lower End of the Os facrum, and of each of the three superior Bones of the Os Coccygis has a small Depression in the Middle; and the upper Part of all the Bones of the Os Coccygis is a little concave, and consequently the interposed Cartilages are thickest in the Middle, to fill up both Cavities; by which they connect the Bones more firmly.-When the Cartilages offify, the upper End of each Bone is formed into a Cavity, exactly a dapted to the protuberant lower End of the Bone immediately above. - From this fort of Articulation, it is evident, that, unless when these Bones grow together, all of them are a pable of Motion; of which, the first and second, especially this last, enjoys the largest Share.

The lower End of the fourth Bone terminates in a rough Point, to which a Cartilage is ap-

di

110

pi

31

tic

The

pended.

To the Sides of these Bones of the Os Cocques, the Coccygæi Muscles (a), and Part of the Levatores Ani, and of the Glutaei maximi, are fixed.

⁽a) Douglas, Myograph. chap. 40. - Eustach. Tab. 34.

The Substance of these Bones is very spongy, and in Children cartilaginous; there being only a Part of the first Bone offissed in a new-born Infant.—Since therefore the *Intestinum rectum* of Children is not so firmly supported as it is in Adults, this may be one reason why they are more subject to a Procidentia Ani than old Peo-

ple (a).

h.

nd

ey

n-

ng

re

he

as

he

or

On.

he

ind

ck.

by

of

1 2.

the

of

hen

ca.

nd,

ates

ap-

occi.

the

are

The

b. 36,

From the Description of this Bone, we see how little it refembles the Vertebræ, fince it feldom has Processes, never has any Cavity for the spinal Marrow, nor Holes for the Passage of Nerves. - Its Connexion hinders it to be moved to either Side; and its Motion backwards and forwards is much confined: Yet, as its Ligaments can be stretched by a considerable Force, it is a great Advantage in the Excretion of the Fæces alvinæ, and much more in Childbearing, that this Bone should remain moveable; and the right Management of it, in delivering Women, may be of great Benefit to them (b). — The Mobility of the Os Coccygis diminishing as People advance in Age, especially when its Ligaments and Cartilages have not been kept flexible by being stretched, is probably one Reason why the Women, who are old Maids before they marry, have generally hard labour in Child-bed.

The Os Coccygis ferves to sustain the Intestimum rectum; and, in order to perform this Office more effectually, it is made to turn with a R 3 Curve

(b) Paaw, ibid .- Deventer, Operat. Chirurg. cap. 27.

⁽a) Spigel. de humani corp. fabric. lib. 2. cap. 22.—
Panw, de Offib. pars 2. cap. 3.

Curve forwards, by which also the Bone itself, as well as the Muscles and Teguments, is preferved from any Injury, when we fit with our

Body reclined back.

The fecond Part of the Trunk of the Skeleton, the PELVIS, is the cylindrical Cavity at the lower Part of the Abdomen, formed by the Os facrum, Os Coccygis, and Offa innominata; which last therefore fall now in Course to be examined.

Though the Name of OSSA INNOMI. NATA* contributes nothing to the Know. ledge of their Situation, Structure, or Office; yet they have been fo long and univerfally known by it, that there is no Occasion for changing it.—They are two large broad Bones, which form the Fore-part and Sides of the Pelvis, and the lower Part of the Sides of the Abdomen. - In Children each of these Bones is evidently divided into three; which are afterwards to intimately united, that scarce the least Mark of their former Separation remains: This notwithstanding, they are described as consisting each of three Bones, to wit, the Os Ilium, Ischium, and Pubis; which I shall first describe separately, and then shall consider what is common to any two of them, or to all the three.

OS ILIUM; or Haunch bone, is fituated highest of the three, and reaches as far down as one third of the great Cavity, into which the

Head of the Thigh-bone is received.

The

il

21

21

25

m

an

OU

te

m

is :

fro

cro

its

fou

Bel

Tra

65

^{*} Zxelluv, προςφίσεις, Sacro conjuncta.

† Λαγόνων, κενίου, Scaphium, lumbare, clupium, clavium, anchas.

The external Side of this Bone is unequally convex, and is called its Dorfum; - the internal concave Surface is by fome (but improperly) named its Costa. The semicircular Edge at the highest Part of this Bone, which is tipped with a Cartilage in the recent Subject, is named the Spine, into which the external, or descending oblique Muscle of the Abdomen, is inferted; and from it the internal ascending oblique and the transverse Muscles of the Belly, with the Glutæus maximus, Quadratus lumborum, and latissimus Dorsi, have their Origin. Some (a) are of Opinion, that it is only the tendinous Crust of all these Muscles, and not a Cartilage, as commonly alledged, that covers this bony Edge. - The Ends of the Spine are more prominent than the Surface of the Bone below them; therefore are reckoned Processes .-From the anterior spinal Process, the Sartorius and Fascialis Muscles have their Rise, and the outer End of the doubled Tendon of the external oblique Muscle of the Abdomen, commonly called Fallopius's or Poupart's Ligament, is fixed to it. — The Infide of the posterior spinal Process, and of Part of the Spine forward from that, is made flat and rough where the Sacro-lumbalis and longiffimus Dorsi rise; and to its outside Ligaments, extended to the Os sacrum and transverse Processes of the fifth and fourth Vertebra of the Loins, are fixed (b).—— Below the anterior spinal Process another Protuberance

⁽a) Winstow, Exposition Anatomique du corps humain, Traité des Os frais, §. 95. (b) Weithrecht. Syndesmolog. Sect. 4. §. 39. 40. 46. 47.

tuberance stands out, which, by its Situation, may be distinguished from the former, by add. ing the Epithet of inferior, where the Mus. culus rectus Tibiæ has its Origin (a). - Betwixt these two anterior Processes the Bone is hollow. ed, where the Beginning of the Sartorius Mus. cle is lodged.—Below the posterior spinal Pro. cess, a fecond Protuberance of the Edge of this Bone is in like Manner observable, which is closely applied to the Os facrum. - Under this last Process a considerable large Niche is oh. fervable in the Os Iliûm; between the Sides of which and the strong Ligament that is stretched over from the Os facrum to the sharp-pointed Process of the Or Ischium of the recent Subject, a large Hole is formed, through which the Musculus-pyriformis, the great sciatic Nerve, and the posterior crural Vessels pass, and are protected from Compression.

t

B

te

ed

fit

fid

cri

Co

COI

Pu

Ato

bro

ed.

hin

per

hum

dow Ligati

The external broad Side or Dorfum of the Os Iliam is a little hollow towards the Forepart; farther back it is as much raised; then is considerably concave; and, lastly, it is convex. These Inequalities are occasioned by the Actions of the Muscles that are situated on this Surface.—From behind the uppermost of the two anterior spinal Processes, in such Bones as are strongly marked by the Muscles, a semicircular Ridge is extended to the hollow Passage of the sciatic Nerve. Between the Spine and this Ridge the Glutaus medius takes its Rise. Immediately from above the lowest of the anterior spinal Processes, a second Ridge

⁽a) Baker, Curf. Offeolog. demonstr. 3.

is stretched to the Niche. Between this and the former Ridge, the Glutaus minimus has its Origin.—On the Outside of the posterior spinal Processes, the Dorsum of the Os Iliam is stat and rough, where Part of the Musculus Glutaus maximus and pyriformis rises.—The lowest Part of this Bone is the thickest, and is formed into a large Cavity with high Brims, to assist in composing the great Acetabulum; which shall be considered, after all the three Bones that constitute the Os Innominatum are described.

The internal Surface of the Os Iliûm is concave in its broadest Fore-part, where the internal Iliac Muscle has its Origin, and some Share of the Intestimum Iliûm and Colon is lodged.—From this large Hollow a small Sinuosity is continued obliquely forwards, at the Inside of the anterior interior spinal Process, where Part of the Psoas and Iliacus Muscles, with the crural Vessels and Nerves, pass.—The large Concavity is bounded below by a sharp Ridge, which runs from behind forwards; and, being continued with such another Ridge of the Os Pubis, forms a Line of Partition between the Aldomen and Pelvis.—Into this Ridge the broad Tendon of the Psoas parvus is inserted.

All the internal Surface of the Os Ilium, behind this Ridge, is very unequal: For the upper Part is flat, but spongy, where the Sacrolumbalis and longissimus Dorsi rise.— Lower down there is a transverse Ridge, from which Ligaments go out to the Os sacrum.—Immediately below this Ridge the rough unequal Cadiately below this Ridge the rough unequal Cadiately

vities

vities and Prominences are placed, which are exactly adapted to those described on the Side of the Os facrum.——In the same Manner the upper Part of this rough Surface is porous, for the sirmer Adhesion of the ligamentous cellular Substance; while the lower Part is more solid, and covered with a thin cartilaginous Skin, for its immoveable Articulation with the Os sacrum.——From all the Circumference of this large unequal Surface, Ligaments are extended to the Os sacrum, to secure more sirmly the Conjunction of these Bones.

The Passages of the medullary Vessels are very conspicuous, both in the Dorsium and Costa of many Ossa Ilium; but in others they are in-

considerable.

The posterior and lower Parts of these Bones are thick; but they are generally exceeding thin and compact at their Middle, where they are exposed to the Actions of the Musculi glutai and Iliacus internus, and to the Pressure of the Bowels contained in the Belly.—The Substance of the Ossa Ilium is mostly cellular, except a thin external Table.

In a ripe Child the Spine of the Os Iliûm is cartilaginous, and is afterwards joined to the Bone in form of an Epiphyse.—The large lower End of this Bone is not compleatly offi-

fied.

dle Bulk between the two other Parts of the os innominatum, is fituated lowest of the three, and is of a very irregular Figure.——Its ex-

tent.

ter

dra

lun

for

gre

mu

gre

I

ûm

out

his of t

inte

Bet'the

pyri

and

Imn

forn

tor 1

the.

the ligar

fmal

Fibr

oute

mis c

Musc

vere

uppe

(a)

^{*} Coxæ, coxendicis, pixis.

tent might be marked by an horizontal Line drawn near through the Middle of the Acetabulum; for the upper bulbous Part of this Bone forms some less than the lower Half of that great Cavity, and the small Leg of it rises to much the same Height on the other Side of the great Hole common to this Bone and the Os Pubis.

From the upper thick Part of the Os Ischium a sharp Process, called by some spinous, stands. out backwards, from which chiefly the Muscuhis coccygæus and superior Gemellus, and Part of the Levator Ani, rise; and the anterior or internal Sacrosciatic Ligament is fixed to it .-Between the upper Part of this Ligament and the Bones, it was formerly observed that the pyriform Muscle, the posterior crural Vessels, and the Sciatic Nerve, pass out of the Pelvis .-Immediately below this Process, a Sinuofity is formed for the Tendon of the Musculus obturator intermus.—In a recent Subject, this Part of the Bone, which ferves as a Pulley on which the Obturator Muscle plays, is covered with a ligamentous Cartilage, that, by two or three small Ridges, points out the Interstices of the Fibres in the Tendon of this Muscle- The outer Surface of the Bone at the Root of this ipinous Process is made hollow by the Pyriformis or Iliacus extermis Muscle.

Below the Sinuosity for the Obturator Muscle, is the great Knob or Tuberosity, covered with Cartilage or Tendon (a).—The upper Part of the Tuberosity gives Rise to the inferior

⁽a) Winflow. Exposit. Anat. des Os frais, § 96.

inferior Gemellus Muscle. To a Ridge at the Infide of this, the external or posterior facrosciatic Ligament is so fixed, that between it, the internal Ligament and the Sinuofity of the Os Ischisten, a Passage is left for the inter. nal obturator Muscle. The upper thick fmooth Part of the Tuber, called by some its Dorsum, has two oblique Impressions on it. The inner one gives Origin to the long Head of the Biceps flexor Tibie and seminervosus Muscles, and the Semimembranosus rises from the exterior one, which reaches higher and nearer the Acetabulum than the other—The lower thinner more scabrous Part of the Knob which bends forwards, is also marked with two flat Surfaces, whereof the internal is what we lean upon in fitting, and the external gives Rife to the largest Head of the Triceps adductor femoris. Between the external Margin of the Tuberosity and the great Hole of the Or innominatum, frequently there is an obtule Ridge extended down from the Acetabulum, which gives Origin to the Quadratus femoris. -As the Tuber advances forwards, it becomes fmaller, and is rough, for the Origin of the Musculus transversalis and Erector Penis.— The finall Leg of it, which mounts upwards to join the Os Pubis, is rough and prominent at its Edge, where the two lower Heads of the Triceps or Quadriceps adductor femoris take their Rife.

The upper and back Part of the Os Ischium is broad and thick; but its lower and fore-Part is narrower and thinner.——Its Substance is of the Structure common to broad Bones.

The

le

T

ed

bec

to

gro

wai

Os ,

Os Mu

upp

with vide

of I

Ridg

paffin

Ners

belov

from

ward

The Os Ilium and Pubis of the same Side, are the only Bones which are contiguous to the

Os Ischium.

The Part of the Os Ischi'm which forms the Acetabulum, the spinous Process, the great Tuber, and the recurved Leg, are all cartilaginous at Birth.—The Tuber with part of the Leg or Process above it becomes an Epi-

physe before this Bone is fully formed.

The OS PUBIS*, or Share-bone, is the least of the three Parts of the Os innominatum, and is placed at the upper Fore-part of it. The thick largest Part of this Bone is employed in forming the Acetabulum; from which becoming much fmaller, it is stretched inwards to its fellow of the other Side, where again it grows larger, and fends a small Branch downwards to join the End of the small Leg of the Os Ischium. — The upper Fore-part of each Os Pubis is tuberous and rough where the Musculus rectus and pyramidalis are inferted. -From this a Ridge is extended along the upper Edge of the Bone, in a continued Line with fuch another of the Os Iliûm, which di vides the Abdomen and Pelvis. The Ligament of Fallopius is fixed to the internal End of this Ridge, and the fmooth Hollow below it is made by the Pfoas and Iliacus internus Muscles passing with the anterior crural Vessels and Nerves behind the Ligament.—Some way below the former Ridge, another is extended from the tuberous Part of the Os Pubis downwards and outwards towards the Acetabulum; between

^{* 23}ng, Pictinis, penis, pudibundum, fenestratum.

te

th

ch

ec

21

fe

T

th

of

in

ne

de

CO

bo

lef

mu

this

mo

wit

ject

bee

ing

ftitt

fore

Cav

between these two Ridges the Bone is hollow and fmooth, for lodging the Head of the Pectineus Muscle.-Immediately below, where the lower Ridge is to take the Turn downwards, a winding Niche is made, which is comprehended in the great Foramen of a Skeleton, but is formed into a Hole by a subtended Ligament in the recent Subject, for the Paffage of the posterior crural Nerve, an Artery, and a Vein. The internal End of the Or Pubis is rough and unequal, for the firmer Adhesion of the thick ligamentous Cartilage that connects it to its Fellow of the other Side: The Process which goes down from that to the Os Ischium is broad and rough before, where the Gracilis and upper Heads of the Triceps, or rather Quadriceps adductor femoris have their Origin.

The Substance of the Os Pubis is the same

as of other broad Bones.

Only a Part of the large End of this Bone is offified; and the whole Leg is cartilaginous

in a Child born at the full Time.

Betwixt the Os Ischium and Pubis a very large irregular Hole is left, which, from its Resemblance to a Door or Shield, has been called Thyroides. This Hole is all, except the Nitch for the posterior crural Nerve, silled up in a recent Subject with a strong ligamentous Membrane, that adheres very firmly to its Circumference. From this Membrane chiesly the two Obsuratores, external and internal, take their Rise.—The great Design of this Hole, besides rendering the bone lighter, is to allow a strong enough Origin to the obsurators.

their Bellies, that there may be no Danger of disturbing the Functions of the contained Viscera of the Pelvis by the Actions of the internal, nor of the external being bruised by the Thigh-bone, especially by its lesser Trochanter, in the Motions of the Thigh inwards: Both which Inconveniencies must have happened, had the Osa innominata been compleat here, and of sufficient Thickness and Strength to serve as the fixed Point of these Muscles.—

The Bowels sometimes make their Way throw the Nitch for the Vessels, at the upper Part of this thyroid Hole, and this causes a Hernia in this Place (a).

In the external Surface of the Osa innominata, near the Outside of the great Hole, a large deep Cavity is formed by all the three Bones conjunctly: For the Os Pubis constitutes about one Fifth; the Os Iliûm makes something less than two Fifths, and the Os Ischium as much more than two Fifths. The Brims of this Cavity are very high, and are still much more enlarged by the ligamentous Cartilage, with which they are tipped in a recent Subject. From this Form of the Cavity it has been called Acetabulum; and, for a distinguishing Character, the name of the Bone that constitutes the largest Share of it is added; therefore Acetabulum ossis Ischii * is the Name this Cavity commonly bears.—Round the Base

⁽a) Memoires de l'Acad. de Chirurgie, tom. 1. p. 709: 60:

^{*} Coxæ, Coxendicis.

of the Supercilia the Bone is rough and unequal. where the capfular Ligament of the Articulation is fixed .- The Brims at the Upper and Back part of the Acetabulum are much larger and higher than any where elfe; which is very necessary to prevent the Head of the Fenner from slipping out of its Cavity at this Place, where the whole Weight of the Body bears upon it, and confequently would otherwise be constantly in danger of thrusting it out. -As these Brims are extended downwards and forwards, they become less; and at their internal lower Part a Breach is made in them: from the one Side of which to the other, a Ligament is placed in the recent Subject; under which a large Hole is left, which contains a fatty cellular Substance and Vessels. The Reason of which Appearance has afforded Matter of Debate. To me it feems evidently contrived for allowing a larger Motion to the Thigh inwards: For if the bony Brims had been here continued, the Neck of the Thigh-bone must have struck upon them when the Thighs were brought across each other; which, in a large strong Motion this Way, would have endangered the Neck of the one Bone, or Brim of the other. Then the Vesfels which are distributed to the Joint may safely enter at the Sinuosity in the Bottom of the Breach; which being however larger than is necessary for that Purpose, allows the large mucaginous Gland of the Joint to escape below the Ligament, when the Head of the Thigh-bone is in Hazard of preffing too much upon it in the Motions

in

th

til

Su

wh

Bo

the

COL

are

thic

0//2

no

con

or 1

Motions of the Thigh outwards (a)—Besides this Disserence in the Height of the Brims, the Acetabulum is otherwise unequal: For the lower internal Part of it is depressed below the cartilaginous Surface of the Upper-part, and is not covered with Cartilage; into the Upper-part, of this particular Depression where it is deepest, and of a semilunar Form, the Ligament of the Thighbone, which is commonly, though improperly, called the round one, is inserted; while in its more superficial lower Part the large mucilaginous Gland of this Joint is lodged. The largest Share of this separate Depression is formed in the Os Ischiam.

From what has been faid of the Condition of the three Bones composing this Acetabulum in new-born Children, it must be evident, that a considerable Part of this Cavity is car-

tilaginous in them.

The Offa innominata are joined at their Back. part to each Side of the Os facrum by a fort of Suture, with a very thin interveening Cartilage, which ferves as fo much Glue to cement thefe Bones together; and strong Ligaments go from the Circumference of this unequal Surface, to connect them more firmly. The Offa innominata are connected together at their Fore-part by the thick firm Cartilage interposed between the two Offa Pubis. These Bones can therefore have no Motion in a natural State, except what is common to the Trunk of the Body, or to the Os facrum. But it has been disputed, whether or not they loofen fo much from each other, and. S. 3"

⁽a) Petit, Memoires de l'Acad, des Sciences, 1722.

and from the Os facrum in Child-birth, by the Flow of Mucus to the Pelvis, and by the Throws of the Labour, as that the Offa Pubis recede from each other, and thereby allow the Passage between the Bones to be enlarged. - Several Ob. fervations (a) shew that this Relaxation some. times happens: But those who had frequently Opportunities of diffecting the Bodies of Women who died immediately after being delivered of Children, teach us to beware of regarding this as the common Effect of Child-birth; for they found such a Relaxation in very few of the Bodies which they examined (b).

Confidering what great Weight is supported in our erect Posture, by the Articulation of the Offa innominata with the Os facrum, there is great Reason to think, that if the conglutinated Surfaces of these Bones were once separated, (without which, the Offa Pubis cannot shuffle on each other) the Ligaments would be violently fretched, if not torn; from whence many Dif-

orders would arise (c).

Each Os innominatum affords a Socket (the Acetabulum) for the Thigh-bones to move in, and the Trunk of the Body tolls here on the Heads of the Thigh-bones. This Articulation is to be more fully described after the Offa femoris are examined.

The

W

fti

an

be

jec

by

na

M

ed

tha

it,

120

Ute

on

hay Ne

tha

Th

its

ope

and

Baf beh

it a

nois that

than

of i the

(a)

⁽a) Bauhin. Theat. Anat. lib. 1. cap. 49. - Spigel. Anat. lib. 2. cap. 24.—Riolan. Anthropogr. lib. 6. cap. 12.—Die-merbrocck, Anat. lib. 9. cap. 16.

⁽b) Hildan. Epist. Cent. obs. 45 .- Dionis. Sixieme Demonstr. des Os .- Morgagn. Advers. 3. Auimad. 15.

⁽b) Ludov, in Ephem, German, Dec. 1, ann. 3. obs. 254.

The Pelvis then has a large Open above where it is continued with the Abdomen, is strongly senced by Bones on the Sides, Back, and Fore-part, and appears with a wide Opening below, in the Skeleton; but, in the recent Subject, a considerable Part of the Opening is filled by the Sacrosciatic Ligaments, Pyrisorm, internal obturator, Levatores ani, Gemini and Coccygaei Muscles, which support and protect the contained Parts better than Bones could have done; so that Space is only left at the lowest Part of it, for the large Excretories, the Vesica urinaria, Intestinum rectum, and, in Females, the

Uterus, to discharge themselves.

The THORAX +, or Cheft, which is the only Part of the Trunk of the Body which we have not yet described reaches from below the Neck to the Belly; and, by means of the Bones that guard it, is formed into a large Cavity: The Figure of which is fomewhat conoidal; but its upper smaller End is not finished; being left open for the Passage of the Wind-pipe, Gullet, and large Blood-vessels; and its lower Part, or Base, has no Bones, and is shorter before than behind; fo that, to carry on our Comparison, it appears like an oblique Section of the Conoid. Besides which we ought also to remark, that the lower Part of this Cavity is narrower than some Way above (a); and that the Middle of its Back-part is confiderably diminished by the Bones standing forwards into it.

The

[†] Pectus caffum.

⁽a) Albin. de Offib. § 169.

The Bones which form the *Thorax* are the twelve dorfal *Vertebræ* behind, the Ribs on the Sides, and the *Sternum* before.

The Vertebræ have already been described as Part of the Spine; and therefore are now to be

passed.

The RIBS, or Costa t, (as if they were Custodes, or Guards, to these principal Organs of the animal Machine, the Heart and Lungs, are the long crooked Bones placed at the Side of the Chest, in an oblique Direction downwards in respect of the Back-bone. Their Number is generally twelve on each Side; tho' frequently eleven or thirteen have been found I never faw fewer or more than the ordinary Number, but in the Skeleton of a Boy, a bout eight Years old, now in my Possession; the fourth and fifth Ribs of the left Side are grown together at their Roots for near an Inch, and, afterwards dividing, have the same Appearance as the Ribs of the opposite Side, which are naturally formed.

The Ribs are all concave internally; where they are also made smooth by the Action of the contained Parts, which, on this Account, are in no Danger of being hurt by them; and they are convex externally, that they might result that Part of the Pressure of the Atmosphere, which is not bellanced by the Air within the Lungs, during Inspiration.—The Ends of the Ribs next the Vertebræ are rounder than they are after these Bones have advanced forwards, when they become flatter and broader, and have

‡ Плепраї, терігерта, от ввац

At Head Ridge lowes most the B Ridge

lages.

on the

mucila Head,

21

ro

in of

th

ter

up

rou

its

the

the

Foff

the

Veff

the]

to fe

plair

grea

the I

than

* K.

an upper and lower Edge, each of which is made rough by the Action of the intercostal Muscles, inserted into them. These Muscles, being all of nearly equal Force, and equally stretched in the Interstices of the Ribs, prevent the broken Ends of these Bones in a Fracture from being removed far out of their natural Place, to interrupt the Motion of the vital Organs. --- The upper Edge of the Ribs is more obtuse and rounder than the lower, which is depressed on its internal Side by a long Fossa, for lodging the intercostal Vessels and Nerves; on each Side of which there is a Ridge, to which the intercostal Muscles are fixed. Fossa is not observable however at either End of the Ribs; for at the Posterior or Root, the Vessels have not yet reached the Ribs; and, at the Fore-end, they are split away into Branches, to serve the Parts between the Ribs: Which plainly teaches Surgeons one Reason of the greater Safety of performing the Operation of the Empyema towards the Sides of the Thorax, than either near the Back or the Breast.

At the posterior End * of each Rib, a little Head is formed, which is divided by a middle Ridge into two plain or hollow Surfaces; the lowest of which is the broadest and deepest in most of them. The two Plains are joined to the Bodies of two different Vertebræ, and the Ridge forces itself into the interveening Cartilages.—A little Way from this Head, we find on the external Surface a small Cavity, where mucilaginous Glands are lodged; and round the Head, the Bone appears spongy, where the cap-

9

^{*} Karioy, Remulus.

to

th

Ca

car

the

ob

Th

to

wit

obl

rab

Ela

into

face

cart

Th

the

Mo

gres

cart

of T

Vert

Cav

larg

in t

bra .-

latio

Mot

dow

(a) (b)

T

fular Ligament of the Articulation is fixed.---Immediately beyond this a flatted Tubercle rifes, with a fmall Cavity at, and Roughness about its Root, for the Articulation of the Rib with the transverse Process of the lowest of the two Ver. tebræ, with the Bodies of which the Head of the Rib is joined.—Advancing further on this external Surface, we observe in most of the Ribs another smaller Tubercle, into which Ligaments which connect the Ribs to each other, and to the transverse Processes of the Vertebra and Portions of the longiffimus Dorsi, are inferted .---- Beyond this the Ribs are made flat by the Sacro-hembalis Muscle, which is inserted into the Part of this flat Surface farthest from the Spine, where each Rib makes a confiderable Curve, called by some its Angle. — Then the Rib begins to turn broad, and continues fo to its anterior End *, which is hollow and spongy, for the Reception of, and firm Coalition with the Cartilage that runs thence to be inferted into the Stermum, or to be joined with some other Cartilages .- In Adults generally the Cavity at this End of the Ribs is smooth and polished on its Surface; by which the Articulation of the Cartilage with it has the Appearance of being defigned for Motion; but it has none.

The Substance of the Ribs is spongy, cellular, and only covered with a very thin external lamellated Surface, which increases in Thickness and Strength as it approaches the Vertebræ.

To the Fore-end of each Rib a long broad and strong Cartilage is fixed, and reaches thence

* Πλάτη, Palmula.

the

to the Stermum, or is joined to the Cartilage of the next Rib. This Course, however, is not in a streight Line with the Rib; for generally the Cartilages make a confiderable Curve, the concave Part of which is upwards; therefore, at their Infertion into the Stermum, they make an obtufe Angle above, and an acute one below.-These Cartilages are of such a Length, as never to allow the Ribs to come to a right Angle with the Spine; but they keep them fituated fo obliquely, as to make an Angle very confiderably obtuse above, till a Force exceeding the Elasticity of the Cartilages is applied.—These Cartilages, as all others, are firmer and harder internally, than they are on their external Surface; and fome times, in old People, all their middle Substance becomes bony, while a thin cartilaginous Lamella appears externally (a). The Offification however begins much oftner at the external Surface. The greatest alternate Motions of the Cartilages being made at their great Curvature, that Part remains frequently cartilaginous, after all the rest is offissed (b).

The Ribs then are articulated at each End, of which the one behind is doubly joined to the Vertebræ; for the Head is received into the Cavities of two Bodies of the Vertebræ, and the larger Tubercle is received into the Depression in the transverse Process of the lower Vertebra.—When one examines this double Articulation, he must immediately see, that no other Motion can here be allowed, than upwards and downwards; since the transverse Process hinders.

(a) Vefal. Lib. 1. cap. 19.

⁽b) Havers. Ofteolog. nov. Difc. 5. p. 289.

in

nu

to

the

wa

liq

its

Ste

gre

abo

Ri

lor

Th

the

firf

fev

dua

pla

Ril

bra fou

170

fro

fcei Bre

and

of

the Rib to be thrust back; the Resistance on the other Side of the Sterman, prevents the Rib's coming forward; and each of the two Joints with the other Parts attached, oppose its turning round. But then it is likewise as e. vident, that even the Motion upwards and downwards, can be but fmall in any one Rib at the Articulation itself. But as the Ribs advance forwards, the Distance from their Center of Motion increasing, the Motion must be larger; and it would be very conspicuous at their anterior Ends, were not they refifted there by the Cartilages, which yield fo little, that the principal Motion is performed by the middle Part of the Ribs, which turns outwards and upwards, and occasions the Twist remarkable in the long Ribs at the Place near their Fore-end where they are most resisted (a).

Connexion which most of the Ribs enjoy, as belonging to all of them; but must now consider the Specialities wherein any of them differ from the general Description given, or from

each other.

In viewing the Ribs from above downwards, their Figure is still streighter; the uppermost being the most crooked of any.—Their Obliquity in respect of the Spine increases as they descend; so that though their Distance from each other is very little different at their Back-part, yet at their Fore-ends the Distance, between the lower ones must increase.—In Consequence too of this increased Obliquity of the lower Ribs,

⁽a) Winstow. Memoires de l'Acad. des Sciences, 1720.

each of their Cartilages makes a greater Curve in its Progress from the Rib towards the Sternum: and the Tubercles, that are articulated to the transverse Processes of the Vertebræ, have their smooth Surfaces gradually facing more upwards .- The Ribs becoming thus more oblique, while the Stermum advances forwards in its Descent, makes the Distance between the Sternum and the anterior End of the lower Ribs greater than between the Stermin and the Ribs above; confequently the Cartilages of those Ribs that are joined to the Breast-bone are longer in the lower than in the higher ones ... These Cartilages are placed nearer to each other as the Ribs descend, which occasions the Curvature of the Cartilages to be greater.

The Length of the Ribs increases from the first or uppermost Rib, as far down as the seventh; and from that to the twelsth, as gradually diminishes. —— The superior of the two plain, or rather hollow Surfaces, by which the Ribs are articulated to the Bodies of the Vertebrae, gradually increases from the first to the sourth Rib, and is diminished after that in each lower Rib. —— The Distance of their Angles from the Heads always increases as they descend to the ninth, because of the greater Breadth of the Sacrolumbalis Muscle (a).

The Ribs are commonly divided into true

and false.

The true * Costæ are the seven upper ones of each Side, whose Cartilages are all gradually

⁽a) Winflow, Exposition Anatomique des Os Secs, §. 643.
• Γνησαι, Germanæ, legitimæ.

dually longer as the Ribs descend, and are joined to the Breast-bone; so that being pressed constantly between two Bones, they are flatted at both Ends, and are thicker, harder, and more liable to offify, than the other Cartilages, that are not subject to so much Pressure. These Ribs include the Heart and Lungs; and therefore are the proper or true Custodes of Life.

t

le

tw

ra

lit

on

wa

is l

Mi

po

of

col

are

cep

ed.

Rit

(6

Offib

The five inferior Ribs of each Side are the false or BASTARD +, whose Cartilages do not reach to the Stermin; and therefore, wanting the Resistance at their Fore-part, they are there pointed; and, on this Account, having less Pressure, their Substance is softer. -The Cartilages of these false Ribs are shorter as the Ribs descend.—To all these five Ribs the circular Edge of the Diaphragm is connected; and its Fibres, instead of being stretched immmediately transversely, and so running perpendicular to the Ribs, are pressed fo as to be often, especially in Exspiration, parallel to the Plane in which the Ribs lie: Nay, one may judge by the Attachments which these Fibres have so frequently to the Sides of the Thorax, a confiderable Way above where their Extremities are inferted into the Ribs, and by the Situation of the Viscera, always to be observed in a dead Subject laid supine, that there is constantly a large Concavity formed on each Side by the Diaphragm within these bastard Ribs, in which the Stomach, Liver, Spleen, &c. are contained; which, being only reckoned

[†] Μαλθάκαι, χουδροδεις, άκανθαι, κλένες, ρόαι, adulterina, Puriæ, illegitimæ.

reckoned among the Viscera naturalia have occasioned the Name of Bastard Custodes to these Bones.

Hence in simple Fractures of the salse Ribs, without Fever, the Stomach ought to be kept moderately silled with Food, less the pendulous Ribs salling inwards, should thereby increase the Pain, Cough, &c. (a).—Hence likewise we may learn how to judge better of the Seat of several Diseases, and to do the Operation of the Empyema, and some others, with more Sasety than we can do, if we sollow the common Directions.

The eight upper Ribs were formerly (b) classed into Pairs, with particular Names to each two, to wit, the crooked, the folid, the pectoral, the twisted: But these Names are of so little Use, that they are now generally neglected.

The first Rib of each Side is so situated, that the stat Sides are above and below, while one Edge is placed inwards, and the other outwards, or nearly so; therefore sufficient Space is lest above it for the Subclavian Vessels and Muscle; and the broad concave Surface is opposed to the Lungs: But then, in Consequence of this Situation, the Channel for the intercostal Vessels is not to be found, and the Edges are differently formed from all the other, except the second; the lower one being rounded, and the other sharp.—The Head of this Rib is not divided into two plain Surfaces by

⁽a) Hippocrat de Articulo. § 51.---Parè, Lib. 15. cap. 11. (b) Laurent. Hist. Avat. lib. 2. cap. 29.-----Paaw. de Ossibus. Part. 3. cap. 2.

a middle Ridge, because it is only articulated with the sirst Vertebra of the Thorax.—
Its Cartilage is ossissed in Adults, and is united to the Stermum at right Angles.—Frequently this sirst Rib has a Ridge rising near the Middle of its posterior Edge, where one of the Heads of the Scalenus Muscle rises.—Farther forward it is statted, or sometimes depressed by the Clavicle.

The fifth, fixth, and feventh, or rather the fixth, feventh, eighth, and fometimes the fifth, fixth, feventh, eighth, ninth Ribs, have their Cartilages at least contiguous; and frequently they are joined to each other by cross Cartilages; and most commonly the Cartilages of the eighth, ninth, tenth, are connected to the former, and to each other by firm

Ligaments.

The eleventh, and sometimes the tenth Rib, has no Tubercle for its Articulation with the transverse Process of the Vertebra, to which it is only loosely fixed by Ligaments.—The Fossa in its lower Edge, is not so deep as in the upper Ribs, because the Vessels run more towards the Interstice between the Ribs.—Its Fore end is smaller than its Body, and its short small Cartilage is but loosely connected to the Cartilage of the Rib above.

The twelfth Rib is the shortest and straightest.—Its Head is only articulated with the last Vertebra of the Thorax; therefore is not divided into two Surfaces.—This Rib is not joined to the transverse Process of the Vertebra, and therefore has no Tubercle, being of

ten

ten

wh

cef

1101

fels

ver To

15 (

mo

tio

nev of

afti

lati

pai dre

vid

per

Th

19 1

ma

10

as 1

the

and

Tu

Con

lar

Suc

are

tenpulled necessarily inwards by the Diaphragm, which an Articulation with the transverse Process would not have allowed.—The Fossa is not found at its under Edge, because the Vessels run below it.—The Fore-part of this Rib is smaller than its Middle, and has only a very small-pointed Cartilage sixed to it.—To its whole internal Side the Diaphragm is connected.

The Motion and Uses of the Ribs shall be more particularly treated of, after the Descrip-

tion of the Stermin.

The Heads and Tubercles of the Ribs of a new-born Child have Cartilages on them; Part of which, becomes afterwards thin Epiphyses.

—The Bodies of the Ribs incroach gradually after Birth upon the Cartilages; so that the latter are proportionally shorter, when compared to the Ribs, in Adults, than in Children.

Here I cannot help remarking the wife Providence of our Creator, in preferving us from perishing as foon as we come into the World. The End of the Bones of the Limbs remain in a cartilaginous State after Birth, and are many Years before they are entirely united to the main Body of their feveral Bones; whereas the Condyles of the occipital Bone, and of the lower Jaw, are true original Processes, and offified before Birth, and the Heads and Tubercles of the Ribs are nearly in the fame Condition: and therefore the Weight of the large Head is firmly supported; the Actions of Sucking, Swallowing, Respiration, &c. which are indiffenfably necessary for us as foon as we come

by

Ca

for

ke

the

Te

tur

Ste

and

tha

lar

cia

fre

OVE

ced

tha

mu

W

giv

the

bei

of

me

rab

but

Danger of separating the Parts of the Bones that are most pressed on in these Motions: Whereas, had these Processes of the Head, Jaw, and Ribs, been Epiphyses at Birth, Children must have been exposed to Danger of dying by such a Separation; the immediate Consequences of which would be the Compression of the Beginning of the spinal Marrow, or want of Food,

or a Stop put to Respiration.

The STERNUM*, or Breaft-bone, is the broad flat Bone, or Pile of Bones, at the Fore-part of the Thorax. — The Number of Bones, which this should be divided into, has occasioned Debates among Anatomists, who have confidered it in Subjects of different Ages.—In Adults of a middle Age, it is composed of three Bones, which eafily separate after the Cartilages connecting them are destroyed. Frequently the two lower Bones are found intimately united; and very often in old People, the Stermum is a continued bony Substance from one End to the other; though we still observe two, sometimes three, transverse Lines on its Surface; which are Marks of the former Divisions.

When we consider the Sternum as one Bone, we find it broadest and thickest above, and becoming smaller as it descends. The internal Surface of this Bone is somewhat hollowed for enlarging the Thorax; but the Convexity on the external Surface is not so conspicuous, because the Sides are pressed outwards by

^{*} Trides, Os Pectoris, ensiforme, scutum cordis.

by the true Ribs; the round Heads of whose Cartilages are received into seven smooth Pits, formed in each side of the Sternum, and are kept firm there by strong Ligaments, which on the external Surface have a particular radiated Texture (a).—Frequently the cartilaginous Fibres thrust themselves into the bony Substance of the Sternum, and are joined by a Sort of Suture.—The Pits at the upper Part of the Sternum, are at the greatest Distance one from another, and, as they descend, are nearer; so

that the two lowest are contiguous.

The Substance of the Breast-bone is cellular, with a very thin external Plate, especially on its internal Surface, where we may frequently observe a cartilaginous Crust spread over it (b). On both Surfaces, however, a strong ligamentous Membrane is closely braced; and the Cells of this Bone are fo small, that a confiderable Quantity of osleous Fibres must be employed in the Composition of it: Whence, with the Defence which the Muscles give it, and the moveable Support it has from the Cartilages, it is sufficiently secured from being broken; for it is strong by its Quantity of Bone; its Parts are kept together by Ligaments; and it yields enough to elude confiderably the Violence offered (c).

So far may be faid of this Bone in general; but the three Bones, of which, according to

the

(b) Jac. Sylv. in Galen. de Offibus, eap. 12.

⁽a) Ruysch, Catalog. Rar. fig. 9.

⁽c) Senac. in Memoires de l'Acad. des Sciences, 1724.

the common Account, it is composed in Adults,

are each to be examined.

The first, all agree, is somewhat of the Figure of a Heart, as it is commonly painted; only it does not terminate in a sharp Point.

This is the uppermost thickest Part of the Sternum.

The upper Middle-part of this first Bone, where it is thickest, is hollowed, to make Place for the Trachea arteria; though this Cavity * is principally formed by the Bone being raised on each Side of it, partly by the Clavicles thrusting it inwards, and partly by the Sterno-mastoidei Muscles pulling it upwards. -- On the Outfide of each Tubercle, there is an oblong Cavity, that, in viewing it transversely from before backwards, appears a little convex: Into these Glenæ the Ends of the Clavicles are received. — Immediately below these, the Sides of this Bone begin to turn thinner; and in each a superficial Caviby or a rough Surface is to be feen, where the first Ribs are received or joined to the Stermum. In the Side of the under End of this first Bone, the half of the Pit for the fecond Rib on each Side is formed. - The upper Part of the Surface behind is covered with a strong Ligament, which secures the Clavicles; and is afterwards to be more particularly taken notice of.

The fecond or middle Division of this Bone, is much longer, narrower, and thinner, than the first; but, excepting that it is a little nar-

rower

10

OV

ne

for

an

ve

me

fro

Mi

Sid

the

of

tha

this

fery

tho

Car

offi

may

the

rece

two

for

it m

Ster

cart

Nan

+, t

calle

tum rioris,

^{*} Yearn, Jugulum, Furcula superior.

rower above than below, it is nearly equal all over in its Dimensions of Breadth or Thickness .- In the Sides of it are compleat Pits for the third, fourth, fifth, and fixth Ribs, and an half of the Pits for the fecond and feventh. The Lines, which are Marks of the former Division of this Bone, being extended from the Middle of the Pits of one Side to the Middle of the corresponding Pits of the other Side.—Near its Middle an unossified Part of the Bone is fometimes found, which, freed of the ligamentous Membrane or Cartilage that fills it, is described as a Hole; and in this Place, for the most Part, we may observe a transverse Line, which has made Authors divide this Bone into two. --- When the Cartilage between this and the first Bone is not offified, a manifest Motion of this upon the first may be observed in Respiration, or in raising the Stermin, by pulling the Ribs upwards in a recent Subject.

The third Bone is much less than the other two, and has only one half of the Pit for the seventh Rib formed in it; wherefore it might be reckoned only an Appendix of the Sternum. —— In young Subjects it is always cartilaginous, and is better known by the Name of Cartilago xiphoides or ensiformist, than any other; though the Antients often called the whole Sternum, Ensiforme, comparing

[†] Clypealis, gladialis, mucronata, malum, granatum, scutum stomachi, epiglottalis, cultralis, Medium Furculæ inferioris, scutiformis, ensiculata.

ring the two first Bones to the Handle, and this Appendix to the Blade of a Sword.— This third Bone is feldom of the fame Figure. Magnitude, or Situation in any two Subjects: for sometimes it is a plain triangular Bone, with one of the Angles below, and perpendicular to the Middle of the upper Side, by which it is connected to the fecond Bone. --In other People the Point is turned to one Side, or obliquely forwards or backwards.—Fre. quently it is all nearly of an equal Breadth, and in feveral Subjects it is bifurcated; whence fome Writers give it the Name of Furcella or Furcula inferior; or else it is unoffified in the Middle.—In the greatest Number of Adults it is offified, and tipped with a Cartilage; in fome one half of it is cartilaginous, and in 0thers it is all in a cartilaginous State. -- Generally several oblique Ligaments fixed at one End to the Cartilages of the Ribs, and by the other to the outer Surface of the xiphoid Bone, connect it firmly to those Cartilages (a).

So many different Ways this small Bone may be formed, without any Inconvenience: But then some of these Positions may be so directed, as to bring on a great Train of ill Consequences; particularly, when the lower End is ossissed, and is too much turned outwards or inwards (b), or when the Conjunc-

tion

tic

to

lef

fta

of

fin

of

gel

Po

dre

fon

till

feri

and

the

the

last

the

W

Mo

to e

Car

of 1

coff

Bones Obl.

(a) Weithrecht, Syndesmolog. p. 121.
(b) Rolfinc. Dissert. Anat. lib. 2. cap. 41.—Peaw, de Offib. part 1. cap. 3. & part 3. cap. 3.

tion of this Appendix with the second Bone is

too weak (c).

The Sternum is joined by Cartilages to the feven upper Ribs, unless when the first coalesces with it in an intimate Union of Substance; and its unequal Cavity on each Side of its upper End is fitted for the Ends of the Clavicles.

The Sternum most frequently has four round small Bones, surrounded with Cartilage, in Children born at the full Time; the uppermost of these, which is the first Bone, being the largest.——Two or three other very small bony Points are likewise to be seen in several Children.—The Number of Bones increases for some Years, and then diminishes but uncertainly, till they are at last united into those above described of an Adult.

The Uses of this Bone are, to afford Origin and Insertion to several Muscles; to sustain the Mediastinum, to defend the vital Organs, the Heart and Lungs, at the Fore-part; and, lastly, by serving as a moveable Fulcrum of the Ribs, to affist considerably in Respiration: Which Action, so far as it depends on the Motion of the Bones, we are now at Liberty to explain.

When the Ribs that are connected by their Cartilages to the Stermum, or to the Cartilages of the true Ribs, are acted upon by the intercostal Muscles, they must all be pulled from the

oblique

⁽c) Paaw. Ibid.—Berrich. Act. Hafn. vol. 5. Obs. 79.—Benet. Sejulchret. Anat. tom. 2. lib. 3. § 5. Append. ad Obs. 8. & ibid. § 7. Obs. 19.

b

ir

C

21

to

01

at

cl

tl

th

712

m

ar

ye

oblique Position which their Cartilages kept them in, nearer to right Angles with the Ver. tebræ and Sternum, because the first or uppe. most Rib is by much the most fixed of any; and the Cartilages making a great Resistance to raising the anterior Ends of the Ribs, their large arched middle Parts turn outwards as well as upwards. The Stermin, pressed strongly on both Sides by the Cartilages of the Ribs, is pushed forwards, and that at its several Parts. in Proportion to the Length and Motion of its Supporters, the Ribs; that is, most at its lower End.—The Stermum and the Cartilages, thus raised forwards, must draw the Diaphragm connected to them; consequently so far stretch it, and bring it nearer to a Plane. The Power that raises this Bone and the Cartilages, fixes them sufficiently to make them resist the Action of the Diaphragin, whose Fibres contract at the same Time, and thrust the Viscera of the Abdo. men downwards .- The arched Part of the Ribs being thus moved outwards, their anterior Ends and the Stermin being advanced forwards, and the Diaphragm being brought nearer to a plain Surface, instead of being greatly convex on each Side within each Cavity of the Thorax, it is evident how confiderably the Cavity, of which the nine or ten upper Ribs are the Sides, must be widened, and made deeper and long. er.—While this is doing in the upper Ribs, the lower ones, whose Cartilages are not joined to the Sternum or to other Cartilages, move very differently, though they conspire to the fame Intention, the Enlargement of the Thorax: For having no fixed Point to which their anterior

anterior Ends are fastened, and the Diaphragm being inferted into them at the Place where it runs pretty streight upwards from its Origin at the Vertebræ, these Ribs are drawn downwards by this strong Muscle, and by the Muscles of the Abdomen, which, at this Time, are refifting the stretching Force of the Bowels; while the intercostal Muscles are pulling them in the contrary Direction, to wit, upwards: The Effeet therefore of either of these Powers, which are Antagonists to each other, is very little, as to moving the Ribs either up or down; but the Muscles of the Abdomen, pushed at this Time outwards by the Viscera, carry these Ribs along with them. Thus the Thorax is not only not allowed to be shortened, but is really widened at its lower Part, to affift in making sufficient Space for the due Distension of the Lungs.

As foon as the Action of these several Muscles ceases, the elastic Cartilages extending themselves to their natural Situation, depress the upper Ribs, and the Stermum subsides;——the Diaphragm is thrust up by the Viscera abdominalia, and the oblique and transverse Muscles of the Belly serve to draw these Ribs inwards at the same time.——By these Causes, the Cavity of the Breast is diminished in all its Di-

menfions.

Though the Motions above described of the Ribs and Stermum, especially of the latter Bone, are so small in the mild Respiration of a healthy Person, that we can scarce observe them; yet they are manifest whenever we designedly increase our Respiration, or are obliged to do it after Exercise, and in several Diseases.

Of the Superior Extremities.

A Uthors are much divided in their Opinions about the Number of Bones which each fuperior Extremity + should be faid to confist of, some describing the Clavicle and Scapula as Part of it, others classing these two Bones with those of the Thorax: But since most Quadrupeds have no Clavicles, and the human Thorax can perform its Functions right when the Scapula is taken away (a), whereas it is impossible for us to have the right Use of our Arms without these Bones; I must think that they belong to the superior Extremities; and therefore shall divide each of them into the Shoulder, Arm, Fore-arm, and Hand.

The SHOULDER confifts of the Clavicle

and Scapula.

crooked Bone, in Figure like an *Italick* f placed almost horizontally between the upper lateral Part of the *Stermum*, and what is commonly called the Top of the Shoulder, which, as a Clavis or Beam, it bears off from the Trunk of the Body.

The Clavicle, as well as other long round Bones, is larger at its two Ends, than in the Middle. The End next to the Stermum † is

triangular:

tr

pi

to

pi

vi til

ev of

na

na bl

bo

liq

an

is wh

the

to

Bo

a fi

Sur

ral

Exp

poli

[†] Κωλα, γυα, εκφυαδες, Enata, adnata, explantata membra, artus.

⁽a) Philosoph. Transact. Num. 449. § 5.

* Os jugulare, jugulum, furcula, ligula, clavis, humerus
quibusdam.

[†] Παρασφάγις.

triangular: The Angle behind is confiderably produced, to form a tharp Ridge, to which the transverse Ligament extended from one Clavicle to the other is fixed (a).—The Side opposite to this is fomewhat rounded.—The Middle of this protuberant End, is as irregularly hollowed, as the Cavity in the Stermin for receiving it is raised; but, in a recent Subject, the irregular Concavities of both, are supplied by a moveable Cartilage, which is much more closely connected every where by Ligaments to the Circumference. of the Articulation, than those of the Lower Jaw are; but grows to the two Bones at both its internal and external End; its Substance at the internal End being foft, but very strong, and refembling the intervertebral Cartilages (b).

From this internal End, the Clavicle, for about two Fifths of its Length, is bended obliquely forwards and downwards. On the Upper and Fore-part of this Curvature a finall Ridge is feen, with a plain rough Surface before it; whence the Musculus sterno-byoideus and sterno-mastoideus have in Part their Origin.—Near the lower Angle a small plain Surface is often to be remarked, where the first Rib and this Bone are contiguous (c), and are connected by a firm Ligament (d).—From this a rough plain Surface is extended outwards, where the pectoral Muscle has Part of its Origin.—Behind,

⁽a) Riolan. Encherid Anat. lib. vi. cap. 13.—Winflow, Expos. Anat. des Os frais, § 248.—Weithrecht. Act. Petropolit. tom. 4. p. 255. & Syndesmolog. Sect. 2. I. 3.

⁽b) Weithrecht, Syndesmotog Seet. 2. I. § 6.

⁽c) Dionis, Sixieme Demonst. des Os.

⁽d) Weitbrecht, Syndesmolog. Sect. 2. I. § 7.

11

te

T

C

0

21

pi

ir

the Bone is made flat and rough by the Infer. tion of the larger Share of the Subclavian Muscle. - After the Clavicle begins to be bended backwards, it is round, but foon after becomes broad and thin; which Shape it retains to its external End.—Along the external Concavity, a rough Sinuofity runs, from which some Part of the Deltoid Muscle takes its Rife: --- Opposite to this, on the convex Edge, a scabrous Ridge gives Infertion to a Share of the cucullaris Muscle. The upper Surface of the Clavicle here is flat; but the lower is hollow, for lodging the Beginning of the Musculus subclavius, and towards its Back-part a Tubercle rifes, to which, and a Roughness near it, the strong short thick Ligament connecting the Bone to the coracoid Process of the Scapula is fixed.

The external End * of this Bone is horizontally oblong, smooth, sloping at the posterior Side, and tipped in a recent Subject with a Cartilage, for its Articulation with the Acromion scapulæ.—Round this the Bone is spongy, for the firmer Connexion of the Ligaments.

The medullary Arteries, having their Direction obliquely outwards, enter the Clavicles by one or more small Passages in the Middle of

their Back-part.

The Substance of this Bone is the same as of

the other round long Bones.

The triangular unequal interior End of each Clavicle, has the Cartilage above defcribed, interposed betwixt it and the irrelar Cavity of the Sternum.—The Ligaments,

[&]quot;Exemis.

which furround this Articulation to fecure it, are fo fhort and strong, that little Motion can be allowed any way: and the strong Ligament that is stretched across the upper Furcula of the Stermum, from the posterior prominent Angle of the one Clavicle, to the fame Place of the other Clavicle, ferves to keep each of these Bones more firmly in their Place. By the Affistance, however, of the moveable interveening Cartilage, the Clavicle can, at this Joint, be raised or depressed, and moved backwards and forwards fo much, as that the external End, which is at a great Distance from that Axis, enjoys very confpicuous Motions .--The Articulation of the exterior End of the Clavicle shall be considered after the Description of the Scapula.

The Clavicles of Infants are not deficient in any of their Parts; nor have they any Epiphyses at their Extremities joined afterwards to their Bodies, as most other such long Bones have, which preserves them from being bended too much, and from the Danger of any unoshined Parts being separated by the Force which

pulls the Arms forwards.

The Uses of the Clavicles are, to keep the Scapulæ, and consequently all the superior Extremities, from falling in and forward upon the Thorax; by which, as in most Quadrupeds, the Motions of the Arms would be much confined, and the Breast made too narrow.—The Clavicles likewise afford Origin to several Muscles, and a Defence to large Vessels.

From the Situation, Figure and Use of the Clavicles, it is evident, that they are much ex-

R

bi

P

th

pe

pr

211

M

tri

fu

an

1720

T

an

ces

the

int

bet

app

its

its

posed to Fractures; that their broken Parts must generally go bye each other, and that they are

difficultly kept in their Place afterwards.

SCAPULA, or Shoulder-blade *, is the triangular Bone situated on the Outside of the Ribs, with its longest Side, called its Base, towards the spinal Processes of the Vertebra. and with the Angle at the upper Part of this Side about three Inches, and the lower Angle at a greater Distance from these Procesfes .- The Back-part of the Scapula has nothing but the thin Ends of the Serratus anticus major and Subscapularis Muscles between it and the Ribs: But as this Bone advances forwards, its Distance from the Ribs increases.—The upper or shortest Side, called the superior Costa of the Scapula, is nearly horizontal, and parallel with the fecond Rib .- The lower Side, which is named the inferior Costa, is extended obliquely from the third to the eighth Rib. -- The inferior Angle is very acute; the upper one is near to a right Angle; and what is called the anterior, does not deserve the Name; for the two Sides do not meet to form an Angle.—The Body of this Bone is concave towards the Ribs, and convex behind, where it has the Name of Dorsim +. Three Processes are generally reckoned to proceed from the Scapula. The first is the large Spine that rises from its convex Surface behind, and divides it unequally.— The fecond Process stands out from the Forepart of the upper Side; and, from its imaginary

^{* &#}x27;Ωμο ελάτος, ἐπινώτιον, Latitudo humeri, scoptulum vel scutulum opertum, spatula, ala, humerus, clypeus, scutum thoracis.

⁺ XEAUNION.

Resemblance to a Crow's Beak, is named Coracoides ‡.—The third Process is the whole thick bulbous Fore-part of the Bone.

After thus naming the several constituent. Parts of the Scapula, the particular Description.

will be more eafily understood.

The Base, which is tipped with Cartilage in a young Subject, is not all streight: For above the Spine, it runs obliquely forwards to the superior Angle; that here it might not be too protuberant backwards, and so bruise the Muscles and Teguments: Into the oblique Space the Musculus patientiæ is inserted.—At the Root of the Spine, on the Back-part of the Base, a triangular plain Surface is formed, by the Pressure of the lower Fibres of the Trapezius.—
Below this the Edge of the Scapula is scabrous and rough, for the Insertion of the Serratus major anticus and rhomboid Muscles.

The Back-part of the inferior Angle is made smooth by the latissimus Dorsi passing over it. This Muscle also alters the Direction of the inferior Costa some way forwards from this Angle; and so far it is statted behind by the Origin of the Teres major.—As the inferior Costa advances forward, it is of considerable Thickness, is slightly hollowed, and made smooth behind by the Teres minor, while it has a Fossa formed into it below by Part of the Subscapularis; and between the two a Ridge with a small Depression appears, where the longus extensor Cubiti has

its Origin.

The fuperior Costa is very thin; and near its Fore-part there is a semilunar Nitch, from one

Anchoroides, ligmoides, digitalis, ancistroides.

one End of which to the other a Ligament is stretched; and sometimes the Bone is continued, to form one, or sometimes two Holes for the Passage of the scapular Blood-vessels and Nerves.—Immediately behind this semilunar Cavity the coraco-byoid Muscle has its Rise.—From the Nitch to the Termination of the Fossa for the Teres minor, the Scapula is narrower than any where else, and supports the third Process. This Part has the Name of Cervix.

The whole Dorsum of the Scapula is always faid to be convex; but, by Reason of the raised Edges that surround it, it is divided into two Cavities by the Spine, which is stretched from behind forwards, much nearer to the superior, than to the inferior Costa—The Cavity above the Spine is really concave where the surface of this Bone below the Spine, on which the infra-spinatus Muscle is lodged; while the Surface of this Bone below the Spine, on which the infra-spinatus Muscle is placed, is convex, except a Fossa that runs at the Side of the inferior Costa.

The internal or anterior Surface of this Bone is hollow, except in the Part above the Spine, which is convex.—The fubscapularis Muscle is extended over this Surface, where it forms feveral Ridges and intermediate Depressions, commonly mistaken for Prints of the Ribs; they point out the Interstices of the Bundles of Fibres of which the subscapular is Muscle is composed (a).

The Spine * rifes fmall at the Base of the Scapulu, and becomes higher and broader as it advances forwards.—On the Sides it is

unequally

ull

on

the

ed

fix

Act

and

Spi

of

ed

the

lon

and

the

its

is

Pal

Th

plai

tus

nal

and

has

Ro

fem

whe

Fro

Apo

bec

fumr

⁽a) Winflow, in Memoires de l'Acad. des Sciences, 1722.
• Ραχις, ὑπερύχη ώμεσπλατῶν, Eminentia scapularum.

nuequally hollowed and crooked, by the Actions of the adjacent Muscles.——Its Ridge is divided into two rough flat Surfaces: Into the upper one, the Trapezius Muscle is inserted; and the lower one has Part of the Deltoid fixed to it.—The End of the Spine, called Acronion †, or Top of the Shoulder, is broad and flat, and is sometimes only joined to the Spine by a Cartilage.—The anterior Edge of the Acromion is flat, smooth, and covered with a Cartilage, for its Articulation with the external End of the Clavicle; and it is hollowed below, to allow a Passage to the infra and supra-spinati Muscles, and free Motion to the Os humeri.

The coracoid & Process is crooked, with its Point inclining forwards; fo that a Hollow is left at the lower Side of its Root, for the Passage of the infra-scapularis Muscle. -The End of this Process is marked with three plain Surfaces. Into the internal, the Serratus minor anticus is inserted: From the external, one Head of the Biceps flexor cubiti rifes; and from the lower one, the Coracobrachialis has its Origin. - At the Upper-part of the Root of this Process, immediately before the femilimar Cavity, a smooth Tubercle appears, where a Ligament from the Clavicle is fixed. From all the external Side of this coracoid Apophyse, a broad Ligament goes out, which becomes narrower where it is fixed to the

Acronion

^{*} Pterygium, crista.

t'Επωμις, άγκυροειδής, κορακοειδής, κατακλείς, Acromii os, summus armus, rostrum porcinum, processus digitalis.

S'Ayxupoeisis, oiquoeisis, Rostriformis.

Acronion.—The sharp Pain, violent Instandam, and tedious Cure of Contusions in this Part, are probably owing to these Tendons

and Ligaments being hurt.

From the Cervix scapulæ the third Process is produced. The Fore-part of this is form. ed into a Glenoid Cavity +, which is of the Shape of the longitudinal Section of an Egg, being broad below, and narrow above. Between the Brims of this Hollow and the Fore-part of the Root of the Spine, a large Sinuofity is left, for the Transmission of the fupra and infra-spinati Muscles; and on the Upper-part of these Brims we may remark a fmooth Surface, where the second Head of the Biceps flexor cubiti has its Origin. — The Root of the Supercilia is rough all round, for the firmer Adhesion of the capsular Ligament of the Articulation, and of the Cartilage which is placed on these Brims, where it is thick, but becomes very thin as it is continued towards the Middle of the Cavity, which it lines all over.

The medullary Vessels enter the Scapula

near the Base of the Spine.

The Substance of the Scapula, as in all other broad flat Bones, is cellular, but of an unequal Thickness; for the Neck and third Process are thick and strong. The inferior Costa, Spine, and coracoid Process, are of a middle Thickness; and the Body is so pressed by the Muscles, as to become thin and diaphanous.

The

Su

ne

tio

cai

wh

bu

the

cal

011

ba

the

par

the

W

Su

the

10

thi be

on wit

abl

Toi

int

Bos

to Ar:

end

wh wa:

The Scapula and Clavicle are joined by plain Surfaces, tipped with Cartilage *; by which neither Bone is allowed any confiderable Motion, being tightly tied down by the common capfular Ligament, and by a very strong one which proceeds from the coracoid Process: but divides into two before it is fixed into the Clavicle, with fuch a Direction, as either can allow this Bone to have a small Rotation, in which its posterior Edge turns more backwards, while the anterior one rifes farther forwards; or it can yield to the Forepart of the Scapula moving downwards, while the Back-part of it is drawn upwards; in both which Cases the oblong fmooth articulated Surfaces of the Clavicle and Scapula are not in the same Plane, but stand a little tranversely, or across each other, and thereby preserve this Joint from Luxations, to which it would be subject, if either of the Bones was to move on the other perpendicularly up and down, without any Rotation. --- Sometimes a moveable ligamentous Cartilage is found in this Joint; otherwhiles fuch a Cartilage is only interposed at the anterior Half of it; and in some old Subjects I have found a sesamoid Bone here (a)—The Scapula is connected to the Head, Os byoides, Vertebræ, Ribs, and Arm-bone, by Muscles, that have one End fastened to these Bones, and the other to the Scapula, which can move it upwards, downwards, backwards or forwards; by the quick Succession of thefe

^{*} Acromion, κατακλεις, Clausuræ.
(a) Jac. Sylv. Isagog. Anat. lib. 1. cap. 2.

these Motions, its whole Body is carried in a Circle. But being also often moved, as upon an Axis perpendicular to its Plane, its Circumference turns in a Circle whose Center this Axis is (a). Whichever of these Motions it performs, it always carries the outer End of the Clavicle and the Arm along with it.—
The glenoid Cavity of this Bone receives the Os humeri, which plays in it as a Ball in a Socket, as will be explained more hereafter.

The Use of the Scapula is, to serve as a Fulcrum to the Arm; and, by altering its Position on different Occasions, to allow always the Head of the Os humeri a right situated Socket to move in; and thereby to affist and to enlarge greatly the Motions of the superior Extremity, and to afford the Muscles which rise from it more advantageous Actions, by altering their Directions to the Bone which they are to move.—This Bone also serves to defend the Back-part of the Thorax, and is often employed to sustain Weights, or to resist Forces, too great for the Arm to bear.

7

11

f

11

ti

11

2

2

The Base, Acromion, coracoid Process, and Head of the Scapula, are all in a cartilaginous State at Birth; and the three first are joined as Epiphyses; while the Head, with the glenoid Cavity, is not formed into a distinct separate Bone, but is gradually produced by the offstation of the Body of this Bone being continued

forwards.

The ARM has only one Bone, best known by

⁽a) See Winflow. Memoires de l'Acad. des Sciences, 1716.

the Latin Name of Os humeri +; which is long,

round, and nearly streight.

11

n.

115

It

of

le.

2

1.

75

d

d

h

V

h

IS

d

d

e

.

The upper End of this Bone * is formed into a large round fmooth Head, whose middle Point is not in a streight Line with the Axis of the Bone, but stands obliquely backwards from it. The Extent of the Head is distinguished by a circular Fossa furrounding its Base, where the Head is united to the Bone, and from which the capfular Ligament of the Joint rises. - Below the Fore-part of its Base two Tubercles stand out: The smallest one. which is fituated most to the Inside, has the Tendon of the fubscapularis Muscle inserted into it. The larger more external Protuberance is divided, at its upper Part, into three smooth plain Surfaces: Into the anterior of which, the Musculus supra-spinatus; into the middle or largest, the infra-spinatus; into the one behind, the Teres minor, is inferted .-Between these two Tubercles, exactly in the Fore-part of the Bone, a deep long Fossa is formed, for lodging the tendinous Head of the Biceps flexor cubiti; which, after passing, in a Manner peculiar to itself, through the Cavity of the Articulation, is tied down by a tendinous Sheath extended across the Fossa; in which, and in the neighbouring Tubercles, are feveral remarkable Holes, which are penetrated by the tendinous and ligamentons Fibres, and by Vessels.—On each Side of this Fossa X

^{† &#}x27;Ακρολία, ωλένη, Os brachii, armi, adjutorium, parvum brachium, cana brachii.

^{*} Acrocolium.

2

n

po

CE

in

tu

fe

gi,

as it descends in the Os humeri, a rough Ridge, gently flatted in the Middle, runs from the Roots of the Tubercles .- The Tendon of the pectoral Muscle is fixed into the anterior of these Ridges, and the latiffmus Dorsi, and Teres major, are inferted into the internal one. - A little behind the lower End of this last. another rough Ridge may be observed, where the Coraco-brachialis is inferted.—From the Back-part of the Root of the largest Tubercle. a Ridge also is continued, from which the Brevis extensor cubiti rises.—This Bone is flatted on the Infide, about its Middle, by the Belly of the Biceps flexor cubiti.——In the Middle of this plain Surface, the Entry of the medullary Artery is feen flanting obliquely downwards. - At the Fore-fide of this Plane the Bone rifes in a Sort of Ridge, which is rough, and often has a great many small Holes in it; where the Tendon of the strong Deltoid Muscle is inferted; on each Side of which the Bone is fmooth and flat, where the Brachiaus internus The exterior of these two flat Surfaces is the largest; behind it a superficial spiral Channel, formed by the Muscular Nerve and the Vessels that acompany it, runs from behind forwards and downwards. The Body of the Os humeri is flatted behind by the Extensors of the Fore-arm. -- Near the lower End of this Bone, a large sharp Ridge is extended on its Out-fide, from which the Musculus spinator radii longus and the longest Head of the Extensor carpi radialis rise.— Opposite to this, there is another small Ridge to which the aponeurotic Tendon, that gives Origin

Origin to the Fibres of the internal and external Brachizei Muscles is fixed; and from a little Depression on the Fore-side of it, the Pronator radii teres rises.

The Body of the Os humeri becomes gradually broader towards the lower End, where it has several Processes: at the Roots of which. there is a Cavity before, and another behind *. The anterior is divided by a Ridge into two; the external, which is the least, receives the End of the Radius: and the internal receives the Coronoid Process of the Ulna in the Flexions of the Fore-arm, while the posterior deep triangular Cavity lodges the Olecranon in the Extensions of that Member. The Bone betwixt these two Cavities is pressed so thin by the Processes of the Ulna, as to appear diaphanous in feveral Subjects. The Sides of the posterior Cavity are stretched out into two Processes, one on each Side: These are called Condyles; from each of which a strong Ligament goes out to the Bones of the Fore-arm. -The external Condyle, which has an oblique Direction also forwards in respect of the internal, when the Arm is in the most natural Posture (a) is equally broad, and has an obtuse smooth Head rising from it forwards.— From the rough Part of the Condyle, the inferior Head of the Bicornis, the Extensor digitorum communis, Extensor carpi ulnaris, Anconœus, and some Part of the Supinator radii brevis take their Rife; and on the smooth X 2 Head

^{*} Βαθμίδεις.
(2) Winstow, Mem ires de l'Acad. des Sciences, 1722.

21

th

N

fc

th

th

to

ft

tic

Su

pe

ru

the

ve

18

Ar

Bo

be

18

of

Head the upper End of the Radius plays. Immediately on the Outfide of this, there is a Sinuolity made by the shorter Head of the Bicornis Muscle, upon which the Muscular Nerve is placed.—The internal Condyle is more pointed and protuberant than the external, to give Origin to some part of the Flexor carpi radialis, Pronator radii teres, Palmaris longus, Flexor digitorum sublimis, and Flexor car. pi ulnaris .- Between the two Condyles, is the Troch. lea, or Pully, which confifts of two lateral Protuberances, and a middle Cavity, that are smooth and covered with Cartilage. - When the Fore. arm is extended, the Tendon of the internal Brachiæus Muscle is lodged in the Fore-part of the Cavity of this Pully. - The external Protuberance, which is less than the other, has a sharp Edge behind; but forwards, this Ridge is obtuse, and only feparated from the little Head, already described, by a small Fossa, in which the joined Edges of the Ulna and Radius move. The internal Protuberance of the Pully is largest and highest; and therefore in the Motions of the Ulna upon it, that Bone would be inclined outwards, was it not supported by the Radius on Side. — Between this Protuberance and Condyle, a Sinuofity may be remarked, where the ulnar Nerve passes.

The Substance and the internal Structure of the Os humeri is the same, and disposed in the

fame Way, as in other long Bones.

The round Head at the upper End of this Bone is articulated with the Glenoid Cavity of the Scapula; which being superficial, and having long Ligaments, allows the Arm a free and

and extensive Motion. — These Ligaments are however confiderably strong: For, besides the common capfular one, the Tendons of the. Muscles perform the Office, and have been described under the Name of Ligaments. Then the Acronion and Coracoid Process, with the strong broad Ligaments Gretched betwixt them, fecure the Articulation above, where the greatest and most frequent Force is applied to thrust the Head of the Bone out of its Place. It is true, that there is not near for ftrong a Defence at the lower Part of the Articulation; but in the ordinary Postures of the Arm, that is, so long as it is at an acute Angle with the Trunk of the Body, there cannot he any Force applied at this Place to occasion. a Luxation, fince the Joint is protected fo well

The Motions which the Arm enjoys by this Articulation, are to every Side; and by the Succession of these different Motions, a Circle may be described. Besides which, the Bone performs a small Rotation round its own Axis. But though this can be performed with the round Head in all Positions; yet as these vary, the Effects upon the Body of the Bone are very different: For, if the Middle of the Head. is the Center of Rotation, as it is when the Arm hangs down by the Side, the Body of the Bone is only moved forwards and backwards; because the Axis of Motion of the Head is nearly at right Angles with the Length of the Bone (a); whereas when the Arm is X 3 raifed

⁽a) Hippocrat. de Articul. § r.

raised to right Angles with the Trunk of the Body, the Center of Motion, and the Axis of the Bone, come to be in the same streight Line; and therefore the Body of the Os humeri performs the same Motion with its Head.—
Though the Motions of the Arm seem to be very extensive, yet the larger Share of them depends on the Motion of the Scapula.—
The lower End of the Os humeri is articulated with the Bones of the Fore-arm, and carries them with it in all its Motions, but serves as a Base on which they perform the Motions peculiar to themselves; as shall be described afterwards.

172

by

Fi

25

of

of

Pro

and

fca

chi

ma

ext

the

or app

cels

as t

and

rund

Trail

Both the Ends of this Bone are cartilaginous in a new-born Infant, and the large Head with the two Tubercles, and the Trochlea with the two Condyles, become Epiphyses before they

are united to the Body of the Bone.

The FORE-ARM * confifts of two long Bones, the Ulna and Radius; whose Situation, in respect of each other, is oblique in the least straining or most natural Posture; that is, the Ulna is not directly behind, nor on the Outside of the Radius, but in a middle Situation between these two, and the Radius crosses it.—The Situation however of these two Bones, and of all the other Bones of the superior Extremity that are not yet described, is frequently altered; and therefore, to thun Repetitions, I desire it may be now remarked, that, in the remaining Account of the superior Extremity, I understand by

[&]quot;Cubitus, angue, alevn, muyav, Ulea, lacertus.

by the Term of posterior, that Part which is in the same Direction with the Back of the Hand; by anterior, that answering to the Palm; by internal, that on the same Side with the Thumb; by external, the Side nearest to the little Finger; supposing the Hand always to be in a middle Position between Pronation and Supination.

as a Measure, is the longest of the two Bones of the Fore-arm, and situated on the Outside

of the Radius.

At the upper End of the Ulna are two Processes. The posterior is the largest, and formed like a Hook, whose concave Surface moves upon the Pully of the Os humeri, and is called Olecranon ||, or Top of the Cubit. -The convex Back-part of it is rough and scabrous, where the Longus Brevis, and Brachiaus externus, are inferted. The Olecranon makes it unnecessary that the Tendons of the extensor Muscles should pass over the End of the Os humeri; which would have been of ill Consequence in the great Flexions of this Joint, or when any confiderable external Force is applied to this Part (a). The anterior Process is not so large, nor does it reach so high, as the one behind; but it is sharper at its End, and therefore is named Coronoid.—Between thefe

[†] Cubitus, πηχυς, προπήχιον, Focile majus, Canna vel arundo major & inferior Brachii.

Ayrar, Gibber, Cubitus, Additamentum necatum.

⁽a) Winflow. Exposition Anatomique du Corps humain, Traite des os secs, § 979.

these two Processes, a large semicircular or figmoid Concavity is left; the Surface of which, on each Side of a middle Rifing, is flanting, and exactly adapted to the Pully of the Bone of the Arm .- Across the middle of it, there is a small Sinuosity for lodging mucilaginous Glands; where, as well as in a small Hollow on the internal Side of it, the Cartilage that lines the rest of its Surface is wanting.-Round the Brims of this Concavity the Bone is rough, where the capfular Ligament of the Joint is implanted.—Immediately below the Olecranon on the Back-part of the Ulna, a flat triangular spongy Surface appears, on which we commonly lean.—At the internal Side of this, there is a larger hollow Surface, where the Musculus Anconæus is lodged; and the Ridge at the Infide of this gives Rife to the Museulus supinator radii brevis. — Between the Top of the Ridge and the coronoid Process is the femilunated smooth Cavity, lined with Cartilage, in which, and a Ligament extended from the one to the other End of this Cavity, the round Head of the Radius plays. —— Immediately below it a rough Hollow gives Lodging to mucaginous Glands. --Below the Root of the voronoid Process, this Bone is feabrous and unequal, where the Brachiæus intermus is inferted. - On the Outfide of that we observe a smooth Concavity, where the Beginning of the Flexor digitorum profundus sprouts out.

The Body of the *Ulna* is triangular.—
The internal Angle is very sharp, where the Ligament that connects the two Bones is fix-

d;

ari

fio

he

Le

pa

ter

co

bli

dif

ſm

To

laft

to

13 1

the

styl

15

to 1

fme

nai

Pro

ma

Ne

and

and

mo

ed;—The Sides, which make this Angle, are flat and rough, by the Action and Adhefion of the many Muscles which are fituated here.—At the Distance of one Third of the Length of the Ulna from the Top, in its Forepart, the Passage of the medullary Vessels is to be remarked slanting upwards.—The external Side of this Bone is smooth, somewhat convex, and the Angles at each Edge of it are blunted by the Pressure of the Muscles equally

disposed about them.

As this Bone descends, it becomes gradually smaller; so that its lower End terminates in a little Head, standing on a small Neck .-Towards the fore but outer Part of which last, an oblique Ridge runs, that gives Rife to the Pronator radii quadratus.— The Head is round, fmooth, and covered with a Cartilage on its internal Side, to be received into the femilianar Cavity of the Radius; while a hyloid Process + rises from its Outside, to which is fixed a strong Ligament that is extended to the Os cuneiforme and pisiforme of the Wrist. -Between the Back-part of that internal smooth Side and this Process, a Sinuosity is left for the Tendon of the Extensor carpiulnaris.—On the Fore-part of the Root of the Proceis, fuch another Depression may be remarked for the Passage of the ulnar Artery and Nerve. The End of the Bone is smooth, and covered with a Cartilage. Between it and the Bones of the Wrist, a doubly-concave moveable Cartilage is interposed; which is a Continu-

[†] Tfapousn, Malleolus externus.

nat

18 .1

the

the

cel

cep

Ric

Su

low

fca

res

cor

wh

fur

of

fac

the

bot

wh

the

litt

011

laft

Pai

Wal

7101

wh

the

its !

In a

Continuation of the Cartilage that covers the lower End of the Radius, and is connected loosely to the Root of the styloid Process, and to the rough Cavity there; in which mu-

caginous Glands are lodged.

The Ulna is articulated above with the lower End of the Os humeri, where these Bones have Depressions and Protuberances correspond. ing to each other, so as to allow an easy and secure Extension of the Fore-arm to almost a streight Line with the Arm, and Flexion to a very acute Angle; but, by the flanting Position of the Pully, the lower Part of the Fore-arm is turned outwards in the Extension, and inwards in the Flexion (a); and a very finall kind of Rotation is likewife allowed in all Positions, especially when the Ligaments are most relaxed by the Fore-arm being in a middle Degree of Flexion.—The Ulna is also articulated with the Radius and Corpus, in a manner to be related afterwards.

RADIUS +, fo called from its imagined Resemblance to a Spoke of a Wheel, or to a Weaver's Beam, is the Bone placed at the Instide of the Fore-arm. Its upper End is formed into a circular little Head, which is hollowed for an Articulation with the Tubercle at the Side of the Pulley of the Os humeri; and the Half of the round Circumference of the Head next to the Uhia is smooth, and covered with a Cartilage, in order to be received into the semilar

(a) Winflow, Memoires de l'Acad. des Sciences, 1722.

† Κερχὶς, παρὰπηχιον, Focile minus, Canna minor, Arundo minor.

nated Cavity of that Bone.—Below the Head, the Radius is much smaller; therefore this Part is named its Cervix, which is made round by the Action of the Supinator radii brevis.—At the external Root of this Neck, a tuberous Process rises; into the outer Part of which the Biceps slexor cubiti is inserted.—From this a Ridge runs downwards and inwards, where the Supinator radii brevis is inserted; and a little below, and behind this Ridge, there is a rough scabrous Surface, where the Pronator radii teres is fixed.

The Body of the Radius is not streight, but convex on its internal and posterior Surfaces; where it is also made round by the equal Presfure of the circumjacent Muscles, particularly of the Extensors of the Thumb; but the Surfaces next to the Ulna are flatted and rough, for the Origin of the Muscles of the Hand; and both terminate in a common sharp Spine, to which the strong Ligament extended betwixt the two Bones of the Fore-arm is fixed.—A little below the Beginning of the plain Surface, on its Fore-part, where the Flexor Muscle of the last Joint of the Thumb takes its Origin, the Pallage of the medullary Vessels is feen slanting upwards.—The Radius becomes broader and flatter towards the lower End, especially on its Fore-part, where its Pronator quadratus Muscle is situated.

The lower End of the Radius is larger than the superior; though not in such a Disproportion, as the upper End of the Ulna is larger than its lower End.——Its Back-part has a flat strong Ridge in the Middle, and Fossa on each Side.— In a small Groove, immediately on the Outside

of the Ridge, the Tendon of the Extensor ter. tii internodii pollicis plays .-- In a large one be. yond this, the Tendons of the Indicator and of the common Extensor Muscles of the Fingers pass—Contiguous to the Uhia, there is a small Depression made by the Extensor minimi digiti. On the Infide of the Ridge there is a broad Depression, which seems again subdivided, where the two Tendons of the Bicornis, or Extensor of this End of the Radius, is also hollowed by the Extensors of the first and second Joint of the Thumb; immediately above which, a little rough Surface shews where the Supinator radii longus is inserted. The Ridges at the Sides of the Grooves, in which the Tendons play, have an annular Ligament fixed to them, by which the several Sheaths for the Tendons are formed.—The Fore-part of this End of the Radius is also depressed, where the Flexors of the Fingers and Flexor carpi radialis pass.-The external Side is formed into a femilunated smooth Cavity, lined with a Cartilage, for receiving the lower End of the Ulna.—The lowest Part of the Radius is formed into an oblong Cavity; in the Middle of which is a small transverse Rising, gently hollowed, for lodging mucilaginous Glands; while the Rifing itself is infinuated into the Conjunction of the two Bones of the Wrist that are received here.-The internal Side of this Articulation is fenced by a remarkable Process * of the Radius, from which a Ligament goes out to the Wrist as the /ty loid

[·] Malleolus internus, Processus styloides.

flyloid Process of the Ulna with its Ligament

guards it on the Outside.

The Ends of both the Bones of the Forearm being thicker than the Middle, there is a considerable Distance between the Bodies of these Bones; in the larger Part of which a strong tendinous, but thin Ligament, is extended, to give a large enough Surface for the Origin of the numerous Fibres of the Muscles situated here, that are so much sunk between the Bones, as to be protected from Injuries, which they would otherwise be exposed to. But this Ligament is wanting near the upper End of the Fore-arm, where the Supinator radii brevis, and Flexor digitorum profundus, are immediately connected (a).

Both Ends of the Bones of the Fore-arm, are first Cartilages, and then Epiphyses in Chil-

dren.

of

rs

d

or

y

lii

٧,

As the head of the Radius receives the Tubercle of the Os humeri, it is not only bended and extended along with the Ulna, but may be moved round its Axis in any Position; and that this Motion round its Axis may be sufficiently large, the Ligament of the Articulation is extended farther down than ordinary on the Neck of this Bone, before it is connected to it; and it is very thin at its upper and lower Part, but makes a firm Ring in the Middle.—This Bone is also joined to the Ulna by a double Articulation; for above a Tubercle of the Radius plays in a Socket of the Ulna, whilst below the Radius gives the Socket and the Ulna the Tubercle:

⁽a) Weithrecht. Syndesmolog. Fig. 10. 11.

But then the Motion performed in these two is very different; for at the upper End, the Radius does no more than turn round its Axis: while at the lower End, it moves in a fort of Ci. cloid, upon the round Part of the Ulna; and as the Hand is articulated and firmly connected here with the Radius, they must move together— When the Palm is turned uppermost, the Radius is faid to perform the Supination; when the Back of the Hand is above, it is faid to be Prone. But then the Quickness and large Extent of these two Motions are assisted by the Ulna, which, as was before observed, can move with a kind of small Rotation on the sloping Sides of the Pulley. This lateral Motion, tho very inconfiderable in the Joint itself, is conspicuous at the lower End of fuch a long Bone; and the strong Ligament connecting this lower End to the Carpus, makes the Hand more readily to obey these Motions. - When we defign a large circular Turn of our Hand, we increase it by the Rotation of the Os humeri, and fometimes employ the Spine and inferior Extremities, to make these Motions of Pronation and Supination of the Hand large enough.

The HAND * comprehends all from the Joint of the Wrist to the Points of the Fingers. Its Back-part is convex, for greater Firmness and Strength; and it is concave before, for containing more surely and conveniently such Bodies as we take hold of:—One Half of the Hand has an obscure Motion in Comparison of what the other has, and serves as a Base to the moveable

^{* &#}x27;Axpoxesp, summa Manus.

moveable Half; which can be extended back very little farther than to a streight Line with the Fore-arm, but can be confiderably bended forwards.

As the Bones that compose the Hand are of different Shapes and Uses, while several of them that are contiguous agree in some general Characters; the Hand is, on this Account, conmonly divided into the Carpus, Metacarpus, and Fingers; among which last the Thumb is reckoned.

The CARPUS * is composed of eight small foongy Bones, fituated at the upper Part of the Hand. I shall describe each of these Bones, under a proper Name taken from their Figure (a), because the Method of ranging them by Numbers, leaves Anatomists too much at liberty to debate, very idly, which ought to be preferred to the first Number; or, which is worse, feveral, without explaining the Order they obferve, differently apply the same Numbers, and fo confound their Readers.—But that the Description of these Bones may be in the same Order as they are found in the Generality of Anatomical Books, I shall begin with the Range of Bones that are concerned in the moveable Joint of the Wrist, or are connected to the Fore-arm, and shall afterwards consider the four that support the Thumb and Offa Metacarpi of the Fingers.

The eight Bones of the Carpus are, Os scaphoides, lunare, cuneiforme, pisiforme, trapezi-

um, trapezoides, magnum, unciforme.

The

[†] Kreis, Brachiale, prima palmæ pars, Rasetta. (a) Lyfer. Cult. Anat. lib. 5. cap. 2.

The Scaphoides is situated most internally of those that are articulated with the Fore-arm.— The Lunare is immediately on the Outside of the former.— The Cuneiforme is placed still more externally, but does not reach so high up as the other two.— The Pissorme stands forwards into the Palm from the Cuneiforme.— The Trapezium is the first of the second Row, and is situated betwixt the Scaphoides and first Joint of the Thumb.— The Trapezium.—

The Os magnum is still more external;— The Unciforme is farthest to the Side of the Little.

V

t

W

11

in

tic

Sic

Its

Sui

pili

finger.

Os Scaphoides * is the largest of the eight ex. cept one. It is convex above, concave and oblong below; from which small Resemblance of a Boat it has got its Name.—Its smooth convex Surface is divided by a rough middle Fossa, which runs obliquely crofs it. The upper largest Division is articulated with the Radius .-Into the Fossa the common Ligament of the Joint of the Wrist is fixed; and the lower Division is joined to the Trapezium and Trapezoides.—The Concavity receives more than an Half of the round Head of the Os magram. --- The external Side of this Hollow is form. ed into a semilunar Plane, to be articulated with the following Bone. The internal, poiterior and anterior Edges, are rough, for fixing the Ligaments that connect it to the furrounding Bones. 05

^{*} Koruhomles, Naviculare.

Os lunare + has a finooth convex upper Surface, by which it is articulated with the Radius.—The internal Side, which gives the Name to the Bone, is in the Form of a Crefcent, and is joined with the Scaphoid; -- the lower Surface is hollow, for receiving part of the Head of the Os magnum. On the Outfide of this Cavity is another smooth, but narrow oblong Sinuofity, for receiving the upper End of the Os unciforme: on the Outside of which a fmall round Convexity is found, for its Connexion with the Os cuneiforme. Between the great Convexity above, and the first deep inferior Cavity, there is a rough Fossa, in which the circular Ligament of the Joint of the Wrift is fixed.

Os cuneiforme * is broader above, and towards the Back of the Hand, than it is below and forwards; which gives it the Resemblance of a Wedge.—The fuperior flightly convex Surface is included in the Joint of the Wrist, being opposed to the lower End of the Ulna .--Below this the cuneiform Bone has a rough Fosfa, wherein the Ligament of the Articulation of the Wrist is fixed. On the internal Side of this Bone, where it is contiguous to the Os lunare, it is smooth and slightly concave .--Its lower Surface, where it is contiguous to the Os unciforme is oblong, somewhat spiral, and concave. Near the Middle of its anterior Surface a circular Plane appears, where the Os piliforme is fustained.

Y 3

Os

[†] Lunatum.

[.] Triquetrum.

01

de

W

is

in

m

la

bo

it

ir

al

fc

T

0

fo

Os pisiforme + is almost spherical, except one circular Plane, or flightly hollow Surface, which is covered with Cartilage for its Motion on the cuneiform Bone, from which its whole rough Body is prominent forwards into the Palm; having the Tendon of the Flexor carpi ulnaris, and a Ligament from the styloid Process of the Ulna, fixed to its upper Part; the transverse Ligament of the Wrist is connected to its internal Side; Ligaments extended to the unciform Bone, and to the Os metacarpi of the Little-finger, are attached to its lower Part; and the Abductor minimi digiti has its Origin from its Fore-part; and, at the internal Side of it, a small Depression is formed, for the Passage of the ulnar Nerve.

Trapezium * has four unequal Sides and Angles in its Back-part, from which it has got its Name. - Above, its Surface is fmooth, flightly hollowed, and femicircular, for its Conjunction with the Os scaphoides.—Its external Side is an oblong concave Square, for receiving the following Bone. The inferior Surface is formed into a Pulley; the two protuberant Sides of which are external and internal. On this Pulley the first Bone of the Thumb is moved.—At the external Side of the external Protuberance, a finall obling fmooth Surface is formed by the Os metacarpi indicis. --- The Fore-part of the Trapezium is prominent in the Palms, and, near to the external Side, has a Sinuofity in it, where the Tendon

† Cartilaginosum, subrotundum, rectum.

^{*} Os cabiforme, trapezoides, multanguium majus.

Tendon of the Flexor carpi radialis is lodged; on the ligamentous Sheath of which the Tendon of the Flexor tertii internodii pollicis plays: And still more externally the Bone is scabrous, where the transverse Ligament of the Wrist is connected, the Abductor and Flexor primi internodii pollicis have their Origin, and Ligaments go out to the first Bone of the Thumb.

Os trapezoides *, so called from the irregular quadrangular Figure of its Back-part, is the smallest Bone of the Wrist, except the Pisisorme.—The Figure of it is an irregular Cube.—It has a small hollow Surface above, by which it joins the Scaphoides; a long convex one internally, where it is contiguous to the Trapezium; a small external one, for its Conjunction with the Os magnum; and an inferior convex Surface, the Edges of which are however so raised before and behind, that a fort of Pulley is formed, where it sustains the Os metacarpi indicis.

Os magnum +, so called because it is the largest Bone of the Carpus, is oblong, having sour quadrangular Sides, with a round upper End, and a triangular plain one below.—
The round Head is divided by a small Rising, opposite to the Connection of the Os scaphoides and lunare, which together form the Cavity for receiving it.—On the Inside a short plain Surface joins the Os magnum to the Trapezoides.—On the Outside is a long narrow concave Surface, where it is contiguous to the Os unciforme.

^{*} Trapezium, multangulum minus» † Maximum. capitatum.

forme.—The lower End, which sustains the metacarpal Bone of the Middle-singer, is triangular, slightly hollowed, and farther advanced on the internal Side than on the external, having a considerable oblong Depression made on the advanced Inside by the metacarpal Bone of the Fore-singer; and generally there is a small Mark of the Os metacar-

pi digiti annularis on its external Side.

Os unciforme + has got its Name from a thin broad Process that stands out from it forwards into the Palm, and is hollow on its Infide, for affording Passage to the Tendons of the Flexors of the Fingers. To this Process also the transverse Ligament is fixed, that binds down and defends these Tendons; and the Flexor and Abdictor Muscles of the Littlefinger have part of their Origin from it .-The upper plain Surface is small, convex, and joined with the Os hmare: — The internal Side is long, and flightly convex, adapted to the contiguous Os magnum: --- The external Surface is oblique, and irregularly convex, to be articulated with the cuneiform Bone: -- The lower End is divided into two concave Surfaces; the external is joined with the metacarpal Bone of the Little-finger, and the internal one is fitted to the metacarpal Bone of the Ring-finger.

In the Description of the preceeding eight Bones, I have only mentioned those plain Surfaces covered with Cartilage, by which they are articulated to each other, or to some o-

ther

fc

21

S

ir

W

At

to

21

b

CI

m

[†] Cuneiforme.

ther Bones, except in some few Cases, where fomething extraordinary was to be observed: and I have defignedly omitted the other rough Surfaces, lest, by crowding too many Words in the Description of such small Bones, the whole should be unintelligible: But these scabrous Parts of the Bones may eafily be underflood, after mentioning their Figure, if it is observed, that they are generally found only towards the Back or Palm of the Hand; that they are all plain, larger behind than before; and that they receive the different Ligaments, by which they are either connected to neighbouring Bones, or to one another: for these Ligaments cover all the Bones, and are fo accurately applied to them, that, at first View, the whole Carpus of a recent Subject appears one smooth Bone (a).

As the Surfaces of these Bones are largest behind, the Figure of the whole conjoined must be convex there, and concave before; which Concavity is still more increased by the Os pisiforme, and Process of the Os unciforme, standing forwards on one Side, as the Trapezium does on the other: And the Bones are securely kept in this Form, by the broad strong transverse Ligament connected to these Parts of them that stand prominent into the Palm of the Hand.—The Convexity behind renders the whole Fabric stronger, where it is most exposed to Injuries; and the large anterior

⁽a) Galen, de usu Part. lib. 2. cap. 8. For a particular Defeription of these Ligaments, see Weithrecht. Syndesmolog. P. 35. -68.

terior Hollow is necessary for a safe Passage to the numerous Vessels, Nerves, and Tendons of the Fingers.

The Substance of these Bones is spongy and cellular, but strong in respect of their

a

ti

11

I

p

to

lu

of

T

10

Bo

Bulk.

The three first Bones of the Carpus make an oblong Head, by which they are articulated with the Cavity at the lower Ends of the Bones of the Fore-arm; so as to allow Motion to all Sides, and, by a quick Succession of these Motions, they may be moved in a Circle. But as the Joint is oblong, and therefore the two Dimensions are unequal, no Motion is allowed to the Carpus round its Axis, except what it has in the Pronation and Supination along with the Radius. -- The Articulation of the first three Bones of the superior Row, with the Bones of the inferior, is such as allows of Motion, especially backwards and forwards; to the Security and Easiness of which the Reception of the Os magnum into the Cavity formed by the Scaphoides and Latnare contributes confiderably: And the greatest Number of the Muscles that serve for the Motion of the Wrist on the Radius, being inserted beyond the Conjunction of the first Row of Bones with the fecond, act equally on this Articulation as they do on the former; but the Joint formed with the Radius being the most easily moved, the first Effect of these Muscles is on it; and the second Row of the Carpus is only moved afterwards. means a larger Motion of the Wrist is allowed

ed than otherwise it could have had safely: For, if as large Motion had been given to one Joint, the Angle of Flexion would have been very acute, and the Ligaments must have been longer than was consistent with the Firmness and Security of the Joint.—The other Articulations of the Bones here being by nearly plain Surfaces, scarce allow of any more Motion, because of the strong connecting Ligaments, than to yield a little, and so elude the Force of any external Power; and to render the Back of the Wrist a little more flat, or the Palm more hollow, on proper Occasions. The Articulations of the Thumb and metacarpal Bones shall be examined afterwards.

The Uses of the Carpus are, to serve as a Base to the Hand, to protect its Tendons, and

to afford it a free large Motion.

All the Bones of the Carpus are in a carti-

laginous State at the Time of Birth.

On Account of the many Tendons that pass upon the lower End of the Fore-arm and the Carpus, and of the numerous Ligaments of these Tendons and of the Bones, which have lubricating Liquors supplied to them, the Pain of Sprains here is acute, the Parts take long Time to recover their Tone, and their Swellings are very obstinate.

METACARPUS + confifts of four Bones which fustain the Fingers.—Each Bone is long and round, with its Ends larger than its Body.—The Upper End, which some call the

Base,

[†] Κτώς, προκάρπιον, επθος, ανδηρον, κτενίον, postbrachiale, pectus, palma, pecten.

Base, is stat and oblong, without any considerable Head or Cavity; but it is however some. What hollowed, for the Articulation with the Carpus: It is made stat and smooth on the Sides where these Bones are contiguous to each other.—Their Bodies are statted on their Backpart by the Tendons of the Extensors of the Fingers.—The anterior Surface of these Bodies is a little concave, especially in their Middle; along which a sharp Ridge stands out, which separates the musiculi interosser placed on each Side of these Bones, which are there made stat and plain by these Muscles.

Their lower Ends are raised into large oblong smooth Heads, whose greatest Extent is forwards from the Axis of the Bone.—At the Forepart of each Side of the Root of each of these Heads, one or two Tubercles stand out, for fixing the Ligaments that go from one metacarpal Bone to another, to preserve them from being drawn asunder:—Round the Heads a rough Ring may be remarked, for the capsular Ligaments of the first Joints of the Fingers to be fixed to; and both Sides of these Heads

are flat, by preffing on each other.

The Substance of the metacarpal Bones is

the fame with that of all long Bones.

At the Time of Birth, these Bones are cartilaginous at both Ends, which afterwards be-

come Epiphyses.

The metacarpal Bones are joined above to the Ossa Carpi and to each other by nearly plain Surfaces. These Connexions are not fit for large Motions—The Articulation of their round Heads at the lower Ends with the Cavities

Cavities of the first Bones of the Fingers, is to be taken Notice of hereaster.

The Concavity on the Fore-part of these metacarpal Bones, and the placing their Bases on the arched Carpus, cause them to form a Hollow in the Palm of the Hand, which is useful often to us.—The Spaces between them lodge Muscles, and their small Motion makes them sit Supporters for the Fingers to play on.

Though the Ossa metacarpi so far agree, yet they may be distinguished from each other

by the following Marks.

1

a

ot

of

10

The Os metacarpi indicis is generally the longest. -- Its Base, which is articulated with the Os trapezoides, is hollow in the Middle. -The small Ridge on the internal Side of this oblong Cavity is smaller than the one opposite to it, and is made flat on the Side by the Trapezium. — The exterior Ridge is also smooth, and flat on its Outside, for its Conjunction with the Os magmim; immediately below which a femicircular smooth flat Surface shews the Articulation of this to the fecond metacarpal Bone. - The Back-part of this Base is flatted, where the long Head of the Extensor carpi radialis is inserted; and its Fore-part is prominent, where the Tendon of the Flexor carpi radialis is fixed.—The external Side of the Body of this Bone is more hollowed by the Action of Muscles, than the internal—The Tubercle at the internal Root of its Head is larger than the external.— Its Base is so firmly fixed to the Bones it is connected with, that it has no Motion.

L

Os metacarpi medii digiti is generally the fecond in Length; but often it is as long as the former; fometimes it is longer; and frequently it only appears to equal the first by the Os magmim being farther advanced down. wards than any other Bone of the Wrift .-Its Base is a broad superficial Cavity, slanting outwards: the internal posterior Angle of which is so prominent, as to have the Appearance of a Process. - The internal Side of this Base is made plain in the same Way as the external Side of the former Bone, while its external Side has two hollow circular Surfaces, for joining the third metacarpal Bone; and between these Surfaces there is a rough Fossa, for the Adhesion of a Ligament, and lodging mucilaginous Glands.—The shorter Head of the Bicornis is inferted into the Back-part of this Bafe. - The two Sides of this Bone are almost equally flatted; only the Ridge on the Fore-part of the Body inclines outwards. --- The Tubercles at the Fore-part of the Root of the Head are equal.—The Motion of this Bone is very little more than the first metacarpal one has; and therefore these two firmly refift Bodies pressed against them by the Thumb, or Fingers, or both.

Os metacarpi digiti annularis is shorter than the second metacarpal Bone.—Its Base is semicircular and convex, for its Conjunction with the Os unciforme.—On its internal Side are two smooth Convexities, and a middle Fossa, adapted to the second metacarpal Bone.—The external Side has a triangular smooth concave Surface, to join it with the fourth

1e

e.

y

n-

g

of

fe

r-

r-

r

i,

gd

rt

e

n

3.

le

n

0

y

n

1

ľ

fourth one. The anterior Ridge of its Body is fituated more to the Out than to the In-fide.

— The Tubercles near the Head are equal.

— The Motion of this third metacarpal Bone is greater than the Motion of the fecond.

Os metacarpi minimi digiti is the smallest and sharpest .- Its Base is irregularly convex, and rifes flanting outwards. -- Its internal Side is exactly adapted to the third metacarpal Bone. The external has no fmooth Surface, because it is not contiguous to any other Bone; but it is prominent where the Extenfor carpi ulnaris is inferted. —As this Os metacarpi is furnished with a proper moving Muscle, has the plainest Articulation, is most loosely connected and least confined, it not only enjoys a much larger Motion than any of the rest, but draws the third Bone with it, when the Palm of the Hand is to be made hollow by its Advancement forwards, and by the Prominence of the Thumb opposite to it.

The THUMB and four FINGERS are

each composed of three long Bones.

The Thumb * is situated obliquely in respect of the Fingers, neither opposite directly to them, nor in the same Plane with them.—All its Bones are much thicker and stronger in Proportion to their Length, than the Bones of the Fingers are: Which was extremely necessary, since the Thumb counteracts all the Fingers.

The first Bone of the Thumb has its Base adapted to the double Pulley of the Trapezi-

Z 2 uni:

^{* &#}x27;Αντίχειρ, δικονδυλός, Magnus digitus, promanus.

um: For, in viewing it from one Side to the other, it appears convex in the Middle; but when considered from behind forwards, it is concave there.—The Edge at the Fore-part of this Base is produced farther than any other Part; and round the Back-part of the Base a rough Fossa may be seen, for the Connection of the Ligaments of this Joint.—The Body and Head of this Bone are of the same Shape as the Ossa metacarpi; only that the Body is shorter, and the Head slatter, with the Tubercles at the Fore-part of its Root larger.

The Articulation of the upper End of this Bone is uncommon: For though it has Protuberances and Depressions adapted to the double Pulley of the *Trapezium*, yet it enjoys a circular Motion, as the Joints do where a round Head of one Bone plays in the orbicular Socket of another; only it is somewhat more confined and less expeditious, but stronger and more

fecure, than fuch Joints generally are.

This Bone of Children is in the same State

with the metacarpal Bones.

The fecond Bone of the Thumb has a large Base formed into an oblong Cavity, whose greatest Length is from one Side to the other.

Round it several Tubercles may be remarked, for the Insertion of Ligaments.—
Its Body is convex, or a Half-round behind; but flat before, for lodging the Tendon of the long Flexor of the Thumb, which is tied down by ligamentous Sheaths that are fixed on each Side to the Angle at the Edge of this flat Surface.—The lower End of this second Bone has

has two lateral round Protuberances, and a middle Cavity, whose greatest Extent of smooth Surface is forwards.

The Articulation and Motion of the upper End of this fecond Bone is as fingular as that of the former.—For its Cavity being joined to the round Head of the first Bone, it would feem at first View to enjoy Motion in all Directions; yet because of the Strength of its lateral Ligaments, oblong Figure of the Joint itself, and Mobility of the first Joint, it only allows Flexion and Extension; and these are

generally much confined.

The third Bone of the Thumb is the smallest, with a large Base, whose greatest Extent is from one Side to the other.—This Base is formed into two Cavities and a middle Protuberance, to be adapted to the Pulley of the some Bone.—Its Body is rounded behind; but is flatter than in the former Bone, for suftaining the Nails.—It is flat and rough before, by the Insertion of the Flexor tertii internodii.—This Bone becomes gradually smaller, till near the lower End, where it is a little enlarged, and has an oval scabrous Edge.

The Motion of this third Bone is confined

to Flexion and Extension.

The orderly Disposition of the Bones of the Fingers into three Rows, has made them generally obtain the Name of three phalanges §.

All of them have half round convex Surfaces, covered with an Aponeurosis, formed by Z 3 the

[§] Scytalidæ, internodia, scuticula, agmina, acies, condyli, articuli.

the Tendons of the Extensors, Lumbricales, and Interossei, and placed directly backwards, for their greater Strength; and their flat concave Part is forwards, for taking Hold more surely, and for lodging the Tendons of the Flexor Muscles.—The Ligaments for keeping down these Tendons are fixed to the Angles that are between the convex and concave Sides.

The Bones of the first *Phalanx* + of the Fingers answer to the Description of the second Bone of the Thumb: Only that the Cavity in their Base is not so oblong; nor is their Motion on the metacarpal Bones so much confined; for they can be moved laterally or circularly, but have no Rotation or a very small Degree of it round their Axes.

Both the Ends of this first *Phalanx* are in a cartilaginous State at the Birth; and the upper one is afterwards affixed in Form of an

Epiphyfe.

The fecond Bone t of the Fingers has its Base formed into two lateral Cavities, and a middle Protuberance; while the lower End has two lateral Protuberances and a middle Cavity; therefore it is joined at both Ends in the same Manner, which none of the Bones of the Thumb are.

This Bone is in the fame Condition with

the former in Children.

The third Bone & differs nothing from the Description

⁺ TIpoxovoususon

[‡] Kondunos.

^{\$} Μετακόκδυλοι, βιζωνέχια.

Description of the third Bone of the Thumb, excepting in the general distinguishing Marks; and therefore the second, and third *Phalanx* of the Fingers enjoy only Flexion and Extension.

The upper End of this third *Phalanx* is a Cartilage in a ripe Child; and is only an *Epi-*physe after, till the full Growth of the Bo-

dy.

All the Difference of the Phalanges of the feveral Fingers consists in their Magnitude.—
The Bones of the Middle-finger ** being the longest and largest,—those of the Fore-finger + come next to that in Thickness, but not in Length, for those of the Ring-finger ‡‡ are a little longer. The Little-finger || has the smallest Bones. Which Disposition is the best Contrivance for holding the largest Bodies; because the longest Fingers are applied to the middle largest Periphery of such Substances as are of a spherical Figure (a).

The Uses of all the Parts of our superior Extremities are so evident in the common Actions of Life, that it is needless to enumerate them here; and therefore I shall proceed to the last Part of the Skeleton. Only, lest I should seem to have forgot the small Bones at the Joints of

the

^{**} Καταπυγών, οφάκκεκος, infamis, impudicus, verpus, famosus, obscænus.

[†] Δεκτικός, indicator, λιχανός, demonstrativus, falutaris. ‡ Ἱατρικός, παράμεσος, δακτυλιώτης, ἐπιβάτης, Annularis, medicus, cordis digitus.

[|] Μύωψ, ωτίτης, Auricularis, minimus.
(a) Galen. de usu Part. lib. 1. cap. 24.

the Hand, I desire now to refer to the Description of them, under the common Title of Sefamoid Bones, which I have placed after the Bones of the Feet.

Of the Inferior Extremities.

THE INFERIOR EXTREMITIES depend from the Acetabula of the Offa innominata; and are commonly divided into three Parts, viz. the Thigh, Leg, and Foot.

The THIGH * has only one Bone; which is the longest of the Body, and the largest and strongest of any of the cylindrical Bones. The Situation of it is not perpendicular: For the lower End is inclined confiderably inwards; fo that the Knees are almost contiguous, while there is a considerable Distance between the Thigh-bones above: Which is of good Use to us, fince sufficient Space is thereby left for the external Parts of Generation, the two great Cloacæ of Urine and Faces, and for the large thick Muscles that move the Thigh inwards: And at the fame time this Situation of the Thigh-bones renders our Progression quicker, surer, streighter, and in less Room; for had the Knees been at a greater Distance from each other, we must have been obliged to describe some Part of a Circle with the Trunk of our Body in making a long Step, and, when one Leg was raifed from the Ground, our Center of Gravity would have been too far from the Base of the other, and we should confequently

it

15

n

ir

13

0

tl

te

fo

ti

^{*} Mrgov, Femen, coxa, agie, anchæ os, crus, femur,

fequently have been in Hazard of falling; for that our Steps would neither have been streight nor firm; nor would it have been possible to walk in a narrow Path, had our Thigh-bones been otherwise placed. In Consequence however of the Weight of the Body bearing so obliquely on the Joint of the Knee, by this Situation of the Thigh-bones, weak rickety Children become in-kneed.

The upper End of the Thigh-bone is not continued in a streight Line with the Body of it, but is fet off obliquely inwards and upwards, whereby the Distance here between these two Bones at their upper Part is confiderably increased. This End is formed into a large smooth round Head +, which is the greater Portion of a Sphere unequally divided.—Towards its lower internal Part a round rough spongy Pit is observable, where the strong Ligament, commonly, but unjustly, called the round one, is fixed, to be extended from thence to the lower internal Part of the receiving Cavity, where it is considerably broader than near to the Head of the Thigh-bone. The small Part below the Head, called the Cervix, of the Os Femoris, has a great many large Holes, into which the Fibres of the strong Ligament, continued from the capfular, enter, and are thereby furely united to it; and round the Root of the Neck, where it rifes from the Bone, a rough Ridge is found, where the capfular Ligament of the Articulation itself is connected.—Below the Back-part of this Root, the large unequal Pro-

[†] Vertebrum.

r

tuberance, called Trochanter major +, stands out; the external convex Part of which is dif. tinguished into three different Surfaces, where. of the one on the Fore-part is feabrous and rough, for the Insertion of the Glutaus mini. mus; the superior one is smooth, and has the Glutæus medius inferted into it; and the one be. hind is made flat and fmooth by the Tendon of the Glutæus maximus passing over it.—The upper Edge of this Process is sharp and pointed at its Back-part, where the Glutæus medius is fixed: but forwards it is more obtuse, and has two superficial Pits formed in it: Into the superior of these, the Piriformis is implanted; and the Obturator internus and Gemini are fixed into the lower one. From the backmost prominent Part of this great Trochanter, a rough Ridge runs backwards and downwards, into which the Quadratus is inferted.—In the deep Hollow, at the internal upper Side of this Ridge, the Obturator externus is implanted.— More internally, a conoid Process, called Trochanter minor t, rifes for the Infertion of the Musculus Psoas, and Iliacus internus, and the Pettineus, is implanted into a rough Hollow below its internal Root.—The Muscles inferted into these two Processes being the principal Instruments of the rotatory Motion of the Thigh, have occasioned the Name of Trochanters to the Processes.—The Tendons that are fixed into, or pass over the great Trochanter, cause Bruises by Falls on this Part to be attended with great Pain

[†] FAUTOS, Rotator natis, malum granatum testiculorum. ‡ Rotator minor.

Pain and Weakness of the Limb, which gene-

rally remain long.

ds

if.

e-

nd

i-

10

e-

of

le

d

is

15

d

0

1-

0

S

The Body of the Os femoris is convex on the Fore-part, and made hollow behind, by the Action of the Muscles that move it and the Leg, and for the Conveniency of Sitting, without bearing too much on these Muscles; and probably the Weight of the Legs depending from the Thighs in that Posture, contributes to this Curvature. The Fore-part of the Thigh-bone is a little flatted above by the Beginning of the Cruræus Muscle, as it is also below by the fame Muscle and the Rectus .--Its external Surface is likewise made flat below by the Vastus extermis, where it is separated from the former by an obtuse Ridge.—The Vastus internus depresses a little the lower Part of the internal Surface.—The posterior concave Surface has a Ridge rifing in its Middle, commonly called Linea aspera, into which the Triceps is inserted, and the short Head of the Biceps flexor tibiæ rises from it.—At the upper Part of it the medullary Vessels enter by a small Hole that runs obliquely upwards. A little above which there is a rough Fossa or two, where the Tendon of the Glutæus maximus is fixed .-The lower End of the Linea aspera divides into two, which descend toward each Side. The two vasti Muscles have Part of their Origin from these Ridges; and the long Tendon of the Triceps is fixed to the internal, by means of Part of the Fascia aponeurotica of the Thigh. -Near the Beginning of the internal Ridge, there is a Discontinuation of the Ridge, where

the crural Artery passes through the Aponeuross.—Between these two rough Lines, the Bone is made flat by the large Blood-vessels and Nerves which pass upon it; and near the End of each of these Ridges, a small smooth Protuberance may often be remarked, where the two Heads of the external Gastrochemius Muscle take their Rise, and where sesamoid Bones are sometimes sound (a); and from the Fore-part of the internal Tubercle, a strong Ligament is extend.

ed to the Infide of the Tibia.

The lower End of the Os femoris is larger than any other Part of it, and is formed into a great Protuberance on each Side, called its Condyles; between which a confiderable Cavity is found, especially at the Back-part, in which the crural Vessels and Nerves ly immersed in -The internal Condyle is longer than the external, which must happen from the oblique Position of this Bone, to give less Obliquity to the Leg.—Each of these Proces. fes feems to be divided in its plain fmooth Surface. The Mark of Division on the external is a Notch, and on the internal a small Protuberance. The Fore-part of this Division, on which the Rotula moves, is formed like a Pulley, the external Side of which is highest .-Behind, there are two oblong large Heads, whose greatest Extent is backwards, for the Motion of the Tibia; and from the rough Cavity between them, but near to the Base of the internal Condyle, the strong Ligament commonly called the cross one, has its Rife.—A little

⁽a) Vesal. Lib. 1. cap. 28. & 30.

little above which a rough Protuberance gives Infertion to the Tendon of the Triceps. The Condyles, both on the Outer and Inner Side of the Knee, are made flat by the Muscles passing along them. On the Back-part of the internal, a flight Depression is made by the Tendons of the Gracilis and Sartorius; and on the external fuch another is formed by the Biceps flexor cruris; behind which a deep Fossa is to be observed, where the Poplitæus Muscle has its Origin.-From a Tubercle immediately before this Cavity, a strong round Ligament goes out to the upper Part of the Fibula .-Round this lower End of the Thigh-bone, large Holes are found, into which the Ligaments for the Security of the Joint are fixed, and Bloodvessels pass to the internal Substance of the Bone.

All the Processes of the Fenuer are cartilaginous in new-born Children, and afterwards become small Apophyses, with large Epiphy-

fes.

ne

nd

nd

tu-

ke

le-

he

id.

to

its

ty

in

an

ne -

h

0.

1.

e

The Thigh-bone is articulated above with the Acetabulum of the Ossa immominata, which affords its round Head a secure and extensive Play, and therefore can be moved to every Side; but is restrained in its Motion outwards, by the high Brims of the Cavity, and by the round Ligament; for otherwise the Head of the Bone would have been frequently thrust out at the Breach of the Brims on the Inside, which allows the Thigh to move considerably inwards.— The Body of this Bone enjoys little or no rotatory Motion, though the Head most commonly moves rounds its own

Axis; because the oblique Progress of the Neck and Head from the Bone is such, that the rotatory Motion of the Head can only bring the Body of the Bone forwards and backwards: Nor is this Head, as in the Arm, ever capable of being brought to a streight Direction with its Body; so far however as the Head can move within the Cavity backwards and forwards, the rest of the Bone may have a partial Rotation.—The Os femoris is articulated below to the Tibia and Rotula in the Manner afterwards to be described.

The Nearness of the small Neck to the round Head of the Thigh-bone, and its upper End being covered with very thick Muscles, make greater Difficulty in distinguishing between a Luxation and Fracture here,

than in any other Part of the Body.

The LEG* is composed, according to the common Account, of two Bones, Tibia and Fibula, though it seems to have a very good Title to a third, the Rotula; which bears a strong Analogy to the Olecranon of the Ulna, and moves always with the other two.

TIBIA‡, fo called from its Resemblance to an old musical Pipe or Flute, is the long thick triangular Bone, situated at the internal Part of the Leg, and continued in almost a

streight Line from the Thigh-bone.

The upper End of the Tibia is large, bulbous, and spongy, and is divided into two Cavities,

^{*} Kvnµn, Crus, tibia.

[‡] Προκνήμιον, εντικνήμιον, focile majus, arundo major, canna major, canna domestica cruris.

1

.

d

d

e

d

a

O

Cavities, by a rough irregular Protuberance +, which is hollow at its most prominent Part, as well as before and behind. The anterior of the two Ligaments that compose the great cross one, is inserted into the middle Cavity, and the Depression behind receives the posterior Ligament. The two broad Cavities at the Sides of this Protuberance are not equal; for the internal is oblong and deep, to receive the internal Condyle of the Thighbone; while the external is more superficial and rounder, for the external Condyle. -In each of those two Cavities of a recent Subject, a femilunar Cartilage is placed, which is thick at its convex Edge, and becomes gradually thinner towards the concave or interior Edge. The Middle of each of these Cartilages is broad, and the Ends of them turn narrower and thinner, as they approach the middle Protuberance of the Tibia. — The thick convex Edge of each Cartilage is connected to the capiular and other Ligaments of the Articulation, but so near to their Rise from the Tibia, that the Cartilages are not allowed to change Place far; while the narrow Ends of the Cartilages becoming almost Ligaments, are fixed at the Infertion of the strong crois Ligament into the Tilia, and feem to have their Substance united with it; therefore a circular Hole is lest between each Cartilage and the Ligament, in which the most prominent convex Part of each Condyle of the Thighbone, moves.—The Circumference of these A a 2 Cavities

[†] Διάρυσις, έξοχη νευροχονδρώδης, Tuber, Tuberculum.

Cavities is rough and unequal for the firm Connexion of the Ligaments of the Joint.—
Immediately below the Edge at its Back-part, two rough flatted Protuberances stand out: Into the internal, the Tendon of the feminembranofus Muscle is inserted; and a Part of the cross Ligament is fixed to the external.—On the Outside of this last Tubercle, a smooth slightly-hollowed Surface is formed by

the Action of the Poplitaus Muscle.

Below the Fore-part of the upper End of the Tibia, a confiderable rough Protuberance + rifes, to which the strong tendinous Ligament of the Rotula is fixed. On the internal Side of this, there is a broad scabrous flightly-hollowed Surface, to which the internal long Ligament of the Joint, the Aponeurosis of the Vastus internus, and the Tendons of the Seninervofus, Gracilis and Sartorius, are fixed. The lowest Part of this Surface is therefore the Place where the Tibia ought to be fawed through in an Amputation, fo as not to have too long and troublesome a Stump, and, at the fame time, to preserve its Motions, by faving the proper Muscles. -Below the external Edge of the upper End of the Tibia, there is a circular flat Surface covered in a recent Subject with Cartilage, for the Articulation of the Fibula; between which and the anterior Knob, there is a rough Hollow from which the Tibialis anticus, and Extensor digitorum longus, take their 0rigin. --- From the smooth flat Surface, a Ridge

^{† &#}x27;Avrixvymiev, anterior Tuber.

Ridge runs obliquely downwards and inwards, to give Rife to Part of the Solæus, Tibialis poficus, and Flexor digitorum longus, and Infertion to the Aponeurolis of the Semimembranofus which covers the Poplitæus, and to some of. the external Fibres of this last named Muscle. -At the Infide of this Ridge, an oblique plain Surface is left, where the greatest Part of the Musculus Poplitaus is inserted. The remaining Body of the Tibia is triangular. The anterior Angle is very sharp, and is commonly called the Spine or Shin ‡. This Ridge is not streight; but turns first inwards, then: outwards, and lastly inwards again. - The plain internal Side is smooth and equal, being little subjected to the Actions of Muscles; but the external Side is hollowed above by the Tibialis anticus, and below by the Extensor digitorum longus and Extensor pollicis longus. The two Angles behind these Sides are rounded by the Action of the Muscles:- the posterior Side comprehended between them, is not fo broad as those already mentioned, but is more oblique and flatted by the Action, of the Tibialis posticus and Flexor digitoriun longus. - Some Way above the Middle of the Bone, the internal Angle terminates, and the Bone is made round by the Pressure of the Musculus solaus.—Near to this, the Passage of the medullary Vessels is feen flanting obliquely downwards.

The lower End of the Tibia is made hollow, but so as a small Protuberance rises in the A a 3 Middle.

[†] Ακανθα, Spina, crea, linea prima ti siæ, angulus acutus.

Middle.—The internal Side of this Cavity. which is smooth, and, in a recent Subject, is covered with Cartilage, is produced into a confiderable Process, commonly named Malleolus internus +; the Point of which is divided by a Notch, and from it Ligaments are fent out to the Foot. -- We ought to obferve here, that this internal Malleolus is fituated more forwards than the internal Condyle of the upper End of this Bone; which is necessary to be remembered in reducing a Fracture of the Leg (a).—The external Side of this End of the Tibia has a rough irregular femilunar Cavity formed in it, for receiving the lower End of the Fibula. The posterior Side has two lateral Grooves, and a finall middle Protuberance. In the internal Depression, the Tendons of the Musculus tilialis posticus and Flexor digitorum longus are lodged; and in the external, the Tendon of the Flexor longus poliicis plays. From the middle Protuberance, ligamentous Sheaths go out, for tying down these Tendons.

The Articulations and Motions of the Tibia shall be explained, after all the three Bones

of the Leg are described.

Both the Ends of the Tibia are Cartilages at the Birth, and become afterwards Epiphyfes.

FIBULA

[†] Doupov, wicov, Talus, clavicula, clavilla interior, eavilla

⁽a) Winflow, Exposition Anatomique, des Os secs, Sect.

the

FIBULA* is the small long Bone, placed on the Outside of the Leg, opposite to the external Angle of the Tibia; the Shape

of it is irregularly triangular.

:1-

d-

re

b-

n-

15

c. of

ar

ng

ri-

all

lelis

he id-

ut,

Ti-

ies

res

by-

A

illa

.B.

The Head of the Fibula has a superficial circular Cavity formed on its Inside, which, in a recent Subject, is covered with a Cartilage, but so closely connected to the Tibia by Ligaments, as to allow only a small Motion backwards and forwards.—This Head is protuberant and rough on its Outside, where a strong round Ligament and the Musculus Biceps are inserted; and, below the Backpart of its internal Side, a Tubercle may be remarked, that gives Rise to the strong tendinous Part of the Solæus Muscle.

The Body of this Bone is a little crooked inwards and backwards, which Figure is owing to the Actions of the Muscles; but is still further increased by Nurses, who often hold Children carelely by the Legs.—The sharpest Angle of the Fibula is forwards, on each Side of which, the Bone is confiderably, but unequally depressed by the Bellies of the feveral Muscles that rise from, or act upon it; and, in old People, these Muscles make distinct Sinuosities for themselves .- The external Surface of the Fibula is depressed obliquely from above downwards and backwards, by the two Peronei. — Its internal Surface is unequally divided into two narrow longitudinal Planes, by an oblique Ridge extended from

^{*} Παρακνήμιον, Perone, Focile minus, arundo minor, canna minor cruris, fura, radius.

the upper Part of the anterior Angle, to join with the lower End of the internal Angle, To this Ridge the Ligament stretched between the two Bones of the Leg is connected .-The anterior of the two Planes is very narrow above, where the Extensor longus digitorum and Extensor longus pollicis arise from it; but is broader below, where it has the Print of the Nomus Vefalii. The posterior Plane is broad and hollow, giving Origin to the larger Share of the Tibialis posticus. The internal Angle of this Bone has a tendinous Membrane fixed to it, from which Fibres of the Flexor digitorum longus take their Rife. -The posterior Surface of the Fibula is the plainest and smoothest, but is made flat a bove by the Solaris, and is hollowed below by the Flexor pollicis longus. -- In the Middle of this Surface the Canal for the medullary Vessels may be seen slanting downwards.

I have taken particular Notice of the Entry and Direction of the medullary Vessels of the large Bones of the Extremities (a); because, in several chirurgical Cases, a Surgeon, who is ignorant of this, may do Mischief to his Patient. Thus, for Example, if these Vessels are opened very near to their Entry into the Bone, or while they are in the oblique Passage through it, an obstinate Hæmorrhagy may ensue: For the Arteries being connected to the bony Passage, Styptics, and other like Corrugators, are vainly applied; compressing Instruments can do no Service, and Ligatures can

11

2

W

d

tl

tl

be

C

T

not

⁽a) Havers, Ofteolog. nov. Difc. 1. p. 59.

nic

le.

en

ar-

gi-

om

he

101

to

en-

res.

fe.

is

2.

by

dle

lry

n-

0

be-

on,

to

ef-

ito

ue

ay

he

01.

[ŋ.

ın-

ot

not be employed. There feems to be a particular Design in the Contrivance of these Canals; those in the Os humeri, Tibia, and Fibula, running obliquely downwards from their external Entry; whereas in the Radius, Ulna, and Os femoris, they slant upwards, whereby the Arteries and Nerves which are lent into these three last Bones, must suffer a confiderable Reflexion before they come the Cancelli. The Reason of this Diversity may perhaps be, that the Arteries which are so small within the Bones as to have no strong contractile propelling Force in their Coats, and where they are not affished by the Action of any moving neighbouring Organ, should have, at least in their Passage through the Bone, a favourable Descent for their Liquids: Which, it is evident, they have in the descending oblique Passages formed for them in the first Class of Bones, to wit, the Os humeri, Tibia and Fibula, which are generally depending; and they also most frequently acquire the like Advantage in the Radius, Ulna, and Os femoris, because the Hand, in the most natural Posture, is higher than the Elbow; and when we fit or lie, the lower End of the Thigh-bone comes to be at least as high raised as the upper. In standing and walking, or when the Arms are moved, the Blood must indeed ascend as it passes through the Bones of the Fore-arm and Thigh; but the Pressure of the Muscles, then in Action, on the Vessels, before they enter the Bones, is sufficient to compensate the Disadvantage of their Course. This Reasoning scems to be still enforced, by observing,

observing, that this Passage is always nearer the upper than the lower Ends of these Bones.

The lower End of the Fibula is extended into a spongy oblong Head, on the Inside of which is a convex, irregular, and frequently a scabrous Surface, that is received by the external Hollow of the Tibia, and fo firmly joined to it by a very thin intermediate Car. tilage and strong Ligaments, that it scarce can move.—Below this, the Fibula is stretched out into a coronoid Process, that is smooth, covered with Cartilage on its internal Side, and is there contiguous to the Outfide of the first Bone of the Foot, the Astragalus, to secure the Articulais. This Process, named Malleolus externus, being fituated farther back than the internal Malleolus, and in an oblique Direction, obliges us naturally to turn the Fore-part of the Foot outwards (a). At the lower internal Part of this Process, a spongy Cavity for mucilaginous Glands may be remarked; from its Point Ligaments are extended to the Afragalus, Os Calcis, and Os Naviculare, Bones of the Foot; and from its Infide short strong ones go out to the Altragahis. On the Back-part of it? Sinuofity is made by the Tendons of the Peronæi Muscles. -- When the Ligament extended over these Tendons from the one Side of the Depression to the other is broke, stretched too much, or made weak by a Sprain, the Tendons frequently start forwards to the Outfide of the Fibula.

The

F

110

th

10

je

8

a

⁽a) Winston, Memoires de l'Acad. des Sciences, 1722.

The Conjunction of the upper End of the Fibula with the Tibia is by plain Surfaces tipped with Cartilage, and at its lower End the Cartilage feems to glue the two Bones together, not, however, fo firmly in young People, but that the Motion at the other End of such a long Radius is very observable.-In old Subjects I often see the two Bones of the Leg

grown together at their lower Ends.

rer

ele

ded

of

tly

the

nly

ar-

can

out

red

ere

he

III-

us,

nal

es

oot

of

9119

Li-

Os

1: 30

IH

2

e-

X

de

h.

he

it.

18

The principal Use of this Bone is to afford Origin and Infertion to Muscles; the Direction of which may be a little altered on proper Occasions, by its upper Part shuffling backwards and forwards.—It likewise helps to make the Articulation of the Foot more secure and firm. The Ends of the Tibia and Fibula being larger than their Middle, a Space is here left, which is filled up with fuch another Ligament as I described extended between the Bones of the Fore-arm; and which is also discontinued at its upper Part, where the Tibialis anticus immediately adheres to the Solæus and Tibialis posticus; but every where else it gives Origin to muscular Fibres (a).

Both the Ends of this Bone are cartilaginous in a ripe Child, and assume the Form of Appen-

dices before they are united to its Body.

ROTULA * is the fmall flat Bone fituated at the Fore-part of the Joint of the Knee.— Its Shape refembles the common Figure of the Heart with its Point downwards.—The anterior

(a) Weitbrecht. Syndesmolog. p. 156.

^{*} Έπιμυλίς, μυλακρίς, κόγχος, έπιγονατίς, πλανησίεδρον, patella, mola, genu, scuti-forme os, cartilaginosum, disciforme, oculus genu.

rior convex Surface of the Rotula is pierced by a great Number of Holes, into which Fibres of the strong Ligament that is spread over it enter.—Behind, its Surface is smooth, covered with Cartilage, and divided by a middle convex Ridge into two Cavities, of which the external is largest; and both are exactly adapted to the Pulley of the Os femoris, on which they are placed in the most ordinary unstraining Po. ftures of the Leg; but when the Leg is much bended, the Rotula descends far down on the Condyles; and when the Leg is fully extended, the Rotula rifes higher, in its upper Part, than the Pulley of the Thigh-bone.—The plain fmooth Surface is furrounded by a rough prominent Edge, to which the capfular Ligament adheres:—Below, the Point of the Bone is scabrous, where the strong tendinous Ligament from the Tubercle of the Tibia is fixed.—The upper horizontal Part of this Bone is flatted and unequal, where the Tendons of the Extenfors of the Leg are inferted.

The Substance of the Rotula is cellular, with very thin external firm Plates: But then these Cells are so similar, and such a Quantity of Bone is employed in their Formation, that scarce any Bone of its Bulk is so strong. Besides, it is covered all over with a thick Ligament, (as it was observed, that this Sort of Bones generally is), to connect its Substance, and is moveable to one Side or other; therefore is sufficiently strong to resist the ordinary Actions of the large Muscles that are inserted into it, or any common external Force applied to it; while a fixed Process, such as the Olecranon, would not have

t

fi

b

been

Y

al

10

ne

d,

In

in

14

d.

a.

nt

le

1d

rs

th

fe

10

14

0-

as

5),

to ly

ned

ve

been sufficient to bear the whole Weight of our Bodies, which frequently fall on it, and would have hindered the rotatory Motion of the Leg. Notwithstanding these Precautions to preserve this Bone from fuch Injuries, yet I have feen a transverse Fracture in it, when, by the Report of the Patient, and of the People about him, and by the Want of Swelling, Discolouring, or other Mark of Bruise or Contusion, it was plain the Bone was broken by the violent straining Effort of the Muscles (a). Though my Patient recovered the Use of the Joint of the Knee, yet I think it reasonable to believe, that this Sort of Fracture is commonly attended with Difficulty of Motion, after the broken Parts of the Rotula are reunited, because the callous Matter probably extends itself into the Cavity of the Joint, where it either grows to some of the Parts, or makes fuch an Inequality on the Surface of this Bone, as does not allow it to perform the necessary Motions on the Condyles of the Femur (b).

At the ordinary Time of Birth, the Rotula is entirely cartilaginous, and scarcely assumes a bony Nature so soon as most Epiphyses do.

The Parts which constitute the Joint of the Knee being now described, let us examine what are its Motions, and how performed.—The two principal Motions are Flexion and Extension—In the former of these, the Leg may be brought to a very acute Angle with the Thigh, by the Condyles of the Thigh-bones B b

⁽a) See Ruysch. Observ. Anat. Chirurg. Obs. 3.

⁽b) Pare, Liv. 15. cap. 22.

being round and made fmooth far backwards. In performing this, the Rotula is pulled down by the Tibia.—When the Leg is to be extended. the Rotula is drawn upwards, and confequent. ly the Tibia forwards, by the Extensor Muscles: which, by Means of the protuberant Joint, and of this thick Bone with its Ligament, have in 'effect the Chord, with which they act, fixed to the Tibia at a confiderable Angle, therefore aft with Advantage; but are restrained from pulling the Leg farther than to a streight Line with the Thigh, by the posterior Part of the cross Liga. ment, that the Body might be supported by a firm perpendicular Column: For at this Time the Thigh and Leg are as little moveable in a rotatory Way, or to either Side, as if they were one continued Bone.—But when the Joint is a little bended, the Rotula is not tightly braced, and the posterior Ligament is relaxed; therefore this Bone may be moved a little to either Side, or with a finall Rotation in the superficial Cavities of the Tibia; which is done by the Motion of the external Cavity backwards and forwards, the internal ferving as a fort of Axis (a). Seeing then one Part of the cross Ligament is fituated perpendicularly, and the posterior Part is stretched obliquely from the internal Condyle of the Thigh outwards, that posterior Part of the cross Ligament prevents the Leg's being turned at all inwards; but it could not hinder it from turning outwards almost round, was not that Motion confined by the lateral Ligaments

in To Su

⁽a) Winflow, Exposition Anatomique du corps humain, traité des Os secs, § 976.

of this Joint, which can yield little. This Rotation of the Leg outwards, is of good Advantage to us in croffing our Legs, and turning our Feet outwards, on feveral necessary Occasions; tho' it is altogether fit this Motion should not be very large, to prevent frequent Luxations here. - While all these Motions are performing, the Part of the Tibia that moves immediately on the Condyles is only fo much as is within the cartilaginous Rings, which, by the Thicknels on their Outsides, make the Cavities of the Tibia more horizontal, by raising their external Side where the Surface of the Tibia flants By this Means the Motions of downwards. this Joint are more equal and steady than otherwife they would have been. The Cartilages being capable of changing a little their Situation, are fit for doing this good Office in the different Motions and Postures of the Member, and likewife contribute to make the Motions larger and quicker.

On Account of the very large Surface of the Bones forming the Joint of the Knee, and the many strong Ligaments connecting them, Luxations seldom happen here. But these very Ligaments, the Aponeuroses passing over this Joint, the Quantity of Fat and mucaginous Glands necessary for lubricating it, make it more subject to White-fwellings, Dropsies, and such other Disorders, than any other Joint of

the Body.

d

in

le

is

e.

al

0-

1.

15

rt

le

of

er

ts

of

ite

The FOOT is divided, as well as the Hand, into three Parts, viz. Tarfus, Metatarfus, and, Toes: In the Description of which, the several Surfaces shall be named, according to their named.

B b 2

tural Situation, viz. the Broad of the Foot, thall be called superior; the Sole, inferior; the Side on which the Great-toe is, internal; that where the Little-toe is, external.

The Tarfus * confifts of feven spongy Bones; to wit, the Astragalus, Os calcis, Naviculare, Cuboides, Cuneiforme externum, Cuneiforme me.

dium, and Cuneiforme internum.

The Astragalus is the uppermost of these Bones.—The Os calcis is below the Astragalus, and is considerably prominent backwards beyond the other Bones of the Tarsus, to form the Heel.—The Os naviculare is in the Middle of the internal Side of the Tarsus.—The Os cuboides is the most external of the Row of sour Bones at the Fore-part of the Tarsus.—The Os cuneiforme externum is placed at the Inside of the Cuboid.—The Cuneiforme medium is between the external and internal cuneiform Bones, and the internal Cuneiform is put at the internal Side of the Foot.

That the Description of these Bones may not be immoderately swelled with Repetition, I desire, once for all, to observe, That where-ever a Ridge is mentioned, without a particular Use assigned, a Ligament is understood to be fixed to it; or where a spongy rough Cavity, Depression or Fossa is remarked, without naming its Use, a Ligament is inserted, and mucilaginous Glands are lodged: For such will occur in

the Detail of each of these Bones.

The

^{*} Raffetta.

The upper Part of the Astragalus + is formed into a large smooth Head ‡, which is slightly hollowed in the Middle; and therefore resembles a superficial Pulley, by which it is sitted to the lower End of the Tibia.—The internal Side of this Head is slat and smooth, to play on the internal Malleolus.—The external Side has also such a Surface, but larger, for its Articulation with the external Malleolus.—Round the Base of this Head there is a rough Fossa; and, immediately before the Head, as also below its internal smooth Surface, we find a con-

fiderable rough Cavity.

it

e

n

r

e

e

S

C

The lower Surface of the Altragalus is divided by an irregular deep rough Fossa; which at its internal End is narrow, but gradually widens, as it stretches obliquely outwards and forwards. -- The fmooth Surface, covered with Cartilage, behind this Fossa, is large, oblong, extended in the fame oblique Situation with the Fossa, and concave, for its Conjunction with the Os calcis. The Back-part of the Edge of this Cavity is produced into two sharp-pointed rough Processes, between which is a Depression made by the Tendon of the Flexor pollicis longus .--The lower Surface before the Fossa is convex,. and composed of three distinct smooth Planes. The long one behind, and the exterior or shortelt, are articulated with the Heel-bone; while the internal, which is the most convex of the three, rests and moves upon a cartilaginous Ligament, B b 3

[†] Aspioc, Talus, balistæ os, malleolus, chaib, quatrio, Ostessaræ, claviculæ, nuciforme.

[‡] Τετρώρος.

that is continued from the Calcaneum to the Os fcaphoides. Without which Ligament, the Astragalus could not be sustained, but would be pressed out of its Place by the great Weight it supports, and the other Bones of the Tarsus would be separated. Nor would a Bone be sit here, because it must have been thicker than could conveniently be allowed; otherwise it would break, and would not prove such an easy bending Base, to lessen the Shock which is given to the Body in leaping, running, &c.

The Fore-part of this Bone is formed into a convex oblong smooth Head, called by some its Process, which is received by the Os naviculare.

—Round the Root of this Head, especially on the upper Surface, a rough Fossa may be re-

marked.

The Altragalus is articulated above to the Tibia and Filula, which together form one Though, in this Articulation, the Cavity. Bones have Prominencies and Cavities for fmall, as might allow Motions in all Directions, yet the Flexion and Extension are the most confiderable, the other Motions being confined by the Malleoli, and by the strong Ligaments which go out from the Points of these Processes to the Astragalus and Os calcis. - When the Foot is bended, to far as it is commonly when we stand, no lateral or rotatory Motion is allowed in this joint; for then the Head of the Aftragalus is funk deep between the Malleoli, and the Ligaments are tenfe; but when the Foot is extended, the Astragalus can move a little to either Side, and with a small Rotation. By this Contrivance the Foot is firm, when the Weight

Weight of the Body is to be supported on it; and when a Foot is raised, we are at Liberty to direct it more exactly to the Place we intend next to step upon.—The Astragalus is joined, below, to the Os calcis; and before, to the Os naviculare, in the Manner to be explained, when these Bones are described.

A confiderable Share of this Bone is offified

in a new-born Infant.

e

it

it

n

it

n

Calcaneum * is the largest Bone of the seven. -Behind, it is formed into a large Knob, commonly called the Heel: The Surface of which is rough behind, where the Tendo Achillis is inferted into it; and above, it is hollow and fpongy. Farther forwards, on the upper Surface of the Calcaneum, there is an irregular oblong smooth Convexity, adapted to the Concavity at the Back-part of the Astragalus: And beyond this a narrow Fossa is seen, which divides it from two small concave smooth Surfaces, that are joined to the Fore-part of the Aftragalus.—Behind the posterior of these smooth Surfaces, which is the largest, a small Sinuosity is made by the Tendon of the Flexor digitorum longus; at the Fore-part of which a small rough Protuberance appears, that gives Rife to the Musculus extensor digitorum brevis.

The external Side of this Bone is flat, with a superficial Fossa running horizontally, in which the Tendon of the Musculus peronaus longus is lodged.—The internal Side of the Heel-bone is hollowed, for lodging the Origin of the Massa carnea Jac. Sylvii, and for the safe Passage of

Tendons,

^{*} Os calcis, πτέρνα, calcar pedis.

Tendons, Nerves, and Arteries.—Under the Side of the internal smooth Concavity, a particular Groove is made by the Tendon of the Flexor pollicis longus; and from the thin Protuberance on this internal Side, the cartilagin. ous Ligament that supports the Astragalus, goes out to the Os naviculare; on which Ligament, and on the Edge of this Bone to which it is fixed, the Groove is formed for the Tendon of

the Flexor digitorum profundus.

The lower Surface of this Bone is pressed state at the Back-part, by the Weight of our Bodies; and immediately before this Plane, there are two Tubercles, from the internal of which the Musculus abductor pollicis, Flexor digitorum sublimis, as also part of the Aponeurosis plantaris, and of the Abductor minimi digiti, have their Origin; and the other Part of the Abductor minimi digiti and Aponeurosis plantaris rises from the external.——Before these Protuberances this Bone is concave, for lodging the Flexormuscles; and at its Fore-part we may observe a rough Depression, from which, and a Tubercle behind it, the Ligament goes out that prevents this Bone to be separated from the Os Cubeides.

The Fore-part of the Os calcis is formed into an oblong Pulley-like fmooth Surface, which is circular at its upper external End, but is pointed below. This finooth Surface is fitted to the

Os cuboides.

Though the Surfaces by which the Aftragalus and Os Calcis are articulated, feem fit enough for Motion; yet the very strong Ligaments by which these Bones are connected, prevent it, and render this principal Part of our Base, which

which rests on the Ground, to wit, the Os calcis, firm.

A large Share of the Heel-bone is oflified at the ordinary Time of Birth, and the large Knob appears afterwards in Form of an Epi-

physe.

the

rti-

the

10.

in-

oes

nt,

15

of

lat

es;

ire

he

ub-

is,

115

ni:

m

cs

or.

2

le

118

is.

10.

is

it-

18

us

h

it,

h

Os naviculare ‡, is fomewhat circular.-It is formed into an oblong Concavity behind for receiving the anterior Head of the Astragalus. —— On the upper Surface there is a rough Fossa.—Below, the Os naviculare is very unequal and rough; but hollow for the Safety of the Muscles. On its Inside a large Knob rises out, from which the Abductor pollicis takes in Part its Origin, and the Tendon of the Tibialis posticus is inserted into it: And to it two remarkable Ligaments are fixed; the first is the strong one, formerly mentioned, which supports the Astragalus; the second is stretched from this Bone obliquely cross the Foot, to the metatarfal Bones of the middle Toe, and of the Toe next to the little one. On the Outside of the Os naviculare there is a semicircular smooth Surface, where it is joined to the Os cuboides.—The Fore-part of this Bone is all covered with Cartilage, and is divided into three smooth Planes, fitted to the three Offa cuneiformia.

The Os naviculare and Astragalus are joined as a Ball and Socket, and the Naviculare moves in all Directions in turning the Toes in-

wards.

[†] Exaposions, Os cymba.

wards, or in raising or depressing either Side of the Foot, though the Motions are greatly restrained by the Ligaments which connect this to the other Bones of the Tarsus.—A Weakness of these Ligaments causes sometimes an unnatural Turn of the Fore-part of the Foot inwards.

The Os naviculare is wholly cartilaginous

in a new-born Infant.

OS CUBOIDES t is a very irregular Cube.—Behind, it is formed into an oblong unequal Concavity, adapted to the Fore-part of the Os calcis. — On its internal Side, there is a fmall femicircular fmooth Cavity, to join the Os naviculare; immediately before which, an oblong fmooth Plane is made by the Os cuneiforme externum. - Below this, the Bone is hollow and rough. On the internal Side of the lower Surface, a round Protuberance and Fossa are found, where the Musculus adductor pollicis has its Origin. the external Side of this fame Surface, there is a round Knob, covered with Cartilage; immediately before which, a smooth Foffa may be observed, in which the Tendon of the Peronæus primus runs obliquely cross the Foot; and on the Knob, the thin flat Cartilage proper to this Muscle plays; in Place of which fometimes a Bone is found:--More externally than the Knob, a rough Hollow is made, for the strong Ligaments stretched betwixt this Bone and the Os calcis.fore,

tl

21

15

[‡] Πολύμορφον, cubiforme, quadratum, grandinosium, varium, tessaræ, multiforme.

fore, the Surface of the Os cuboides is flat, smooth, and slightly divided into two Planes, for sustaining the Os metatarsi of the little Toe, and of the Toe next to it.

The Form of the Back-part of the Os culoides, and the Ligaments connecting the Joint there with the Os calcis, both concur in allowing little Motion in this Part.

The Offification of this Bone is scarcely be-

gun at the Birth.

ľ

e

V

e

d

e

n

;

16

le

r.

ce

W

ed

e-

e,

ım,

Os cuneiforme externum *, if we regard its Situation or medium by its Bulk, is much of the Shape of a Wedge, being broad and flat above, with long Sides running obliquely downwards, and terminating in a sharp Edge. -The upper Surface of this Bone is an oblong Square. The one behind is nearly a Triangle, but not compleat at the inferior Angle, and is joined to the Os naviculare .-The external Side is an oblong Square divided as it were by a Diagonal; the upper Half of it is fmooth, for its Conjunction with the Os cuboides: The other is a scabrous Hollow, and in its superior anterior Angle as mall smooth Impression is made by the Os metatarsi of the Toe next to the little one. — The ternal Side of this Bone is also quadrangular, with the Fore-part of its Edge made flat and smooth by the Os metatersi of the Toe next to the great one, and the Back-part is also flat and smooth where the Os cuneiforme medium is contiguous to it. The Fore-part of this Bone

^{*} Chalcoideum externum.

Bone is an oblong Triangle, for fustaining the

Os metatarsi of the middle Toe.

Os cuneiforme medium, or minimum, is still more exactly the Shape of a Wedge than the former.—Its upper Part is Square;—its internal Side has a flat smooth Surface above and behind, for its Conjunction with the sollowing Bone; with a small rough Fossa below; and a considerable Share of it is rough and hollow.—The external Side is smooth and a little hollowed, where it is contiguous to the last described Bone.—Behind, this Bone is triangular, where it is articulated with the Os naviculare; and it is also triangular at its Fore-part, where it is contiguous to the Os metatarsi of the Toe next to the great one.

Os cuneiforme maximum, or internum, differs from the two former in its Situation, which is more oblique than their's.—Besides, its broad thick Part is placed below, and the finall thin Point is above and outwards; while its under broad Surface is concave, for allowing a fafe Passage to the Flexors of the great Toe. - The Surface of this Os cuneiforme behind, where it is joined to the Os naviculare, is hollow, fmooth, and of a circular Figure below, but pointed above. - The external Side confifts of two smooth and flat Surfaces, whose Direction is nearly at right Angles with each other. With the posterior, that runs obliquely from below forwards and upwards, the Os cuneiforme minimum is joined; and with the anterior, whose Direction is longitudinal, the Os metatarsi of the Toe next to the great one is connected. The Fore-

part

part of this Bone is femilunar, but flat and smooth, for sustaining the Os metatarsi of the great Toe.—The internal Side is scabrous, with two remarkable Tubercles below, from which the Musculus abductor pollicis rises; and the Tibialis anticus is inserted into its upper Part.

The three cuneiform Bones are all so secured by Ligaments, that very little Motion is allowed in any of them, and they are carti-

laginous in a Fætus of nine Months.

S

S

5

ıt

These seven Bones of the Tarfus, when joined, are convex above, and leave a Concavity below, for lodging fafely the feveral Muscles, Tendons, and Vessels that lie in the Sole of the Foot. —— In the recent Subject, their upper and lower Surfaces are covered with strong Ligaments, which adhere firmly to them, and all the Bones are fo tightly connected by these and the other Ligaments, which are fixed to the rough Ridges and Fossa mentioned in the preceeding Description of the particular Bones, that, notwithstanding the many Surfaces covered with Cartilage, some of which are of the Form of the very moveable Articulations, no more Motion is here allowed, than only to prevent too great a Shock of the Fabric of the Body in walking, leaping, &c. by falling on too folid a Base; which, if it was one continued Bone, would likewise be much more liable to be broken; and, in order to make our Foot accommodate itself to the Surfaces we tread on, by becoming more or less hollow, or by raising or depresfing either Side of it, as might be judged by what was faid of the particular Bones. Sprains Sprains here occasion, as in the Wrist, great Pain and obstinate Tumors, which too

often cause carious Bones.

METATARSUS * is composed of five Bones, which, in their general Characters, agree with the metacarpal Bones; but may be distinguished from them by the following Marks: 1. They are longer, thicker, and stronger. 2. Their anterior round Ends are not so broad, and are less in Proportion to their Bases. 3. Their Bodies are sharper above, and slatter on the Sides, with their inferior Ridge inclined more to the Outside. 4. The Tubercles at the lower Parts of the round Head are larger.

The first or internal metatarsal Bone is easily distinguished from the rest by its Thickness.—The one next to it is the longest, and with its sharp Edge almost perpendicular.—
The others are shorter and more oblique, as their Situation is more external. Which general Remarks, with the Description I am now to give of each, may teach us to distin-

guish them from each other.

Os metatarsi pollicis is by far the thickest and strongest, as having much the greatest Weight to sustain. Its Base is oblong, irregularly concave, and of a semilunar Figure, to be adapted to the Os cuneiforme maximum.—

The inferior Edge of this Base is a little prominent and rough, where the Tendon of the Peronæus princus Muscle is inserted.——On

^{*} Στηθος, πεδίον, Planta, planum, vestigium, solium, petus, præcordium, pectusculum.

its Outside an oblique circular Depression is made by the second metatarsal Bone.—Its round Head has generally on its Fore-part a middle Ridge, and two oblong Cavities, for the Ossa sejamoidea; and on the external Side a Depression is made by the following Bone.

Os metatarsi of the second Toe, is the longest of the five, with a triangular Base supported by the Os cuneiforme medium, and the external Side produced into a Process; the End of which is an oblique fmooth Plane, joined to the Os cuneifornie externum.-Near the internal Edge of the Base, this Bone has two small Depressions, made by the Os cuneiforme maximum, between which is a rough Cavity. —Farther forwards we may observe a smooth Protuberance, which is joined to the foregoing Bone. On the Outside of the Base are two oblong smooth Surfaces, for its Articulation with the following Bone; the superior fmooth Surface being extended longitudinally, and the inferior perpendicularly; between which there is a rough Fossa.

Os metatarsi of the middle Toe, is the second in Length.——Its Base, supported by the Os cuneiforme externum, is triangular, but slanting outwards, where it ends in a sharp-pointed little Process; and the Angle below is

not compleated.

The internal Side of this Base is adapted to the preceeding Bone; and the external Side has also two smooth Surfaces covered with Cartilage, but of a different Figure; for the upper one is concave, and, being round behind, turns smaller as it advances forwards;

B 2

and the lower Surface is little, smooth, convex, and very near the Edge of the Base.

Os metatarsi of the fourth Toe, is near as long as the former, with a triangular slanting Base joined to the Os cuboides, and made round at its external Angle, having one hollow smooth Surface on the Outside, where it is pressed upon by the following Bone, and two on the internal Side, corresponding to the former Bone; behind which is a long narrow Surface impressed by the Os cuneiforme externum.

Os metatarsi of the little Toe, is the shortest, situated with its two slat Sides above and below, and with the Ridges laterally. — The Base of it, Part of which rests on the Os cuboides, is very large, tuberous, and produced into a long-pointed Process externally, where Part of the Abductor minimi digiti is fixed; and into its upper Part the Peronæus secundus is inserted.—Its Inside has a flat conoidal Surface, where it is contiguous to the preceeding Bone.

When we stand, the fore Ends of these metatarsal Bones, and the Os calcis, are our only Supporters; and therefore it is necessary they should be strong, and should have a con-

fined Motion.

The Bones of the TOES are much a-kin to those of the Thumb and Fingers: Particularly the two of the great Toe are precisely formed as the two last of the Thumb; only their Position, in respect of the other Toes, is not oblique; and they are proportionally much stronger, because they are subjected to a greater

greater Force; for they Justain the Force with which our Bodies are pushed forwards by the Foot behind at every Step we make; and on them principally the Weight of the Body is supported, when we are railed on our Tiptoes.

The three Bones in each of the other four Toes, compared to those of the Fingers, differ from them in these Particulars. They are less, and smaller in Proportion to their Lengths: - Their Bases are much larger than their anterior Ends:—Their Bodies are more narrow above and below, and flatter on the Sides. — The first Phalanx is proportionally much longer than the Bones of the fecond and third, which are very short.

Of the four, the Toe next to the great one, has the largest Bones in all Dimensions, and more externally the Toes are less.—The little Toe, and frequently that next to it, have the fecond and third Bones intimately united into one; which may be owing to their little Motion, and the great Pressure they are

subjected to.

on-

as

nt-

de

ol-

it

VO

he

V/

ſt,

e-

le

es,

a

rt

0

t-

e,

e.

le.

ir

n

i-

The Toes are of good Use to us in walking; for, when the Sole is raised, they bring our Body, with its Center of Gravity, perpendicu-

lar to the advanced Foot.

The Bones of the Metatarfus and Toes, are in the fame Condition in Children as those of

the Metacarpus and Fingers.

The only Bones now remaining to complete the Descriptio: of the Skeleton, are the small ones, which are found at the Joints of the

C c 3 Fingers Fingers and Toes, and in fome other Parts

called,

OSSA SESAMOIDEA, which are of very different Figures and Sizes, though they are ge. nerally faid to refemble the Seed of the Sefa. muni. They feem to me nothing else than the Ligaments of the Articulations or the firm Tendons of strong Muscles, or both, become bony, by the Compression which they suffer. Thus the fesamoid Bones at the Beginning of the Gastrocnemii Muscles, are evidently composed of the tendinous Fibres only.—These, at the first Joint of the great Toe, are as plainly the fame continued Substance with the Ligaments and the Tendons of the Abductor, Flexor brevis, and Adductor.—That which is fometimes double at the fecond Joint of that Toe, is Part of the capfular Ligament; and if we enumerate the other fesamoid Bones that are at any time found, we may observe all of them formed in this Manner. Their Number, Figure, Situation, and Magnitude, are fo uncertain, that it were in vain to infift on the Differences of each; and therefore I shall only in general remark.

r. That where-ever the Tendons and Ligaments are firmest, the Actions of the Muscles strongest, and the Compression greatest, there

fuch Bones are most commonly found.

2. That, cateris paribus, the older the Subject is in which they are fought, their Number

is greater, and their Size is larger.

3. The more Labour any Person is inured to, he has, cateris paribus, the most numerous and largest Ossa sefamoidea.

However,

irts,

ge-

efa-

han irm

fer.

the

the

the

nts

bre-

nes art

mere, in, ces

les re

iber

o,

F. 9

However, as the two at the first Joint of the great Toe are much larger than any other, are early formed, and are feldom wanting in an Adult, we may judge, that besides the more forcible Cause of their Formation, there should also be some particular Advantage necessary at this Place, rather than elsewhere, which may possibly be, to allow the Flexor Muscles to send their Tendons along this Joint, secure from Compression in the Hollow between the two oblong fefamoid Bones; while, by removing thefe Tendons from the Center of Motion, and giving them the Advantage of an Angle at their Infertion, the Force of the Muscles is increased, and therefore the great superincumbent Weight of our Body in Progression is more easily raised.

ÁPPEN.

APPENDIX.

soft word? copyrignor about two the

Of the Marks of a FEMALE SKELETON.

O finish the Description of the Bones, is generally to conclude the Ofteology; but, that no Part of the Subject may be left untouched, I think it necessary to subjoin the di-Hinguishing Marks of the Male and Female Skeletons; and have chosen to illustrate them principally in the latter; because Women having a more delicate Constitution, and affording Lodging and Nourishment to their tender Fætuses, till they have sufficient Strength and Firmness to bear the Injuries of the Atmosphere, and Contact of other more folid Substances, their Bones are frequently incomplete, and always of a Make in some Parts of the Body different from those of the robust Male; which agree to the Description already given, unless where the proper Specialities of the Female were particularly remarked; which could not be done in all Places where they occur, without perplexing the Order of this Treatife: Therefore I chole rather to fum them up here by Way of Appendix.

The Causes of the following Specialities of the Female Bones, may be reduced to these three. V

three. 1. A weak lax Constitution. 2. A sedentary unactive Life, increasing that Constitution. 3. A proper Frame for being Mothers.

The Bones of Women are smaller in Proportion to their Length than those of Men; because the Force of their Muscles is not so great, nor is such strong external Force applied to them to prevent their stretching out in Length.

N.

15

ut,

ın-

di-

ale

em

17-

ng

u/-

m-

nd

eir

ys.

nt

to

he.

u-

in .

ng

ole

11-

of

ile.

e.

The Depressions, Ridges, scabrous Surfaces, and other Inequalities made by the Muscles, are not so conspicuous in them; because their Muscles are neither so thick nor strong, nor so much employed, to make so strong Prints on their Bones.

Their Os frontis is more frequently divided by a Continuation of the fagittal Suture, which depends on the first and second general Causes assigned above, for the Specialities in their Bones; as will appear after resecting on the Account given formerly of the middle internal Spine of this Bone.

Their Clavicles are less crooked; because their Arms have been less forcibly pulled forwards, which in our European Women, especially those of Distinction, is more hindered by their Garb.

Their Sternum is more raised by long Cartilages below, that the Thorax might be there widened in some Proportion to what it is shortened by the Pressure upon the Diaphragm, when they are with Child.

The Defect of Bone, or the Hole in the Middle of the Stermum, is oftenest found in them, to allow the Passage of the mammary

Vessels.

Vessels, say some; but, in my Opinion, this is owing to a lax Constitution, by which the Ossistation is not so soon completed as in Men, where the Action of the Solids is vigorous, and the Circulation of the Fluids is brisk; for a much smaller Hole might have served this Purpose; and the Branches of the internal mammary Vessels which are sent to the external Parts of the Thorax, do not pass here, but between the Cartilages of the Ribs, before these are joined to the Stermum.

The Cartilago Xiphoides, is oftner bifurcated in Women than Men, for the Reason assigned in the preceeding Paragraph, viz. a less for-

cible Power of Offification.

The superior Cartilages of the Ribs sooner offify, to support the Weight of the manma.

The middle Cartilages are more flat and broad by the Weight of the Breafts.

The inferior Cartilages are longer, for en-

larging the Chest.

Weak Women who have born many Children when young, often have the Vertebra of their Back bended forwards, and their Sternum depressed, or become round-shouldered and flat-breasted (a) by the Pressure and Weight of the impregnated Uterus, and by the strong Action of the abdominal Muscles.

The Os facrum is broader and turned much more backwards, for enlarging the Pelvis.

The

⁽a) Cheselden, Anatomy, book 1. chap. 3.

The Os Coccygis is more moveable, and much less bended forwards, to facilitate the Birth.

The Ossa Riven are more hollow, and more reflected outwards, and consequently further removed from each other, in order to widen the lower Part of their Abdomen, and for the better Support of the impregnated Uterus.

The Ridge on the upper Part of the Os Pubis is larger in such Women as have born Children, being extended by the strong Acti-

on of the Musculi recti abdominis.

The Cartilage between the two Offa Pubis, especially in Women who have born Children, is thicker than in Men, by which the Pelvis is

more capacious in Females.

The conjoined Surfaces of the Ossa Pubis, and of the Ossa innominata and sacrum are less, that with the streight Os sacrum a larger Passage might be lest for the Exclusion of the Child in Birth.

The great Tuberosity of the Ossa Ischium is statter in Women than in Men, because it is more pressed upon in the sedentary Life which

Females enjoy.

thie

the

Men.

and

or a

Pur.

am.

rnal

but

fore

ated

rned

for-

ner

and

en-

hil.

bræ

ter.

red ght

ong

ach

he

2.

In Consequence of the *Pelvis* of Women being wider, the Articulations of their Thighbones must be farther removed from each other; and therefore a larger Space is left for the Procreation and Birth of Children (a); which Distance of the Thighs, may be one Reason why Women in running generally shuffle more from

⁽a) Albin. De Offib. § 339.

from one Side to the other than Men, to pre ferve the Center of Gravity of their Bodies from falling too far to a Side of the Joint of the Thigh that supports them when the o. ther is raised, which would endanger their tumbling to the Ground.

F I N I S.

INDEX

A

A Cetabulum of Ossa innominata, 207.

Analysis, chemical, of Bones, 14. Phænomenon from it, 15.

Apophyses, their different Sorts, Uses, 23. Many so named

are Epiphyses, 24.

Appendices, vid. Epiphyses. Arm-bone, vid. Humeri os.

Arthrodia, 40, 43.

dies oint oheir

Articulations, 37. Symphysis, Synarthrosis, 38. Diarthrosis, 39.

42. Dispute concerning them, 40.

Astragalus, 293. Its Articulation, 294. Of Children 295.

Atlas, or the first Vertebra of the Neck, 174; of Infants, 176.

Axis, or third Vertebra Colli, 181.

B

Bones, what, 5. Their Plates, ibid. Fibres, Claviculi, 6. Cancelli, 6. Arteries, 8. Veins, 10. Nerves, ibid. Circulation of their Liquors, Nourishment, Increase, Decrease, 11. Phænomena and Diseases, 12, 21, 34. Transverse Canals, 9, 12. Longitudinal ones, 12. Analysis, 14. Phænomenon, Uses, 15. Periosteum internum and Marrow, ibid. Distinguished into broad and round, 21. Strength increased by being hollow, 22. Processes, 23. Cavities 25. Epiphyses, 26. Ossiscation, 29. Articulations, 37.

Breast-bone, vid. Sternum.

Bregma, 87.

Broad Bones, their Structure, 21.

C

Cancelli of Bones, 7. Corrugati, cribriformes, reticulares, Uses, 8.

Canini dentes, 155.

Dd

Carpus

Carpus consists of eight Bones, 255. Its Figure, 261. Substance, Articulation, 262. Motions, Uses; of Infants, 263.

Cartilages, what; their Plates, Fibres, 48. Vessels, ibid. Nerves, 49. Offiscation, how prevented, ibid. Offiscation accounted for, 50. Their Uses, 52. Diseases, 53.

Cavities of Bones, their different Kinds, Uses, 25.

Cervical Vertebra, their distinguishing Marks, 172. Atlas, 174. Dentata, 176. Axis, 181. Seventh, 182.

Cheek-bones, vid. Malarum Offa.

Clavicles, 230. Internal End, ibid. Body, 231. External End, Medullary Vessels, Substance, 232. Articulation, ibid. Uses; of Children, 233.

Claviculi of Bones, perpendicular, oblique, headed, crooked, 6. Coccygis Os. 194. Its 1st Bone, 195. Its 2d, 3d, 4th Bones, 196. Substance, Motion, Diseases, Uses, 197.

Collar bone, vid. Clavicle.

Coronal Suture, 66.

Costa, vid. Ribs.
Cranium, its Figure, 61. Surfaces, 63. Tables, 64. Diploe, 65. Consists of six proper, 65. and two common Bones,

104. Its Sutures, 66.

Cubitus, vid. Forearm.

Cuboides Os, 298. At the Birth, 299. Cuneiforme Os, of the Wrist, 257.

Cuneiformia Offa, of the Foot, Externum, medium, 299. internum, 300. At the Birth, 301.

D

DEntata, or 2d Vertebra of the Nock, its distinguishing Marks, 176; of Infants, 178.

Dentes, vid. Teeth.

Diarthrosis, 39, 42. Its three Species, Enarthrosis, 39, 41.
Artorodia, 40, 43. Ginglimus, 40, 43.

Digiti, vid. Fingers and Toes.

Diploe of the Scull, 65.

Dorfal Vertebra, their distinguishing Structure, 182. Wherein they differ from each other, 184.

E

Eniphyses, 25. Several have Apophyses, 26. Their Uses, ibid.
Ethmoidal Suture, 71.

Ethmoides

Sub. 5, 263. ibid.

Cation
Atlas.

ternal ation,

ed, 6. Bones,

iploe, ones,

nter-

hing

42.

rein

heir

ides

Ethmoides Os, 104. Its cribriform Plate, 105. Nasal Plate, 105. Cellulæ, Ossa spongiosa, 106. Connexion, 107. Uses, Morbid Phænomena; of a Child, 108.

Extremities inferior, consist, each, of Thigh, 272. Leg. 278. Foot, 291.

Arm, 240. Fore-arm, 246. Hand, 254. Uses, 271.

F

FAce, 118. Composed of upper Jaw, 119. Lower Jaw,

Female Skeleton, distinguishing Marks of it, 308.

Femoris Os, 272. Its upper End, 273. Body, 275. Lower End, 276. Of Infants, its Articulation and Motions, 277. Fibula, Its upper End, Body, 283. Inferior End, 286. Its Articulation, Uses; of Infants, 287.

Fingers, their Bones, Phalanges, 269, 1st, 2d, 3d, 270. Indicis, of middle, annular, little Finger, 271.

Fontanelle, 87.

Foot, composed of Tarsus, 292. Metatarsus, 302. Toes, 304. Its Motion, 294.

Fore-arm, 246. Confifts of Ulna, 247. Radius, 250.

Frontal Bone, 74. Its external Surface, Processes, 75, Cavities, 75. Foramina, 76. Internal Surface, 78. Foramen, 79. Substance, Sinuses, 80. Connexion, Uses; of Children, 83.

G

G Inglimus, its Species, 40, 43.
Gomphosis, 39.

H

HAnd, 254. Consists of Carpus, 255. Metacarpus, 263.

Harmonia, 39.

Haunch-bones, vid. Ilium Offa.

Head, consists of Cranium, 61. Face, 118. Its motions, 179. Heel-bone, vid. Calcis Os.

Hip bone, vid. Os Ischium.

Humeri Os, 240. Its upper End, 241. Body, 243. Lower End, ibid. Substance, Articulation, 244. Motion, 245. Of Children, 246.

Hyoides Os, 160. Its Body, ibid. Cornua, 161. Appendices, Ligaments, Substance, Connexion, Uses; of Children, 162. D d 2 Jaw

I

JAw, lower, vid. Maxilla inferior.

— upper, vid. Maxilla superior.

Riûm Ossa, 188. Spine, Dorsum, 199. External Surface, 200. Interior Surface 201. Medullary Vessels, Substance; of a Child, 202.

Incifores dentes, 154.

Innominata Osa, 198. Composed of Os Iliúm, ibid. Ischium, 202. Pubis, 205. Their great Foramen, 206. Acetabulum, 207. In Infants, 198, 209. Connexion, ibid. Phanomena, 210. Uses, ibid.

Ischium Os, 202. Process, Depressions, Tuber, 203. Substance, 204. Of Infants, 205.

L

L Ambdoid Suture, 66. Its Additamenta, ibid. Lattice work of Bones, vid. Cancelli.

Leg, composed of Tibia, 278. Fibula, 283. Rotula, 287. Its Articulation and Motions, 305.

Ligaments, 45. Their Fibres, Vessels, ibid. Nerves, Difeates, 46. Uses, Phænomena, 47.

Lumbar Vertebra, their particular Structure, 186. Differences, 187.

Lunare Os, 257.

M

Malaram Offa, their Processes, 125. Holes, Substance, Connexion, 126. Of Infants, 127.

Marrow, what, its Arteries, Veins, 16. Nerves, 17. Transverse and longitudinal Canals for conveying it, 13, 18.

Uses, ibid. Phænomena and Diseases, 18, 20.

Maxilla inferior, 142. The Chin, Sides, Base, 143. Phanomena, Angles, Processes, 144. Foramina, Substance, 145. Articulation, 146. Motions, 147. Of Infants, Uses, 148.

nexion, 120.

Macillaria Offa, their Processes, 127. Cavities, 128. Foramina, 130. Sinuses, 131. Morbid Phænomena, 132. Substance, Connexion, 133. Uses; of Children, 134.

Metacarpus, composed of four Bones, 263. Substance, at the Birth, Articulation, 264. Figure, Uses, 265. Digiti indicts, ibid. Medii, Annularis, 266. Minimi, 267.

Metatarfus, composed of five Bones, 302. Pollicis, ibid. 1, 2, 303. 3, 4, Digiti. 304.

Molares Dentes, 155.

Mucilaginous

Mucilaginous Glands, 54. Liquor secreted by them, their cellular Substance, Vessels, Nerves, Diseases, 55. Uses, 361 Phænomena, 57.

N

N Asi Ossa, their Sides, 122. Connexion, Uses; of an Infant, 123. Naviculare Os, of the Tarfus, 297. Of Children, 298. Neck, vid. Cervical Vertebra.

O Ccipitis Os, 96. Its external Surface, 197. Processes, ibid. Internal Surface, 100. Holes, 101. Substance, 102. Connexion, 103. Uses, in Infants, ibid.

Offification of Bones accounted for from particular Disposition of Vessels, 29; and from Pressure, 30. Phanomena, 28, 33.

Its History, 35.

-- of Cartilages explained, so. Osteogenea, its History, 35. Of what Use, 36, 37.

Process 125 North Land Process, 135. Substance, Connexion, 137. Uses; of Children; Diseases, 138.

Parietal Bones, their external Surface, 84. Internal Surface, 85. Substance, Connexion, 86. Uses, 87. Of Children, Practical Observations concerning Bregma, 87.

Patella, vid. Rotula.

Pelvis, 198, 211. Confifts of Os Sacrum, 189. Coccygis, 192. Offa innominata, 198.

Perichondrium, its Vessels, 48:

Periosteum externum, 1. Its Fibres, ibid. Arteries, Veins, Nerves, 3. Uses, 4. Diseases, 4. 5.

--- internum, its Structure, 15. Uses, 16.

Phalanges of the Fingers, 269. - Toes, 304.

Pifforme Os of the Wrift, 258.

Plates of Bones, their Fibres, 5.

Pollicis manus Offa, vid. Thomba

--- pedis Offa, vid. Toes.

Pores transverse and longitudinal of Bones, 8, 12. Their

Processes of Bones, vid. Apophyses.

Pubis Os, 205. Substance; of Children, 205.

Dd3

Raditts.

200. of a

hiûm, ulum. æno.

ance.

287.

Dif Diffe-

Sub-

apf-18.

hæ-145. 48. Con-

ora-Sub-

the icis,

. I,

lous.

R Adius, its upper End, 250. Body, 251. Lower End, ibid. Of Children, Articulation, 253.

Ribs. 212. Their Middle, ibid Posterior End, 213. Anterior End, Substance, Cartilages, 214. Articulation and Motion, 215. Differ from each other, 216. Diffinguifhed into true and false, 217. Distinguishing Marks of ift, ad. 3d, 4th, 5th, 6th, 7th, 11th, 12th, 219. Of Infants, 221, Motion, 221, 227.

Rotula described, 287. Its Substance, 288. Articulation; of

Children, 289.

Round Bones, their Structure, 21. Strength, how increased, ibid.

C Acrum Os, its Body, 189. Processes, oblique, transverse, 100. Spinal, 191. Foramina, 1924 Substance, Articulation, 193. Uses; of Children, 194.

Sagittal Suture, 68.

Scaphoid Bone of the Wriff, 256.

Scapula, 234. Its Base, inferior Costa, Superior, Costa, 235. Dorsum, anterior Surface, Processes, Spine, 236. Acromion, coracoid Process, 237. Third Process, Medullary Vessels, Substance, 238. Articulation, 239. Uses; of Children, 240. their contract someth

Schyndelefis, 39.

Scull, vid. Cranium.

Sefamoid Bones, how formed, 306. Where to be found, ibid. Share Bone, vid. Offa Putis.

Shoulder consists of Clavicle, 230, and Scapula, 234.

Shoulder-blade, vid. Scapula.

Sinus frontales, 80. Sphenoidal, 122. Maxillary, 131.

Skeleton, natural, artificial, 59 Situation of the Bones, 60. Divided into Head, 61. Trunk, 163. Extremities supe-

rior, 230. inferior, 272.

-- of a Female, its Differences from that of the Male, 308. Sphenoid Bone, its external Surface, Processes, 109. Cavities, 111. Internal Surface, ibid. Holes proper, 113. Common, 115. Sinufes, 116. Substance, Connexion, 117. Uses; of Children, 118.

Sphenoidal Suture, 71.

Spine, its Shape, 163. Confifts of true Vertebra, ibid. and falfe, 189.

Spongiofa inferiora Offa, 138. Their Substance, Connexion, Uses; of Children, 139.

Spongiofa

Spongiosa fuperiora Offa, 106.

Squamous Sutures, 69. How formed, 70.

Sternum, 222. Its Substance, 223. First Bone, second Bone, 224. Xiphoid Cartilage , 225. Diseases, 226. Connexion; of Children; Uses, Motion, 227.

Sutures, 38. How formed, 39.

of the Cranium, 66. Coronal, Lambdoid, ibid. Sagittal, 68. Squamous, 69. Ethmoidal, sphenoidal, trans-

verse, Zygomatic, 71. Uses, 72.

--- of the Face, distinguished by Names, 120. Uses, 121. Symplysis, divided into Synchondrofis, Synneurofis, Sysfarcofis, 38. Synarthrofis, divided into Suture, Harmonia, Gomphofis, Schindelefis, 38, 39.

Synchondrofis, 38.

Synneurofis, 38.

End,

nte-

and

hed

ad. 21.

of

fed.

rfe,

cu-

35.

ion.

els.

eu,

d.

so.

pe-

8.

es,

m-

7.

nđ

m,

fa

Synovia of joints, what composed of, 56. Its Uses, Diseases, 57. Syfarcofis, 38.

TArjus consists of seven Bones, 292. Its Articulation, Ules, 294 301.

Teeth, their Base, Roots, Periosteum, Substance, 149. Canals, 150. Vessels, 151. Formation, 152. Shedding. Connexion, Uses, 153. Distinguished into Incisores, 154. Caniai, Molares, 155. Phanomena, 158.
Temporal Bones, 88. Their external Surface, Processes,

8. Cavities, 90 Holes, 91. Internal Surface, 93. Processes, Cavities, Holes, 94. Substance, 95. Connexion. Uses; of Infants, 96.

Thigh-bone, vid. Femoris Os.

Thorax, 211. Confifts of Vertebra, 182. Ribs, 212. Ster-

num, 222. Its Motion, 227.

Thumb confifts of three Bones; the first, 267, its Articulation; at the Time of Birth, 268. Second Bone, 268. Its Articulation, 269. Third Bone, ibid.

Tibia, its upper End, 278. Cartilages, 279. Body, 280. Lower End, 281. Articulation; of Children, 282.

Toes, 304: differ from the Fingers, Uses; of Children, 305. Transverse Suture, 71.

Trapezium Os, 258.

Trapezoides Os, 259.

Triquetra Offa, 66.

Trochoides Ginglimus, 43.

Trunk of the Skeleton, consists of the Spine, 163. Pelvis, 198, 211. Thorax. ibid.

Turbinata Offa, vid. Spongiefa.

Ulna,

ULna, its upper End, 247. Body, 248. Lower End, 249. Articulation, 250. Of Children, 253.

Unciforme Os of the Wrist, 260.
Unguis Osa, their Surfaces, 123. Substance, Connexion, Uses, 124. Morbid Phænomena; of Children, 125.

TErtebra cervical, vid. Cervical. --- Dorfal, vid. Dorfal.

--- False, composed of Os sacrum, 189. Coccygis, 104.

---- Lumbar, vid. Lumbar.

--- True, 163. Their Bodies, Cartilages, 164. Processes, Ligaments, Foramina, 166. Articulations, Phanomena, 167. Of Infants, 169. Diseases, 170. Divided into Cervical, 172. Dorfal, 182. Lumbar, 186. Uses, fecured from Luxations, 188.

Vessels of Periofteum, 3. of the Bones, 8. of the Marrow, 17. 284. of Ligaments, 45. Cartilages, 48. Mucaginous

Glands, 55.

Vomer, its Figure, Situation, 140. Substance, Connexion, 141. Ules, 142...

Women, the Differences of their Bones from those of Men, 308. Wormiana Offa, vid. Triquetra. Wrist, vid. Carpus.

X

X Iphoid Cartilage, 225. Its Diseases, 226.

Z

Ygoma, 89. L Zygomatic Suture, 71:

THE

THE

ANATOMY

OFTHE

HUMAN NERVES,

WITH

An Account of the reciprocal Motions of the HEART,

AND

A Description of the HUMAN LAC-TEAL SAC and DUCT.

The SIXTH Edition.

By ALEXANDER MONRO Senior, M. D. and P. A.

249.

Ufes,

194.

Proexnovided

rrow,

inous

ofe of

H E

ADJECT FOR SOME THE APPLIES i caragina di darangan

PREFACE.

Being informed that the following Essays have been useful to the Students in Anatomy, I have caused them to be reprinted with such A-

mendments as I thought necessary.

That Offence might not be given, I have treated all the Opinions concerning the disputed Physiology of the Nerves with that Deference which the Uncertainty of the Subject required, and have not only concealed the Names of the Writers whose Sentiments were different from mine, but have shunned Quotations from those whom I approve, lest the Knowledge of the latter should be a Key to discover the former by.

Descriptions of the very minute Ramifications of the Nerves are obscure to the young Gentlemen for whose Use I write; and therefore I have taken Notice only of the larger Branches

in the Description of the particular Nerves.

I have endeavoured to illustrate my great Master, Dr. Boerhaave's Doctrine, concerning the
Systole and Diastole of the Heart, on Account
of its beautiful Simplicity; though it will appear
from what I have said in several Editions of this
little Essay on the Nerves, that I have long thought
it insufficient for the Solution of the Phænomena.
The Description of the Phænomena

The Description of the Receptaculum chyli

PREFACE.

and thoracic Duct is more accurate, than in the common Systems of Anatomy; and on that Ac.

count is here republished.

The Corrections and Additions made in this Edition of the Anatomy of the Bones, and of these Essays, shew, that my own Errors and Omissions are rather too great a Burden for me to struggle with: It is cruel to load me with a great Number more; and many of them such as the youngest Pupil under my Care would be ashamed of I must therefore beg the Public to impute momore Faults to me as an Author, than what are contained in the preceeding Anatomy of the Bones, in the following Essays, and in those Papers of the medical Essays and Observations, and Essays physical and literary, to which my Name is prefixed.

ANATOMY

OF THE

HUMAN NERVES.

Of the NERVES in general.

and vertebral Arteries make before they pass through the Dura Mater, these Arteries having neither swelling Muscles nor Pressure of the Atmosphere to affect the Course of the Blood in them after they enter the Scull, and their Division into innumerable communicating Branches in the Pia Mater and its Processes, shew, that the Liquors must move more slowly and equally in them than in most other Parts of the Body.

2. By the Affistance of Injections and Microscopes, the very minute Branches of these Vessels (§ 1.) are discovered to go from the Pia Mater, into the Cortex, cineritious, or ashy coloured Part of the Cerebrum, Cere-

E e bellum

Millet

in the at Ac-

this these issions uggle

Vumoungd of.

te no

ers of

ffays pre-

ΗE

bellum, and fpinal Marrow; whereas we can only fee longitudinal Vessels, without numerous Ramifications or reticular Plexuses, in the white

medullary Substance of these Parts.

3. The Continuity of the Cortex with the Medulla of the Encephalon and Spinal Marrow is observable with the naked Eye, and is more distinctly seen with the Assistance of a Microscope.

4. In diffecting the Brain and Cerebellum, we see the small Beginnings of the Medulla proceeding from the Cortex, and can trace its gradual Increase by the Addition of more such white Substance coming from the Cortex.

5. Both these Substances (§ 4.) are very succellent; for being exposed to the Air to dry, they lose more of their Weight than most o

ther Parts of the Body do.

6. In several Places we can observe the Medulla to be composed of Fibres laid at each others Sides.

7. The medullary Substance is all employed in forming the white sibrous Cords, which have now the Name of Nerves appropriated to them. Within the Scull, we see the Nerves to be the medullary Substance continued; and the spinal Marrow is all employed in forming Nerves.

8. The common Opinion concerning the Rife of the Nerves, founded on a superficial Inspection of those Parts, is, that the Nerves are propagated from that Side of the Eucephalon, at which they go out of the Scull. But it having been remarked, after a more strict Inquiry, and preparing the Parts by Maceration

111

can

OUS

hite

the

וניסיו

ore

ro-

ım,

illa

its

ich

uc-

ry,

0-

he

ich

Dy.

ch

ed

res

nd

ng

he

ial

es

10-

it

n-

on

in

(1900) (1900) (1900) (1900) (1900)

in Water, that the medullary Fibres decussate or cross each other in some Parts of the Medulla; as for Example, at the Corpus annulare, and Beginning of the Spinal Marrow: And practical Objervators having related feveral Examples of People whose Brain was hurt on one Side, while the morbid Symptom, Palfy, appeared on the other Side of the Eody. of which I have feen fome Inflances; and Experiments made on Brutes having confirmed these Observations, it has been thought, that the Nerves had their Rife from that Side of the Encephalon, which is opposite to their Egress from the Scull. It may however still be faid, that this last Opinion is not fully demonstrated, because a Decussation in some Parts is not a Proof that it obtains univerfally; and if there are Examples of Palfy of the Side opposite to where the Lesion of the Brain was, there are also others, where the Injury done to the Brain and the Palfy were both on the fame Side.

9. The Nerves are composed of a great many Threads lying parallel to each other, or nearly so, at their Exit from the Medulla.

This fibrous Texture is evident at the Origin of most of the Nerves within the Scull; and in the Cauda equina of the spinal Marrow, we can divide them into such small Threads, that a very good Eye can scarce perceive them; but these Threads, when looked at with a Microscope, appear each to be composed of a great Number of smaller Threads.

Nerves is, we know not: but when we consider

E e 2 that

that every, even the most minute Part of the Body is sensible, and that this must depend on the Nerves (which all conjoined, would not make a Cord of an Inch Diameter) being divided into Branches or Filaments to be dispersed through all these minute Parts, we must be convinced that the nervous Fibrils are very small. From the Examination of the minimum visibile, it is demonstrated, that each Fibre in the Retina of the Eye or expanded optick Nerve, cannot exceed the Size of the 32,400 Part of a Hair.

nervous Fibrils are composed, is very tender, and would not be able to resist such Forces as the Nerves are exposed to within the Bones, nor even the common Force of the circulating Fluids, were not the Pia Mater and Tunica Arachnoides continued upon them; the former giving them Firmness and Strength, and the latter surnishing a cellular Coat to connect the Threads of the Nerves, to let them ly soft and moist, and to support the Vessels which go with them.

It is this cellular Substance that is distended when Air is forced through a Blow-pipe, thrust into a Nerve, and that makes a Nerve appear all spongy, after being distended with Air till it dries; the proper nervous Fibrils shrivelling so in drying, that they scarce can be observed.

12. These Coats (§ 11.) would not make the Nerves strong enough to bear the Stretching and Pressure they are exposed to in their Course to the different Parts of the Body; and therefore

fore,

fore, where the Nerves go out at the Holes in the Cranium and Spine, the Dura Mater is generally wrapt closely round them, to collect their different Fibres into tight firm Cords; and that the Tension which they may happen to be exposed to, may not injure them before they have got this additional Coat, it is firmly fixed to the Sides of the Holes in the Bones through which they pass.

13. The nervous Cords thus composed of nervous Fibrils, cellular Coat, Pia and Dura Mater, have such numerous Blood-vessels, that after their Arteries only are injected, the whole Cord is tinged of the Colour of the injected Liquor; and if the Injection is pushed violently, the cellular Substance of the Nerves is at last

distended with it.

14. A nervous Cord, fuch as has been just now described, (§ 13.), has very little Elasticity, compared with several other Parts of the Body. When cut out of the Body, it does not become observably shorter, while the Blood-vessels contract three eighths of their

Length.

the

d on

not

z di-

per-

t be

very

nmi-

ibre

tick

400

the

der,

as

nes,

ing

nca

ner

the

the

and

ith

led

pe,

rve

ith

rils

be

he

ng

rie

re-

e,

Access Sections Secti

lar or fatty Substance, and have their Course in the Interstices of Muscles, and other active Organs, where they are guarded from Pressure; but in several Parts they are so placed, as if it was intended that they should there suffer the vibrating Force of Arteries, or the Pressure of the contracting Fibres of Muscles.

16. The larger Cords of the Nerves divide into Branches as they go off to the different Parts; the Branches being smaller than the

Ee 3

Trunk from which they come, and making generally an acute Angle where they separate.

17. In several Places, different Nerves unite into one Cord, which is commonly larger than

any of the Nerves which form it.

18. Several Nerves, particularly those which are distributed to the Bowels, after such Union, (§ 17.), suddenly form a hard Knot considerably larger than all the Nerves of which it is made. These Knots were called Corpora olivaria, and are now generally named Ganglious.

numerous and larger Blood-vessels, than the Nerves; so that they appear more red and muscular. On dissecting the Ganglious, Fibres are seen running longitudinally in their Axes, and other Fibres are derived from their Sides in an oblique Direction to the longitudinal ones.

which conjunctly are not equal to the Size of the Ganglian, are fent out from it, but with a Structure no Way different from that of other

Nerves.

21. The Nerves sent to the Organs of the Senses, lose there their firm Coats, and terminate in a pulpy Substance. The optick Nerves are expanded into the soft tender Webs, the Retina. The auditory Nerve has scarce the Consistence of Mucus in the Vestibulum, Cochlea and semicircular Canals of each Ear. The Papillae of the Nose, Tongue and Skin, are very soft.

22. The Nerves of Muscles can likewise be traced till they seem to lose their Coats by becoming very soft; from which, and what we ob-

ferved

ferved of the fenfatory Nerves, (6 21.), there is Reason to conclude, that the muscular Nerves are also pulpy at their Terminations, which we cannot indeed profecute by Diffection.

ge.

nite

an

ich

on,

er-

is

va-

H'e

he

nd

es

es,

les

al

es,

of

2

er

10

te

X-

₽.

e

iof

e

.

22. It would feem necessary that the Extremities of the Nerves should continue in this foft flexible State, (§ 21. 22.), in order to perform their Functions right: For, in Proportion as Parts become rigid and firm by Age, or any other Cause, they lose of their Sensibility, and the Motions are more difficultly performed.

24. Though the Fibres in a nervous Cord are firmly connected, and frequently different Nerves join into one Trunk, or into the fame Ganglion; yet the Senfation of each Part of the Body is fo very diffinct, and we have fo much the Power of moving the Muscles separately, that, if the Nerves are principal Agents in these two Functions, which I shall endeavour to prove they are, we have Reason to believe that there is no Union, Confusion, or immediate Communication of the proper nervous Fibrils, but that each Fibre remains distinct from its Origin to its Termination.

25. Changes produced any Way upon the Coats of the Nerves, cannot however miss to The cellular Subaffect the nervous Fibrils. stance may be too full of Liquor, or may not supply enough; the Liquor may not be of a due Confistence, or it may be preternaturally obstructed and collected. The Pia or dura Muter may be too tense, or too lax; their Vessels may be obstructed; their proper Nerves may be violently irritated, or lose their Power of acting; and a great many other such Changes

may happen, which will not only occasion Diforders in particular Nerves, but may cause the Sympathy so frequently observed among the Nerves; which is so necessary to be attentively regarded in a great many Diseases, in order to discover their true State and Nature, that, without this Knowledge, very dangerous Mistakes in the Practice of Physick and Surgery

may be committed.

26. Many Experiments and Observations concur in proving, that when Nerves are compressed, cut, or any other way destroyed, the Parts served by such Nerves, farther from the Head or Spine than where the injuring Caufe has been applied, have their Senfations, Motions, and Nourishment, weakened or lost, while no fuch Effects are feen in the Parts nearer to the Origin of those Nerves; and in fuch Experiments where the Cause impeding the Nerves to exert themselves could be removed, and the Structure of the Nerves not injured; as for Example, when a Ligature made upon a Nerve and stopping its Influence has been taken away, the Motion and Sensation of the Parts foon were restored. From which it would appear, that the Nerves are principal Instruments in our Sensations, Motions, and Nourishment; and that this Influence of the Nerves is not inherent in them, without the Communication between these Cords and their Origin is preferved.

This Conclusion is just, notwithstanding that fometimes, upon cutting a Nerve, the Esseds above mentioned have been felt for a short Time; but afterwards the Person was sensible

10

n Dif.

fe the

the the

itive.

order

that,

Mif.

gery

tions

com.

the

the

aule

Mo-

oft,

arts

l in

ing.

10-

1n-

ade

has

of

It

n-

nd

he

ne

ir

t

e

The state of the s

of no Numbness or Immobility; for where-ever this is said to have happened, the cut Nerve was only one of several which were sent to the Member; the want of whose Influence was felt no longer, than till the Habit was acquired of performing the Functions easily by the other Nerves.

Nor is it of greater Weight as an Objection, that sometimes when a Ligature is drawn very hard upon a Nerve, and then is taken away, the Nerve never again recovers its Insluence upon the Parts it is distributed to beyond the Ligature, but is of as little Effect as if it had been cut through; which is to say, that its Texture has been altered beyond Recovery. The same Thing is to be seen by tying a Thread tight round a tender Twig of any Vegetable; it decays.

27. Experiments and Observations shew too, that when Parts of the Encephalon or Spinat Marrow have been irritated, compressed, or destroyed, the Parts of the Body, whose Nerves had their Origin from such affected Parts of the Eucephalon or spinal Marrow, became convulsed, paralytic, infensible or wasted; and in fuch Cases where the injuring Cause could be removed from the Origin of the Nerves, the morbid Symptoms observed in the Parts to which these Nerves were distributed, went off upon the Removal of that Cause. From which it is thought reasonable to conclude, that the Nerves must not only have a Communication with their Origin, but that the Influence they have upon the Parts they are distributed to, de-

pends

pends on the Influence which they derive from

the Medulla Encephali and Spinalis.

28. Though the fpinal Marrow has its own Vessels and cineritious Substance which assists to form its Medulla; yet a very large Share of the medullary Substance within the Spine is derived from the Encephalon, whose Medulla oblongata descends from the Head, and the Influence of the spinal Marrow on its Nerves depends in a great Measure on this Medulla oblongata of the Head. Hence an Injury done to any Part of the spinal Marrow, immediately affects all the Parts whose Nerves have their Origin below where the injuring Caufe is applied. tion of a Vertebra in the Loins makes the lower Extremities foon paralytic; a transverse Section of the Medulla at the first Vertebra of the Neck, foon puts an End to Life.

29. If such Causes produce constantly such Effects (6 26. 27. 28.) in us and other Creatures living in nearly the same Circumstances as we do, the Conclusions already made must be good, notwithstanding Examples of Children and other Creatures being born without Brains or spinal Marrow; or notwithstanding that the Brains of adult Creatures can be much changed in their Texture by Diseases, that Tortoifes, and some other Animals, continue to move a considerable Time after their Heads are cut off. We may be ignorant of the particular Circumstances requisite or necesfary to the Being or Well-being of this or that particular Creature, and we may be unable to account for a great many Phanomena; but we must believe our Eyes in the Examination

of Facts; and if we see constantly such Consequences from fuch Actions, we cannot but conclude the one to be the Caufe, and the other the Effect. It would be as unjust to deny the Conclusions made in the three preceeding Articles, because of the seemingly preternatural Phænomena mentioned in the Beginning of this Article, as it would be to deny the Necessity of the Circulation of the Blood in us and most Ouadrupeds, because a Frog can jump about, or a Tortoise can walk long after all the Bowels of its Thorax and Abdomen are taken out, or because the different Parts of a Worm crawl after it has been cut into a great many Pieces. It is therefore almost universally allowed that the Nerves are principal Instruments in our Senfations, Motion and Nourishment; and that the Influence which they have is communicated from their Origin, the Encephalon and Medulla spinalis. But Authors are far from agreeing about the Manner in which this Influence is communicated, or in what Way Nerves act to produce these Effects.

30. Some alledge, that the nervous Fibres are all solid Cords acting by Elasticity or Vibration. Others maintain, that those Fibres are small Pipes conveying Liquors, by means of which their Ef-

fects are produced.

e from

S OWn

ifts to

of the

crived

mgata

ce of

ina

f the

rt of

1 the

eiow

UXa-

OWer

ecti-

the

fuch

rea-

nces

nust

Iren

ains

that

uch

and

nti-

eir

of

cef-

hat

ble

out

011

of

10000 Page 10000 Page

31. The Gentlemen, who think the nervous Fibres solid, raise several Objections to the other Doctrine; which I shall consider afterwards; and endeavour to shew the Fitness of their own Doctrine to account for the Effects commonly observed to be produced by the Nerves.

The

The Objects of the Senses plainly (fay they) make Impulses on the Nerves of the proper Organs, which must shake the nervous Fibrils; and this Vibration must be propagated along the whole Cord to its other Extremity or Origin, as happens in other tense Strings; and these Vibrations being differently modified, according to the Difference of the Object, and its different Application, produce the different

Ideas we have of Objects.

32. To this Account of the Sensations, it is objected, 1st, That Nerves are unfit for Vibrations, because their Extremities, where Objects are applied to them, are quite soft and pappy (§ 21.), and therefore not susceptible of the Vibrations supposed; and if there could be any little Tremor made here by the Impulse of Objects, it could not be continued along the nervous Cord, because the cellular Substance by which each particular Fibre is connected to the neighbouring ones (§ 11.), and the fatty Substance in which the nervous Cord is immersed (§ 15.), would soon stifle any such vibratory Motion.

A 2d Objection to this Doctrine is, that, supposing the Nerves capable of Vibrations by the Impressions of Objects, these Vibrations would not answer the Design. For if what we know of other vibrating Strings, to wit, that their Tone remains the same, unless their Texture, Length or Tension is altered, and that different Substances striking them, do no more than make the Sound higher or lower; if these Properties are to be applied to Nerves, then it will follow, that the same Nerve would constantly convey the same Idea, with no other Variety

than

than of its being weaker or stronger, whatever different Objects were applied to it; unless we supposed the Nerve changed in its Texture, Length or Tenfion, each Time a different Objed is applied; which, it is prefumed, no body will undertake to prove does happen.

Nay, 3dly, If ever fuch a Variety of Vibrations could be made, our Senfations would notwithstanding be confused and indistinct, because the tremulous nervous Fibre being firmly connected and contiguous to feveral other Fibres of the fame Cord, would necessarily shake them too, by which we should have the Notion of the Object as applied at all the different Parts where the Extremities of these Fibres terminate.

33. In whatever Way the Favourers of the Doctrine of folid Nerves please to apply the Elasticity of Nerves to the Contraction of Mus cles; their Adversaries infift that Nerves are too weak to refift fuch Weights as the Mufcles fuftain; they would furely break, especially as they are in a great Measure, if not wholly, deprived of their strong Coats before they come to the Part of the Muscle they are immediately to accupon (§ 22.).—The Nerves being found to have little or no Elasticity to shorten themselves (§ 13.), thews them altogether unfit for fuch an Office as this of contracting Muscles in the Way proposed of their acting by Elasticity; and when a Nerve is viewed with a Microscope while the Muscles it serves are in Action, no Contraction or Motion is observed in it.

34. As a further Objection against either Motion or Sensation being owing to the Elasticity of the Nerves, it is faid, that if this Doctrine

Ff

hey)

oper

rils:

ong

Ori-

and

ac-

and

rent

it is

bra-

ects

ppy

the

anv Ob-

ner-

nich gh.

e in 5.),

upi

the

uld WO

eir

ire,

ent

nan

ro-

vill

tly

ety

lan

was true, the Sensations would be more acute, and the Contractions of Muscles would be greater and stronger, when the Parts become firmer and more rigid by Age; for then their Elasticity is increased: Whereas, on the contrary, it appears (§ 23.), that then the Sensations are blunted, and muscular Contraction be-

comes less and weaker.

35. If the Nerves were granted to be elastic, and to communicate a springy Force to all the Parts they are distributed to, they might appear necessary in this View to assist the Application of the nutritious Particles of the Fluids to the Sides of the Vessels which these Particles were to repair; and so far might well enough account for the Share which Nerves are thought to have in Nutrition: But if we cannot make use of Elasticity in the other two Functions, Sensation and Motion, we must also endeavour to find out some other Way for the Nerves to act in Nutrition; which will be done afterwards.

36. Having thus stated the Reasons for and against the Nerves acting as solid Strings, let us likewise relate the Arguments for Nerves being Pipes, and the Objections to this

Doctrine.

A great Argument of those who think the Nerves to be Tubes conveying Liquors, is the strong Analogy of the Brain and Nerves to other Glands of the Body and their Excretories, where a manifest Secretion of Liquor is made in the Glands, to be conveyed by the Excretories to the proper Places in which it ought to be deposited: They think that the vascular Tex-

ture

cute,

d be

come

their

con.

ısatiı be-

Iftic,

the

ap-

lica-

ds to

icles

ugh

ight

iake

ons,

our

s to

ter-

and

let

rves

this

the

the

0-

ies,

e 111

ries

be

ex-

ure

ture of the Cortex of the Encephalon and spinal Marrow (§ 2.), the Continuation of the Cortex in forming the medullary Substance (§ 3. 4.), the sibrous Texture (§ 6.) and succulent State of this Medulla (§ 5.), and its being wholly employed to form the Nerves (§ 7.), where the sibrous Texture is evident (§ 9.); all these Things, say they, conspire to shew such a strong Analogy between these Parts and the other Glands of the Body, as carries a Conviction that there is a Liquor secreted in the Encephalon and spinal Marrow, to be sent out by the Nerves to the different Parts of the Body.

37. The following Objections are raised to this Argument in favour of Liquor conveyed in the Nerves, from the Analogy of the Glands. ift, Other Glands, it is faid, have their Excretories collected into a few large Pipes, and not continued in fuch a great Number of separate Pipes, as far as the Places where the Liquors are deposited; which last must be the Case, if the Nerves are the Excretories of the glandular Brain. 2dly, We fee the Cavities, and can examine the Liquois in the Excretories of other Glands much smaller than the Brain; which cannot be done in the Nerves. 3dly, If the Nerves were Pipes, they would be fo small, that the Attraction of the Liquors to their Sides, would prevent that Celerity in the Motion of the Liquors, which is requifite to Senfations and Motions. 4thly, If the Nerves were Pipes, they would be cylindrical ones, and confequently not subject to Diseases; or at least we could have no Comprehension of the Diseases in them.

Ff 2

38. The

38. The Answer to the 1st of these Objecti. ons is, That there are other Glands where there is a manifest Secretion, and in which the Difposition of the Excretories is in much the same Way as in the Encephalon: The Kidneys, for Example, have a reticulated Cortex of Veffels. from which the Eustachian or Bellinian Medulla. confifting of longitudinal Fibres and a few Blood-vessels in the same Direction, proceeds: and this Medulla is collected into ten, twelve or more Papille, each of which is formed of nemerous imall feparate Pipes, which fingly difcharge the Urine into the large membranous Tubes; and these united form the Pelvis. Upon comparing this Texture of the Kidneys with that of the Encephalon (§ 2. 3. 4. 5. 6. 7. 9.), the Analogy will be found very strong.

39. In Answer to the 2d Objection, in § 37. it is granted, that Microscopes, Injections, and all the other Arts hitherto employed, have not thewn the Cavities of the nervous Fibrils, or the Liquors contained in them; and from what was faid (§ 10.) of the Smallness of the neryous Fibrils, it is not to be expected that ever they should be seen. But so long as such a Number of little Animals can every Hour be brought to the Objectors, in which they can as little demonstrate the Vessels or contained Fluids, it will not be allowed to be conclusive Reasoning, that because ocular Demonstration cannot be given of either the Tubes or their Contents, therefore they do not exist. For if we have any Notion of an Animal, it is its being an Hydraulick Machine, which has Liquors moving in it as long as it has Life; if therefore fuch

fuch little Animals have Vessels and Liquors which we cannot see, why may not some of the Vessels and Liquors of the human Body be also invisible to us.

ſ.

16

r

s,

4,

14

n

To avoid this Answer to the Objection, it is further urged, That though we might not fee the nervous Tubes or the Liquors they contain, as they naturally flow; yet if fuch Liquors really exist, they ought to discover themselves, either by a Nerve's swelling when it is firmly tied; or that, however subtile their Fluids are, they might be collected in some Drops at least, when the cut End of a Nerve of a living Animal is kept some Time in the exhausted Receiver of an Air-pump. It is affirmed, that neither did the tied Nerve swell between the Brain and Ligature, nor was there any Liquor collected in the Receiver of the Air-pump; from which it is concluded, that there is no Liquor in the Nerves.

Some, who fay they have tried these Experiments, affirm, that in young Animals the Nerve does swell above the Ligature, and that a Liquor does drill out upon cutting a Nerve.—Whether Swelling or Liquor is seen or is not seen in these Experiments, no Conclusion for or against a nervous Fluid can be made from them; for the Swelling of the Nerve after it is tied, or the Essure of Liquors from its Extremity, will never prove either to be the Essect of the Fluid in the proper nervous Fibrils, so long as they might be occasioned by the Liquors in the larger Vessels of the cellular Substance of the Nerves; and if these same Vessels of the Coats of the Nerves do

Ff3

not discover their Liquors by these Experiments, it is far less to be expected, that the much more subtile Nerves will discover their.

40. The 3d Objection to the Doctrine of the Brain being a Gland, and the Nerves its Excretories, supposes a more rapid Motion necessary in the Fluid of the Nerves, than what most of the Defenders of the nervous Fluid will now allow; and is afterwards to be considered particularly in a more proper Place.

41. The 4th Objection being, That if Nerves are Excretories of a Gland, they must be cylindrical Pipes, in which no Obstructions or Difeases would happen; but fince we daily fee Diseases in the Nerves, they must therefore not be such Excretories. The Answer is, That Difeases happen often in the Excretories of other Glands, as of the Liver, Kidneys, &c. notwithstanding their cylindrical Form, and their much shorter and less exposed Courfe. When we consider the very tender Substance of the Brain, the vast Complication of Vessels there, the prodigious Smallness of the Pipes going out from it, the many moving Powers which the Nerves are to undergo the Shock of, and the many Chances which the Vessels, Membranes and cellular Substance accompanying the Nerves have of being difordered, and then affecting the nervous Fibrils, we have very great Reason to be furprifed, that these cylindrical Pipes are not much more frequently put out of Order, by too great or too fmall a Quantity of Liquors; by too viscid or too thin Fluids; by Liquors confifting of too mild and fluggish Particles,

ne

of

ts

n

n

d 1-

1-

1

r

-

1

1-

n

o h

e

t

or of too acrid pungent ones; by too great or too little Motion given to the Liquors; by the Diameters of the Pipes being too much straitened, or too much inlarged; and by a great many other Varieties of Circumstances which might be thought capable of disturbing the Functions of the Nerves, supposing them to be cylindrical Excretories of the Gland, the Brain.

42. The numerous Vessels of the Encephalon have brought some of the Gentlemen who assert the Nerves to be solid, to acknowledge, that there is a Liquor secreted in the Brain: But then they will not allow that this Liquor is sent out by the proper nervous Fibrils; but that it is poured into the cellular Substance in which the Nerves lie, to keep them moist and supple, and therefore sit for exerting their Elasticity, Vibration, &c. by which, in their Opinion, the Essets commonly ascribed to Nerves are produced.

43. Besides the Objections already mentioned (§ 32. 33.) against the Nerves acting as elastic Strings, this Opinion has some other Dissiculties, which may be objected to it: For Instance, there is not one analogous Example in the whole Body of Liquors secreted in a large Gland, to be poured into a cellular Substance, as is here supposed; the Liquors in the Cells of the tela cellularis of other Parts are separated from the little Arteries which are

distributed to these Cells.

Further, it cannot be imagined, how a Liquor secreted in the Cortex of the Brain, should make its Way through the Medulla, to come

Out

out into the cellular Membranes on the Surface of that Medulla.

Lastly, A very simple Experiment, of injecting Water by the Artery of any Member, and thereby filling the cellular Substance of the Nerves of that Member, shews evidently, that the Liquor of the cellular Substance of the Nerves has the same Fountain as the Liquor has in the tela cellularis any where else, that is, from the little Arteries dispersed upon it.

44. The Doctrine of a Fluid in the Nerves, is not only thus supported by the Analogy of the Brain and Nerves to the other Glands and their Excretories, but those who maintain this Doctrine mention an Experiment which they think directly proves a Fluid in the Nerves. It is this: After opening the Thorax of a living Dog, catch hold of and press one or both the phrenic Nerves with the Fingers, the Diaphragm immediately ceases to contract; cease to compress the Nerves, and the Muscle acts again: A fecond Time, lay hold of the Nerve or Nerves some Way above the Diaphragm, its Motion stops. Keep firm the Hold of the Nerve, and, with the Fingers of the other Hand, strip it down from the Fingers which make the Compression, towards the Diaphragm, and it again contracts: A Repetition of this Part of the Experiment three or four Times, is always attended with the fame Effects; but it then contracts no more, strip as you will, unless you remove the Pressure, to take hold of the Nerves above the Place first pinched; when the Muscle may again be made

1-

t-

e

,

n

d

to contract, by stripping the Nerve down towards it. This Experiment I have done with the Success here mentioned. Let any one try if he can imagine any other reasonable Account of these Appearances, than that the Pressure by the Fingers stopped the Course of a Fluid in the Nerve; that fo much of this Fluid as remained in the Nerve, betwixt the Fingers and Diaphragm, was forced into that Muscle by stripping, and when it was all preffed away, the Fingers above preventing a Supply, the Muscle contracted no more till the Fingers were removed, and a fresh Flow by that Means was received from the spinal Marrow, or from that Part of the Nerve which had yet not been so stripped.

It has been objected to the Conclusions from this Experiment, 1. That the Diaphragm is set in Motion by stripping the Nerve from, as well as towards, this Muscle, and this may be well expected; for a Liquor in such small Pipes hindered to slow backwards by Ligature, pinching Fingers, or even the Flow of their Liquors from the Fountain, will regurgitate forwards with Velocity, when pressed backwards. We see it happen in the Stalks of tender succulent Plants.

2. It is said, that Muscles cease to act when their Veins are tied, as well as when their Arteries or Nerves are tied or cut, but that Muscles continue to act when their Veins are cut; by which it would appear, that the overloading of the Vessels is an Impediment to the Action of Muscles, and therefore the ceasing of their Action when their Arteries or Nerves are tied or cut, may also be owing to the Liquor

quor in the Branches of these Pipes of Muscles stagnating when it is not propelled by the Flow of more Liquor from their Trunks, and not to any Influence or moving Power, which now

ceases to be conveyed to them.

It is to be observed in making the Experiments just now mentioned, that the Contracti. on of the Muscles ceases soonest when the Nerves, and latest when the Veins are tied. That when Veins are tied, not only are the Vessels overloaded, but all the cellular Sub. stance of the Muscles is filled with coagulated Blood; whereas when the Arteries or Nerves are tied, the reverse is seen, the Muscles are lax and of less Bulk. So that in these Cases the Ceasing of the Contraction of the Muscles feems to depend on very different Causes, to wit, a Deprivation of necessary Liquors in the one, and a Redundancy of superfluous Blood in the other. An elastic Stick may be deprived of its Elasticity by being made either too dry or too wet.

45. Some Gentlemen, convinced of the Reasonableness of the Secretion of a Liquor in the Brain to be sent out by the Nerves, but not comprehending how a Fluid could have such a rapid retrograde Motion as they imagined was necessary for conveying the Impressions of Objects made on the Extremities of Nerves to the Sensorium, supposed two Sorts of Nerves; one that conveyed a Liquor for muscular Motion and Nutrition; the other composed of solid Nerves, that were to serve for Organs of the Senses, to convey the Vibrations communicated from Objects to the Sensorium.

46. Ta

11

is

fa

C

121

te

P

V

ta

ir

Sp

n

46. To this Opinion (§ 45.) the Objections against the sensatory Nerves acting by Vibration (§ 32.) may be made; and there is so little Reason to suspect any Difference in the Texture of the different Parts of the Brain or Nerves, that, on the contrary, the Structure is every where similar, and Branches of the same Nerve often serve both for Sensation and Motion.

How little Necessity there is for supposing extremely rapid Motions of the nervous Fluid,

is to be examined foon.

47. The Hypothesis of great Celerity in the Motion of the Fluid of the Nerves being neceffary, gave also Rise to another Division of the Nerves, into arterious or effluent, and venow or refluent. It was faid, that muscular Motion and Nutrition depended on the arterious Nerves; and that the Senfations depended on an accelerated Motion of the nervous Fluid towards the Brain, by the Impreffions which the Objects of the Senses make upon the venous Nerves. By this Supposition the Absurdity of rapid Fluxes and Ressuxes in the fame Canal was prevented, and an Advantage was thought to be gained by it, of faving too great a Waste of the Fluid of the Nerves, which otherwise the Encephalon and spinal Marrow could not supply in sufficient Quantity to answer all the Exigencies of Life.

48. To this Opinion (§ 47.) it has been objected, 1st, That there is no Example in the Body of a secreted Liquor being returned immediately and unmixed to the Gland by which

it was originally separated from the Mass of Blood; which would be the Case were there venous Nerves. 2th, There is no Occasion for faving the Fluid of the Nerves in the Way proposed; the Organs for secreting that Fluid being large enough to supply all that is necessary of it in the common Functions of Life. 3 dly, If the Fluid of the Nerves was to be thuskept in a perpetual Circulation, it would foon become too acrid for continuing with Safety in such sensible tender Vessels as the Brain and Nerves are composed of. 4thly, This Hypothelis will not answer the Delign for which it was proposed: For though the momentary Application of an Object might canse an Acceleration in the Fluid of venous Nerves, yet if the Object was kept applied to the Nerves, it would stop their Fluid, so that it could not go forward to the Brain; and therefore, according to this Doctrine, we should be sensible of no Objects, except those whose Application to the Organs of the Senses was momentary.

Encephalon and fpinal Marrow secen a Liquor from the Blood which is sent into all the Nerves, and that by the Means of this Liquor the Nerves perform the Offices commonly assigned to them; it is next necessary to inquire what Kind of Liquor this is, and how it moves, in order to determine how well its Nature and Motion are sitted for performing what is expected from

t. I de Militar

50. The Liquor of the Nerves has been fancied by some to be of a very strong acid or

or alcaline Nature: But fince none of our Juices appear to be of this Sort, and fince such Liquors irritate and destroy the Parts of the Body which they are applied to, we cannot conceive how the Brain can separate, or the Nerves could bear any thing of such an acrid Nature. This Tenderness and Sensibility of these Organs must hinder us absolutely from supposing that the Liquor of the Nerves can be acrid or pungent, or of the Nature of Spi-

rit of Wine, Hartshorn, &c.

d

C

n

n

ľ

Nerves to be capable of vast Explosion like Ganpowder, or of violent sudden Rarefaction like Air, or of strong Ebullition like boiling Water, or the Mixture of Acids with alcaline Liquors. But as the Mass of Blood from which this Fluid is derived, is not possessed of any such Properties, we cannot suppose the Blood to surnish what it has not in itself. Besides, all these Operations are too violent for the Brain or Nerves to bear; and when once they are begun, they are not so quickly controuled or restrained, as Experience teaches us the Nerves can be made to cease from acting.

52. We are not sufficiently acquainted with the Properties of an Æther or of electrical Effuvia pervading every Thing, to apply them justly in the Animal Occonomy; and it is as difficult to conceive, how they should be retained or conducted in a long nervous Cord. These are Difficulties not to be surmounted.

53. The furest Way of judging what Kind of Liquor this of the Nerves must be, is to G g examine

examine the Liquors of fimilar Parts of the Body. All the Glands feparate Liquors from the Blood much thinner than the compound Mass itself; such is the Liquor poured into the Cavity of the Abdomen, Thorax, Ventricles of the Brain, the Saliva, pancreatic Juice, Lymph, &c. Where-ever there is Occasion for se. creted Liquors being thick and viscid, in order to answer better the Uses they are intended for, Nature has provided Refervoirs for them to stagnate in, where their thinner Parts may be carried off by the numerous absorbent Veins dispersed on the Sides of those Cavities; or they may exhale where they are exposed to the open Air. The Mucus of the Nose becomes viscid by Stagnation; for, when it is immediately fecreted, it is thin and watry; as appears from the Application of Sternutatories, &c. The Cerumen of the Ears is of a watry Confistence, when just squeezed out. The Mucus of the alimentary Canal grows thick in the Lacuna. The Bile in the hepatic Duct has little more Confistence than Lymph; that in the Gall-bladder is viscid and firong. The Urine is much more watry as it flows from the Kidneys, than when it is excreted from the Bladder. The Seed is thin as it comes from the Testicles, and is concocted in the Vesiculæ seminales, &c.

54. Hence (§ 53.) we may fafely conclude, that a thin Liquor is fecreted in the Cortex Encephali and fpinal Marrow; and feeing the Thinness of secreted Liquors is generally, as the Divisions of the Vessels, into small subtile

Branches

Branches, and that the Ramifications within the Scull are almost infinitely subtile, the Liquor secreted in the *Encephalon* may be determined to be among the finest or thinnest Fluids.

75. Seeing also that we can observe no large Reservoir, where the Liquor secenced in the cortical Substance is deposited, to have its siner Parts taken off, we have Reason to think, that it goes forward into the Nerves in the

fame Condition in which it is fecerned.

for the or fubtile animal Liquors, is meant no more than those which are very fluid, and which seem to consist of a large Proportion of watry Particles, and a lesser one of the oily, saline and terrestrious Particles. Some of the Liquors which we can have in sufficient Quantity to make Experiments with, are so sluid, and have so little Viscidity or Cohesion of Parts, that when laid upon a Piece of clean Mirror, they evaporate without leaving a Stain; such is the Liquor outing out from the Surface of the Pleura, the Lymph, and several others.

If then these Liquors, which are subject to our Examination, the secenting Vessels of which are so large that we can see them, have such a small Cohesion of Parts, it might not be unreasonable to say, that the Liquor of the Nerves is as much more fine and sluid than Lymph, as the Vessels separating it are smaller; and therefore that the Fluid of the Nerves is a desecated Water, with a very small Proportion of the other Principles extremely sub-

tilized.

10

of

r

t

d

Gg 2

57. Two

- dict this Opinion of the Liquor of the Nerves being fo fluid and subtile. One is, that upon cutting the Cauda equina of a living Animal, a Liquor as viscid as the White of an Egg drops out: The other is, that a wounded Nerve yields a glairy Sanies. But these don't appear to be the proper Fluid of the Nerves; since it is evident, that what is discharged in both these Cases, comes out of the cellular Substance involving the nervous Fibrils.
- 58. Confidering how many Experiments make it evident, that there is a constant uninterrupted Stream of Liquors flowing through all the Canals of Animals, which convey Liquors composed of Particles smaller than the Diameter of their Canal, which is always the Case of the Nerves in a natural State; it is furprifing how it ever could be thought that the Liquid of the Nerves should be obliged to flow from the Brain to each Muscle the Moment we will; or that this Liquor should flow back with the like Swiftness from the Extremity of each Nerve, to which an Object of Sensation is applied. The Nerves, as well as the other Excretories of the Glands, always are full of Liquor; the Degree of Distension of the Canals not being at all Times alike even in a found State. But this happens without Inconvenience, as the Sides of the Canals have a Power to accommodate themselves to the present Quantity, unless it is very much above

or below the natural Standard; in both which

Cases, Diseases ensue.

59. The Motion of the Fluid in the Nervesis therefore not only constant, but it is also equal, or nearly fo: For, though the Blood in the larger Arteries is moved unequally by the unequal Forces, the Contraction of the Ventricle of the Heart, and the weaker Power, the Systole of the Arteries; yet the Difference between these two moving Powers comes to be less and less perceptible, as the Arteries divide into smaller Branches; because of the numerous Resistances which the Liquors meet with, and because the Canals they move in become larger, till in the very imaliarterious Branches, there is no sensible Difference in the Velocity of the Liquors from the Effect of the Heart or Arteries. The Motion of the Fluids must still be more equal in the Excretories of Glands, and particularly in those where the Vesfels have divided into very minute Branches, and the Liquors have no other propelling Force but the Heart and Arteries, (see § 1.), therefore the nervous Fluid moves constantly, equally, and flowly, unless when its Course is altered by the Influence of the Mind, or by the Pressure of some neighbouring active Organ.

ty of the Valves supposed by some in Nerves, we are not to assume them in accounting for

any Phænomena.

61. We have not, and perhaps cannot have any Idea of the Manner in which Mind and Body act upon each other; but if we allow that the one is

Gg3

affected

affected by the other, which no Body denies, and that the Fluid of the Nerves (whatever Name People please to give it) is a Principal Instrument which the Mind makes Use of to influence the Actions of the Body, or to inform itself of the Impressions made on the Body, we must allow that the Mind can direct this Instrument differently, particularly as to Quantity and Celerity, though we must remain ignorant of the Manner how many Phænomena depending on this Connexion of Mind and Body are produced. Thus we would in vain attempt to account for Animals continuing after their Heads were struck off or their Hearts were cut out, to perform Actions begun before they suffered any Injury.

fuch as has been argued for, to wit, a very fluid saponaceous Water, moving in a constant, equal, slow Stream, from the Eucephalon and spinal Marrow, in each of the proper nervous Fibres, except when the Motion is changed by some accessory Cause, such as the Mind, Pressure of other Parts, &c.; and let us examine how well such a Supposition will agree with the Phænomena of the three great Functions, Nutrition, Sensation, and muscular Motion, which

the Nerves are principal Instruments of.

63. In general, we may fay, that Nerves can carry Fluids to the most minute Part of the Body, to supply what is wasted in any of the Solids; that the Impression made by the Objects of the Senses on the very soft pulpy Extremities of the Nerves of the Organs of the

Senses,

Senses, must make such a Stop in the equalflowing nervous Fluid, as must instantaneously be perceptible at the Fountain-head from which the Pipes affected arise; that the constant Flow of the Liquor of the Nerves into the Cavities of the muscular Fibrillæ, occasions the natural Contraction of the Muscles, by the as constant misus it makes to increase the transverse, and to shorten the longitudinal Diameter of each Fibre; and that it is only to allow the Mind a Power of determining a greater Quantity of this same Fluid with a greater Velocity, into what muscular Fibres it pleases, to account for the voluntary strong Action of the Muscles.

64. But fince such a superficial Account would not be satisfactory, it will be expected, that the principal Phænomena of these three Functions, should be explained by the Means of such a Fluid as has been supposed; and that the several Objections against this Doctrine should be answered: Let us attempt this; and where we cannot extricate ourselves from Dissipulties which may be thrown in, let us honestly ac-

knowledge Ignorance.

e

65. a. If Water, with a very small Proportion of Oils and Salts from the Earth, proves a structure of Nourishment for Vegetables, such a Liquor as the Fluid of the Nerves has been described, (§ 56.) may not be unsit for repairing the Waste in Animals.

B. The flow continual Motion of this nervous Fluid (§ 58. 59.) to the most minute Parts of the Body (§ 10.), is well enough calculated to supply the Particles that are constantly worn

off from the Solids by the Circulation of the

Liquors, and necessary Actions of Life.

r. The greater proportional Size of the Encephalon in young Creatures, than in Adults, feems calculated for their greater proportional Growth: For the younger the Animal is, the larger Encephalon and speedier Growth it has.

s. A Palfy and Atrophy of the Members generally accompanying each other, shew, that Nourishment, Sensation, and Motion, depend

on the same Cause.

It was faid (§ 26.), that the Nerves were principal Instruments in Nutrition: It was not affirmed, that they were the fole Instruments; and therefore an Atrophy may proceed from the Compression or other Lesion of an Artery, without being an Objection to the Doctrine here laid down.

63. a. All Objects of Sense, when applied to their proper Organs, act by Impulse; and this Action is capable of being increased by increafing the impelling Force. In tangible Objects, this is clearly evident; the closer they are pressed to a certain Degree, the more distinct Perception enfues. Odorous Particles need the Afliftance of Air moved rapidly, to affect our Nose: Sapid Substances, that are scarce sufficient to give us any Idea of their Tafte by their own Weight, are affifted by the Pressure of the Tongue upon the Palate: The Rays of Light collected drive light Bodies before them: Sound communicates a Vibration to all Bodies in harmonic Proportion with it.

The

The Impulse made thus by any of these Objects on the foft pulpy Nerves (§ 21.), which are full of Liquor, presses their Sides or Extremities, and their Liquor is hindered to flow fo freely as it did. The Canals being all full (§ 58.), this Refistance must instantaneously affect the whole Column of Fluids in the Canals that are pressed, and their Origins, and have the same Effect, as if the Impulse had been made upon the Origin itself. To illustrate this by a gross Comparison; let any one push Water out of a Syringe, through a long flexible Pipe fixed to the Syringe, and he is fensible of Resistance or a Push backwards, the Moment any one stops the Orifice of the Pipe, or closes the Sides of it with his Fingers. This Impulse made on the Nerves, and thus communicated to their Origin, varies according to the Strength or Weakness, the Quickness or Slowness, the Continuance or speedy Removal, the Uniformity or Irregularity, the Constancy or Alternation, &c. with which Objects are applied to the Nerves.

b. Whenever any Object is regularly applied with due Force to a Nerve, rightly disposed to be impressed by it, and is communicated, as just now explained, to the Sensorium, it gives a true and just Idea of the Object to the Mind.

c. The various Kinds of Impulses which the different Classes of Objects make, occasion in Animals which ought to have accurate Perceptions of each Object, a Necessity of having the different Organs of the Senses variously modified, so that the several Impulses may be regularly applied to the Nerves in each Organ; or

in

in other Words, we must have different Organs of the Senses fitted to the different Classes of

Objects.

d. As the Objects have one common Property of Impulse, so all the Organs have most of the Properties of the Organ of Touching in common with the Papillæ of the Skin. In the Nose and Tongue, this is evident: In some Operations on the Eyes we can also perceive this; as we may likewise do in some Cases where Matter is collected in the internal Ear.

e. These Properties common to the different Objects and Organs, occasion frequently uncommon Effects in the Application of an Object to an Organ proper to another Object of Sensation; for sometimes we have the same Idea, as if the Object had been applied to its own proper Organ: At other Times the Object is as it were changed, and we have the Idea as if the Organ had had its own proper Object applied to it. Thus, for Example, Light is the proper Object to be applied to the Eye, to give us any Idea of Colours; yet when all Light is excluded from the Eyes, an Idea of Light and Colours may be excited in us by Coughing, Sneezing, Rubbing or Striking the Eye-ball.—A Cane vibrating, fo as not to excite Sound perceptible to the Ear, applied to the Teeth, raises a strong Idea of Sound; as a little Insect creeping in the Meatus auditorius also does. The Fingers applied to two rough Surfaces, rubbing on each other, are fensible of the Sound they make; Surgeons of any Practice in the Cure of fractured Bones, can bear Witness to the Truth of this. The Fingers dip-

f. If the Impulse of an Object is applied with due Force, but irregularly, a confused Idea of the Object is raised. Distant Objects are confused to Myopes, as very near ones are to Pres-

bytæ.

g. If the Application of the Impulse is regular, but the Force with which it is applied is too weak, our Perception of the Object is too faint. One may whisper so low, as not

to be heard.

h. If the Application of Objects is too violent, and there is any Danger of the tender Organs of our Senses being hurt or destroyed, an uneasy Sensation, we call Pain, is raised, whatever the Organ thus injured is. The Object of seeling affects every Organ: Thus Pressure, stretching, cutting, pricking, acrid Salts, pungent Oils, great Heat, violent Cold, &c. occasion Pain, where-ever they are applied. Besides this, every particular Organ can be affected with Pain, by the too violent Application

Light pains the Eyes; very loud Sound stuns the Ears; very odorous Bodies and too sapid Objects hurt the Nose and Tongue. A pretty sure Proof this, that the Objects of our Senses all act, and that the Organs are all impressed,

in nearly the same Way.

i. Since a middle Impulse, neither too small nor too great, is necessary for a clear Percen tion of Objects, we would often be in Danger of not distinguishing them, if we were not subjected to another Law, to wit, that numerous Impulses made at once, or in a quick Succession to each other, increase our Perceptions of Ob jess. Thus fuch Sound as would not be heard on a Mountain Top, will be distinctly heard in a Wainscotted Chamber.—We feel much more clearly a tangible Object when our Finger is drawn alongst it, than when applied with the same Force, but by a fingle Pressure upon it. --- We make repeated Applications of odorous and fapid Objects, when we wish to fmell or taste accurately.

k. Whenever the uneafy Sensation, Pain, is raised by the too strong Application of Objects, a Sort of Necessity is as it were imposed upon the Mind, to endeavour to get free of the injuring Cause, by either withdrawing the grieved Part of the Body from it, as one retires his Hand when his Finger is pricked or burnt; or the injuring Cause is endeavoured to be forced from the Body, as a Tenesmus pushes acrid Faces out of the Rectum. In both these Operations, a convulsive Contraction is immediately made in the lesed Part, or in the Neigh-

tuch

apid

etty

nses led,

nall

cep.

ger ub-

ous

Db.

ard

ard

in-

ed

of of

to

15 b-

ed

of

1e

10

or

es

1-

1-

Neighbourhood of it; and if the Irritation is very strong or permanent, the greater Part of the nervous System becomes affected in that spasmodic or convulsive Way.—Is it this Necessity which obliges the Mind to exert herself in Respiration, or in the Action of the Heart, when the Lungs or Heart are gorged with Blood? or the Iris to contract the Pupil, when the Eye is exposed to strong Light? or Successing to be performed when the Nose is tickled? &c.

This Nisus of the Mind to free the Body of what is in Danger of being hurtful, may ferve to explain the Phænomena of a great many Diseases, when we are acquainted with the Distribution of the particular Nerves; and from this we can understand the Operation of Medicines that stimulate; and may learn how, by exciting a sharp, but momentary Pain, we may free the Body of another Pain that would be more durable; and that, by having it thus in our Power to determine a Flow of the Liquor of the Nerves to any particular Part, for the Benefit of that Part, or the Relief of any other difeased Part, we can do considerable Service by a right Application of the proper Medicines.

l. If a Pain-giving Cause is very violent or long continued, it destroys the Organs either irrecoverably, or puts them so much out of Order, that they only gradually recover. People have been made blind or deaf for all their Lives after a violent Essect of Light on their Eyes, or of Sound on their Ears; and we are frequently exposed to as much Light and Sound

Hh

25

as to make us unfit to fee or hear for a confiderable Time. I would explain this by a Ligature put round the tender Branch of an Herb. This Ligature drawn to a certain Degree, may weaken the Canals fo as to be unfit for the Circulation of the Juices a good While, till they are gradually explicated and made firm by these Juices: A stricter Ligature would disorder the Structure of the Fibres so much, that the Liquors could not recover them. The Analogy is so plain, that it needs no Commentary.—Thus the Insluence of a Nerve tied with an Artery in the Operation of an Aneurism may cease for some Time, but be afterwards recovered.

66. 1. In applying the Fluid of the Nerves to the Action of Muscles, it was said, that the natural or involuntary Contraction of Muscles was the Nisus which the nervous Fluid slowing constantly into the muscular Fibres makes to distend these Fibrils, by enlarging their transverse Diameters and shortning their Axes; and that voluntary Contraction was owing to a greater Quantity of that nervous Liquor determined towards the Muscle to be put in Action, and poured with greater Momentum into the muscular Fibrils, by the Power of the Mind willing to make such a Muscle to act, or obliged to do it by an irritating Pain giving Cause (§ 65. k.)

2. It has been objected to this Account of muscular Motion, that, if it was true, the Volume of the Muscle in Contraction necessarily would be considerably increased by so much Liquor poured into its Fibrils; whereas it does

not appear, by any Experiment, that the Volume of a Muscle is increased by its being put into Action.

ji-

b.

ay

ne

ill

by

r.

at

4-

n.

u-

1-

es

at

id

es

V-

1-

it

r

0

11

).

3. To this it has been answered, r. That when the Axes of muscular Fibres are shortened, and their transverse Diameters are enlarged; the Capacities of their Fibres, and confequently their Volume may not be changed, the Diminution one Way ballancing the Increase in the other.—2. That the Spaces between the muscular Fibres are sufficient to lodge these Fibres when they swell, during the Contraction of a Muscle, without any Addition to its Bulk; and that it plainly appears that these Spaces between the Fibrils are thus occupied, by the Compression which the larger Vessels of Mulcles, which run in those Spaces, luffer during the Action of the Muscle; it is so great as to drive the Blood in the Veins with a remarkable accelerated Velocity.

4. Another Objection to the Action of Muscles being owing to the Influx of a Fluid into their Fibrils is, That muscular Fibres are distractile, or capable of being stretched; and therefore, when a Fluid is poured into their hollow Fibrils, they would be stretched longitudinally, as well as have their transverse Diameters increased; that is, a Muscle would become longer, as well as thicker, when it is put into Action; whereas it is known to every Body,

that a Muscle is shortened while it acts.

5. In Answer to this, it has been remarked, That though muscular Fibrils are distractile, yet they will not yield to, or be stretched by every Force, however small, that might be H h 2 applied

applied to them. A Cord that can be stretch. ed in Length by the Weight of a Pound or two, would not yield in the least to an Ounce or two; and it must likewise be observed, that gradually as any Body is stretched, its Resistance to the stretching Force increases. Rope may be stretched to a certain Length by a Pound Weight appended to it, which would require two Pounds to stretch it very little further; and therefore the general Observation of Animal Fibres being distractile, cannot be a reasonable Objection to the Account of muscular Motion above mentioned, unless a Proof is brought that the Force which the Liquid of the Nerves must exert upon each Fibre of a Muscle in Order to make a Muscle act, is capable of distracting or stretching the Fibres; which has not yet been attempted to be proved.—It would appear from the Pain caufed by too great an Effort of Muscles, especially in weak People, that muscular Fibres can bear very little Distraction without Danger of a Solution of Continuity.

6. Muscles ceasing to act when their Arteries are tied or cut, and being brought into Motion by injecting Liquors into these Arteries even of a dead Animal, has been mentioned as Objections to the nervous Influence causing

their Contractions.

To the first of these Experiments it may be answered, That the tying or cutting of the Nerves sooner produces the Essect of making the Contraction cease than stopping the Insux of the arterious Blood does; and it will be universally allowed, that the Insux of Blood into

into Muscles is necessary for performing their

Functions right.

or

at

A

y

d

n

f

Whoever compares the Motion, which injecting Water or any other Liquor into the Arteries of a dead Animal, causes in its Muscles, will not compare it to what Contraction, whether voluntary or excited by Irritation, he may see in a living one.

7. If muscular Motion depends on the Influx of the nervous Liquid, the instantaneous Contraction of a Muscle, when the Mind wills to make it act, will be easily understood, from the Nerves being always full of their Li-

quor (§ 58. 64. a.)

8. If either the Nerves of any Muscle do not furnish a sufficient Quantity of their Liquor, or if the Fibres of a Muscle become too easily distractile, such a Muscle will be unactive or paralytic.

9. If too great a Quantity of the Liquor of the Nerves is determined to a Muscle or Muscles, by any Cause which the Mind cannot command, such Muscle or Muscles will be con-

vulled.

Nerves is not uniform, but by Disease becomes irregular, an alternate Relaxation and Contraction of Muscles may be the Consequence. Hence trembling Palsies, chorea fancti Viti, &c. Hence also the convulsive Tremors which Animals have when they lose much Blood.

Mr. Though the Nerves may not furnish as much Liquor as may be sufficient to make Muscles contract with Strength enough to overcome the Resistances to their Action, yet there may

H h 3

be a sufficient Quantity of Liquor in the Nerves to allow the Impressions of Objects to be conveyed to the Sensorium. This may be one Cause of a Member's being sometimes sensible after it cannot be moved.

quires some Energy in the Brain, which we have no Reason to think the Circulation of the Fluids in the Vessels can give it, or unless it has other Properties than what we can discover in it, or unless there is an Agent determining its Momentum towards some particular Parts which we are not conscious of; if some of these, I say, do not obtain, the Action of the Heart continuing of equal Force to propel our Liquors, notwithstanding all the Resistances that are to it, is not to be explained.

13. All Muscles, but especially the Heart, continue to contract in an irregular Way, aster they are cut away from the Animal to whom they belonged; which may be owing to the Liquors continuing to flow in the small Vessels, and being poured irregularly into the

muscular Fibrillæ.

14. It is faid that a Muscle cut out of the Body continues some Time to be capable of Contraction, whereas by tying its Arteries or Nerves, while it is otherwise entire in the Body, it loses its contracting Power, which therefore does not depend on these Organs, the Arteries or Nerves.

The Loss of the Power of acting when the Arteries or Nerves are tied while the Muscle is in the Body, is denied by some who made the Trial, and it might be expected that the

Motion

Motion would be more conspicuous when there is no Resistance to it, as is the Case when it is cut away from all the Parts it is connected with, which make greater Resistance to its contractible Essorts while it is entire in the Body, than when it is separated from these, and has no Resistance.

15. After the Heart, or any other Muscle cut away from an Animal, has ceased to contract, its Contraction may again be restored, by breathing upon it, or pricking it with any sharp Instrument. That Heat or Pricking should, by their Stimulus (§ 64. k.), occasion Contraction in a living Creature, may be understood; but how they should have the same Essect in a Muscle separated from an Animal, I know not.

67. Some have thought the Ganglions of Nerves (§ 18. 19, 20.) to be glandular, and to perform a Secretion. — Others from their firm Texture suppose them to be muscular, and to ferve to accelerate the Motion of the Liquor in the Nerves which proceed from them; but as no Proof is offered of either of these Opinions, they cannot be maintained.—Others would make them ferve, 1. To divide a small Nerve into many Nerves, and by these Means to increase the Number of nervous Branches. 2. To make Nerves come conveniently by different Directions to the Parts to which they belong. 3. To reunite several small nervous Fibres into one large Nerve. - If these three Things happen where there are no Ganglions, which is the Cafe, and

if no Proof is brought that these Things could not be done unless by the Interposition of a Ganglion; we must continue to acknowledge Ignorance concerning the Uses of these Knots, the Ganglious.

Of the PARTICULAR NERVES.

Pair of Nerves in all; of which ten come out from the Encephalon, and the other thirty have their Origin from the spinal Marrow.

Of the ten Pair of Nerves which come from the Encephalon, the first is the OLFACTORY, which long had the Name of the Mammillary Processes of the Brain, because in the Brutes, Cows and Sheep, which were most commonly diffected by the Antients, the anterior Ventricles of the Brain are extended forwards upon these Nerves, and adhere so firmly to them, that they feem to make the upper Side of the Nerves. Each of them being large, where it begins to be stretched out; and gradually becoming finaller, as it approaches the cribriform Bone, was imagined to refemble a Nipple. Those who mistook the Ventricles for Part of the Nerves, observing the Cavity in them full of Liquor, concluded, that thefe olfactory Nerves ferved to convey the fuperfluous Moisture of the Brain to the Holes of the Ethmoid Bone through which it passed into the Noie. But in Man, the Ventricles of whole

whose Brain are not thus extended forwards, these Nerves are small, long, and without any Cavity, having their Origin from the corpora striata, near the Part where the internal carotid Arteries are about to fend off their Branches to the different Parts of the Brain: and in their Course under the anterior Lobes of the Brain, which have each a Depression made for lodging them, the human olfactory Nerves become larger, till they are extended to the cribriform Bone; where they split into a great Number of small Filaments, to pass through the little Holes in that Bone; and being joined by a Branch of the fifth Pair of Nerves, are spread on the Membrane of the Nofe.

The tender Structure and sudden Expansion of these Nerves on such a large Surface, render it impossible to trace them far; which has made some Authors deny them to be Nerves: But when we break the Circumference of the cribriform Lamella, and then gently raise it, we may see the Distribution of the Nerves some Way on

the Membrane of the Nose.

The Contrivance of defending these long soft Nerves from being too much pressed by the anterior Lobes of the Brain under which they lie, is singular; because they have not only the prominent orbitar Processes of the frontal Bone to support the Brain on each Side, with the Veins going into the longitudinal Sinus, and other Attachments bearing it up, but there is a Groove formed in each Lobe of the Brain itself for them to lodge in.—Their splitting into so many small Branches before they enter the Bones

Bones of the Scull, is likewise peculiar to them; for generally the Nerves come from the Brain in disgregated Filaments, and unite into Cords, as they are going out at the Holes of the Bones. This Contrivance is the best for answering the Purpose they are designed for, of being the Organ of Smelling; for had they been expanded upon the Membrane of the Nose into a medullary Web, such as the optic Nerve forms, it would have been too sensible to bear the Impressions of such Objects as are applied to the Nose; and a Distribution in the more common Way, of a Cord sending off Branches, would not have been equal enough for such an Organ of Sensation.

The 2d Pair of Nerves, the OPTIC, rising from the thalani Nervorum opticorum, make a large Curve outwards, and then run obliquely inwards and forwards, till they unite at the Forepart of the Sella Turcica; where they divide, and each runs obliquely forwards and outwards to go out at its proper Hole in the sphenoid Bone, accompanied with the ocular Artery, to be extended to the Globe of the Eye, within which each is expanded into a very fine Cuplike Web, that lines all the Inside of the Eye as far forwards as the ciliary Circle, and is universally known by the Name of Retina.

Though the Substance of this Pair of Nerves feems to be blended at the Place where they are joined, yet Observations of People whose optic Nerves were not joined, and of others who were blind of one Eye from a Fault in the optic Nerve, or in those who had one of their Eyes taken out, make it appear, that there is

110.

n

le

d

Nerve of the affected Side only being wasted, while the other was large and plump. And the same Observations are contradictory to the Doctrine of a Decustation of all the Nerves (§ 8.); for the Disease could be traced from the affected Eye to the Origin of the Nerve on the same Side. In many Fishes indeed, the Doctrine of Decustation is favoured; for their optic Nerves plainly cross each other, without any Union at the Part where they are joined in Men and most Quadrupeds.

Those People whose optic Nerves were not joined, having neither seen Objects double, nor turned their Eyes different Ways, is also a plain Proof, that the Conjunction of the optic Nerves will not serve to account for either the uniform Motions of our Eyes, or our seeing Objects single with two Eyes, though it may be one Cause of the remarkable Sympathy of the one

Eye with the other in many Diseases.

The Retina of a recent Eye, without any Preparation, appears a very fine Web, with some Blood-vessels coming from its Center to be distributed on it; but, after a good Injection of the Arteries that run in the Substance of this Nerve, as is common to other Nerves, it is with Difficulty that we can observe its nervous medullary Substance.——The Situation of these Vessels in the central Part of the optic Nerve; the Want of medullary Fibres here, and the Firmness of this Nerve before it is expanded at its Entry into the Ball of the Eye, may be the Reason why we do not see such Bodies, or Parts of Bodies, whose Picture falls on this cen-

tral Part of the Retina. - An Inflammation in thole Arteries of the Retina, which feveral Fevers and an Opthalmia are generally attended with, may very well account for the Tender. ness in the Eyes, and Inability to bear the Light, which People have in these Diseases .-The Over-distension of these Vessels may like. wife ferve to account for the black Spots obferved on bright-coloured Bodies especially, and for that fmoaky Fog through which all Oh. jects are feen by People in fome Fevers.——If those Vessels lose their Tone, and remain preternaturally diftended, no Objects affect our Retina, though the Eye externally appears found; or this may be one Caufe of an Aman. rosis or Gutta serena. --- From a partial Distenfion of these Vessels or Paralysis of a Part of the Retina, the central Part, or the Circumference, or any other Part of Objects, may be loft to one or both Eves.

The THIRD PAIR rife from the anterior Part of the Processus annularis, and piercing the Dura Mater a little before, and to a Side of the Ends of the posterior clinoid Process of the sphenoid Bone, run along the Receptacula, or cavernous Sinuses, at the Side of the Ephippium, to get out at the Foramina lacera; after which each of them divides into Branches, of which one, after forming a little Ganglion, is distributed to the Globe of the Eye; the others are fent to the Musculus rectus of the Palpebra, and to the Attollens, Adductor, Deprimens, and obliquus minor Muscles of the Eye-ball. These Muscles being principal Instruments in the Motions of the Eye-lid and Eye-ball, this Nerve

Nerve has therefore got the Name of the Motor oculi. - I have frequently observed in Convulfions the Eye-lids widely opened, the Cornea turned upward and outwards, and the Eye-balls fank in the Orbit; which well described the conjunct Action of the Muscles which this Pair of Nerves ferves.—The Distension of a confiderable Branch of the Carotid, which passes over this Nerve near its Origin on each Side, may possibly be the Reason of that Heaviness in the Eye-lids and Eyes, after drinking hard,

or eating much.

eb-

7. If

e.

ll' 1'5

u-

nle

e,

to

of

a,

er

of

is

rs

a,

nd

le

10

is

re

The FOURTH PAIR, which are the smallest Nerves of any, derive their Origin from the Back-part of the Base of the Testes; and then making a long Course on the Side of the annular Protuberance, enter the Dura Mater a little farther back, and more externally than the third Pair, to run also along the Receptacula, to pass out at the Foramina lacera, and to be entirely spent on the Musculi trochleares, or superior oblique Muscles of the Eyes. These Muscles being employed in performing the rotatory Motions, and the Advancement of the Eye-balls forward, by which feveral of our Passions are expressed, the Nerves that serve them have got the Name of PATHETICI. -Why these small Nerves should be brought fo far to this Muscle, when it could have been supplied easily by the Motor oculi, I know not.

The FIFTH PAIR are large Nerves, rifing from the annular Processes, where the medullary Processes of the Cerebellum join in the Formation of that Tuber, to enter the Dura

Mater

Mater near the Point of the petrous Process of the Temporal Bones; and then sinking close by the Receptacula at the Sides of the Sella Turcica, each becomes in Appearance thicker, and goes out of the Scull in three great Branches.

The first Branch of the fifth is the OPH. THALMIC, which runs through the Foramen lacerum to the Orbit, having in its Passage thither a Connexion with the fixth Pair. It is afterwards distributed to the Ball of the Eye with the third; to the Nose, along with the Olfactory, which the Branch of the fifth that passes through the Foramen orbitarium internum joins, as was already mentioned in the Description of the first Pair. This ophthalmic Branch likewise supplies the Parts at the internal Canthus of the Orbit, the Glandula lacrymalis, Fat, Membranes, Muscles and Teguments of the Eye-lids; its longest farthest extended Branch passing through the Foramen superciliare of the Os frontis, to be distributed to the Fore-head.

The small Fibres which this sirst Branch of the fifth and the third Pair of Nerves send to the Eye-ball, being situated on the optic Nerve, and, after piercing the sclerotic Coat, running along the choroid Coat on the Outside of the Retina in their Course to the Uvea or Iris, may be a Cause of the Sympathy between the optic Nerve and the Uvea; by which we more readily acquire the Habit of contracting the Iris, and thereby lessen the Pupil, when too strong Light is excluded, and, on the contrary, enlarge the Pupil, when the Light is too faint.—This, with the Sympathy which must arise from

fome

t

17

0

some of the Nerves of the Membrane of the Nostrils, being derived from this first Branch of the fifth Pair of Nerves, may also be the Cause, why an Irritation of the Retina, by too strong Light, may produce Sneezing, as if a Stimulus had been applied to the Membrane of the Nose itself; --- why pressing the internal Canthus of the Orbit, sometimes stops Sneezing; -why Irritation of the Note causes the Eyelids to shut convulsively, and makes the Tears to flow plentifully; and why Medicines put into the Nose, do often great Service in Diseases of the Eyes.—In the Megrim all the Branches of this Nerve discover themselves to be affected; for the Forehead is racked with Pain, the Eye-ball is pained, and feels as if it was fqueezed, the Eye-lids that convulfively, the Tears trickle down, and an uneasy Heat is felt in the Nofe. Hence we can understand, where external Medicines will have the best Effect, when applied to remove this Disease, to wit, to the Membrane of the Nose, and to the Forehead; -why alternate Pressure near the superciliary Hole of the frontal Bone, or Sneezing, sometimes gives immediate Relief in the Megrim; -why the Sight may be loft by an Injury done to the fupra-orbitar Branch; how it may be restored by Agitation of that Branch of this Nerve.

The fecond Branch of the fifth Pair of Nerves may be called MAXILLARIS SUPERIOR, from its ferving principally the Parts of the Upper Jaw. It goes out at the round Hole of the sphenoid Bone, and fends immediately one Branch in the Channel on the Top of the

Ii 2

Antrunz:

Antrum maxillare; the Membrane of which and the upper Teeth are supplied by it in its Passage. As this Branch is about to go out at the Foramen orbitarium externum, it fends a Nerve through the Substance of the Os maxillare to come out at Steno's Duct, to be distributed to the Fore-part of the Palate; and what remains of it escaping at the external orbitar Hole, divides into a great many Branches, that fupply the Cheek, upper Lip, and Nostril. The next considerable Branch of the Superior maxillary Nerve, after giving Branches to join the Intercostal where it is passing through the Skull with the carotid Artery, is fent into the Nose by the Hole common to the Palate and sphenoidal Bone; and the remaining Part of this Nerve runs in the Palato-maxillaris Canal, giving off Branches to the Temples and pterygoid Muscles, and comes at last into the Palate to be lost. Hence the Ach in the Teeth of the Upper Jaw occasions a gnawing Pain deepfeated in the Bones of the Face, with Swelling in the Eye-lids, Cheek, Nose, and upper Lip; and on the other hand, an Inflammation in these Parts, or a Megrim, is often attended with sharp Pain in the Teeth.-Hence an Obstruction in the Duct of the maxillary Sinus, which obliges the Liquor fecreted there to find out a preternatural Route for itself, may be occasioned by the Pain of the Teeth.-Hence the upper Lip often fuffers when the Palate or Nose is ulcerated.

The third or MAXILLARIS INFERI-OR Branch of the fifth Pair going out at the oval Hole of the fphenoid Bone, ferves the

Muscles

Muscles of the Lower Jaw, and the Muscles. fituated between the Os byoides and Faw: All. the falivary Glands, the Amygdalæ, and the external Ear, have Branches from it: It has a large Branch loft in the Tongue, and fends another through the Canal in the Substance of the Lower Jaw, to serve all the Teeth there, and to come out at the Hole in the Fore-part of the Jaw, to be lost in the Chin and under Lip.—Hence a convulsive Contraction of the Muscles of the Lower Jaw, or the Mouth's being involuntarily thut, a great Flow of Spittle or Salivation, a Pain in the Ear, especially in Deglutition, and a Swelling all about the Throat, are natural Consequences of a violent Irritation of the Nerves of the lower Teeth in the Toothach; and Pain in the Teeth and Ear, is as natural a Consequence of an Angina. -- Hence alternate Pressure on the Chin may sometimes relieve the Violence of a Tooth-ach. -- Hence destroying the Nerves of a Tooth by actual or potential Cauteries, or pulling a carious Tooth, fo often removes immediately all these Symptoms. -- Hence no Cure is to be found for fome Ulcers in the Upper or Lower Jaw, but by drawing a Tooth.—Hence in Cancers of the under Lip, the falivary Glands are in Danger of being affected, or the Disease may be occasioned to the Lip by its beginning in the Glands .-Perhaps the Sympathy of the Organs of Tafting and Smelling, may in some Measure depend on their both receiving Nerves from the fifth Pair.

The SIXTH PAIR, which is the smallest except the fourth, rises from the Fore-part of Li 2 the

the Corpora pyramidalia; and each entering the Dura Mater, some Way behind the posterior clinoid Process of the sphenoid Bone, has a long Courfe below that Membrane, and within the Receptaculum at the Side of the Sella Turcica, where it is immersed in the Blood of the Receptacle, but for what Purpose I am ignorant. It goes afterwards out at the Foramen lacerum into the Orbit, to serve the abductor Muscle of the Eye. A Defect in this Nerve may therefore be one Caufe of a Strabifmus. In the Passage of this Nerve below the Dura Mater, it lies very contiguous to the internal carotid Artery, and to the ophthalmic Branch of the fifth Pair of Nerves. At the Place where this fixth Pair is contiguous to the Carotid, a Nerve either goes from each of them in an uncommon Way, to wit, with the Angle beyond where it rifes obtuse, to descend with the Artery, and to form the Beginning of the intercostal Nerve, according to the common Description; or, according to other Authors, this Nerve comes up from the great Ganglion of the Intercostal, to be joined to the fixth here.

The Arguments for this latter Opinion are, That, according to the common Doctrine, this Beginning of the intercostal Nerve, as 'tis called, would rise in a Manner not so ordinary in Nerves. In the next Place, it is observed, that the sixth Pair is larger nearer to the Orbit, than it is before it comes to the Place where this Nerve is said to go off; and therefore it is more probable, that it receives an Addition there, rather than gives off a Branch. Lastly, It is found,

that

that upon cutting the intercostal Nerves of living Animals, the Eyes plainly were affected; they lost their bright Water; the Gum, or Gore, as we call it, was separated in greater Quantity; the Pupil was more contracted; the cartilaginous Membrane, at the internal Canthus, came more over the Eye; and the Eyeball itself was diminished.

To this it is answered, in Defence of the more common Doctrine, 1st, That other Branches of Nerves go off in a reflected Way, as well as this does, supposing it to be the Beginning of the Intercostal; and that the Reflexion would rather be greater, if it is thought to come up from the Intercostal to the fixth. adir. It is denied that this Nerve is for ordinary thicker at its Fore than its Back part; and if it was supposed to be thickest nearest to the Orbit, the Conclusion made above could not be drawn from this Appearance, because other Nerves enlarge fometimes where there is no Addition made to them, as in the Instance already mentioned of the Trunk of the fifth Pair while below the Dura Mater. 3dly, The Experiments on living Animals shew indeed, that the Eyes are affected upon cutting the intercostal Nerve, but not in the Way which might have been expected, if the Intercostal had furnished such a Share of the Nerve that goes to the Abductor Muscle of the Eye; for it might have been thought, that this Muscle would have been so much weakened immediately upon cutting the Intercostal, that its Antagonist the Adductor would have greatly prevailed over it, and have turned the Eye strongly in towards the Noie; which

is not faid to be a Confequence of this Experiment. So that the Arguments are still equivocal; and more Observations and Experiments must be made, before it can be determined with Certainty, whether the sixth Pair gives or receives a Branch here. In the mean Time, I shall continue to speak about the Origin of the Intercostal with the Generality of Anatomists.

At this Place where the Intercostal begins, the fifth Pair is contiguous and adherent to the fixth; and it is generally faid, that the ophthalmic Branch of the fifth gives a Branch or two to the Beginning of the Intercostal, or receives fuch from it. Others deny any fuch Communication between them, and those who affirm the Communication confess, that in Some Subjects they could not see it. After examining the Nerves here in a great many Sabjects, I cannot determine whether or not there are nervous Filaments going from the one to the other. Sometimes I have thought that I traced them evidently; at other times I obferved that what I diffested for nervous Filaments, was collapsed cellular Substance; and in all the Subjects where I had pushed an Injection successfully into the very small Arteries, I could only observe a Plexus of Vessels connecting the one to the other. In any of these Ways, however, there is as much Connexion, as we are affired from many Experiments and Observations on other Nerves, is sufficient to make a very great Sympathy among the Nerves here.—Possibly the Appearances in the Eyes

Eyes of Dogs, whose intercostal Nerves were

cut, might be owing to this Sympathy.

eri.

VO-

nts

ed

res

ie,

of

to-

ns,

he

h-

10

e-9

ch

10

in

4

1

re

0

I

b--

a.

nd

ŋ-

S,

1-

(e

n,

ts

It

e

e

35

The SEVENTH PAIR comes out from the lateral Part of the annular Process, behind where the medullary Processes of the Cerebellum are joined to that Tuber; and each being accompanied with a larger Artery than most other Nerves, enters the internal Meatus auditorius, where the two large Bundles of Fibres, of which it appeared to confift within the Scull, foon separate from each other; one of them entering by feveral small Holes into the Vestible, Cochlea, and semicircular Canals, is stretched on this inner Camera of the Ear in a very foft pulpy Substance; and being never feen in the Form of a firm Cord, fuch as the other Parcel of this and most other Nerves become, is called PORTIO MOLLIS of the auditory Nerve.

The other Part of this seventh Pair passes through Galen's Foramen cacum, or Fallopius's Aquaduct, in its crooked Passage by the Side of the Tympanum; in which Passage, a Nerve fent from the lingual Branch of the inferior maxillary Nerve, along the Outside of the Tuba Eustachiana, and cross the Cavity of the Tympanum, where it has the Name of Chorda Tympani, is commonly faid to be joined to it. The very acute Angle which this Nerve makes with the fifth, or the fudden violent Reflexion it would fuffer on the Supposition of its coming from the fifth to the feventh, appears unusual; whereas, if we suppose that it comes from the seventh to the fifth, its Course would be more in the ordinary Way, and the Chorda

Tympani

Tympani would be esteemed a Branch of the seventh Pair going to join the fifth. This smaller Bundle of the seventh gives Branches to the Muscles of the Malleus, and to the Dura Mater, while it passes through the bony -crooked Canal, and at last comes out in a firm Chord named PORTIO DURA, at the End of this Canal, between the styloid and mafloid Processes of the temporal Bone, giving immediately Filaments to the little oblique Muscles of the Head, and to those that rife from the styloid Process. It then pierces through the parotid Gland, and divides into a great many Branches, which are difperfed in the Muscles and Teguments that cover all the Side of the upper Part of the Neck, the whole Face and Cranium, as far back as the Temples, including a confiderable Part of the external Its Branches having thus a confiderable Connexion with all the three Branches of the fifth Pair, and with the fecond Cervical, occasion a considerable Sympathy of these Nerves with it .- Hence, in the Tooth-ach, the Pain is sometimes very little in the affected Tooth, compared to what it is all along the Side of the Head and in the Ear. Hence probably the Relief of the Tooth-ach from Blisters applied behid the Ear, or a hot Iron touching the Antibelix of the Ear .- By this Communication or Connexion possibly too it is, that a vibrating String, held between one's Teeth, gives a strong Idea of Sound to the Person who holds it, which no body else can perceive. Perhaps too the Distribution of this

the

his ies

Du-

ny

2

he

20-

ng

ue

le

es

2

in

ne.

le

S,

al

3

e

C.

e

d

e

1

S

this Nerve occasions the Head to be so quickly turned upon the Impression of Sound on our Fars.

The EIGHT PAIR of Nerves rise from the lateral Bases of the Corpora olivaria in disgregated Fibres; and as they are entring the anterior internal Part of the Holes common to the Os occipitis and Temporum, each is joined by a Nerve which afcends within the Dura Mater from the tenth of the Head, the first and fecond cervical, and fometimes from inferior Nerves: This every Body knows has the Name of the NERVUS ACCESSORIUS. When the two get out of the Scull, the Accessorius separates from the eight, and, descending obliquely outwards, passes through the Sternomastoideus Muscle, to which it gives Branches, and afterwards terminates in the Trapezius and Rhomboid Muscles of the Scapula. In this Course it is generally more or less joined by the second cervical Nerve. - Why this Nerve, and feveral others which are distributed to Muscles, are made to pierce through Muscles, which they might have only passed near to, I do not know.

The large eight Pair, soon after its Exit, gives Nerves to the Tongue, Larynx, Pharynx, and Ganglion of the Intercostal Nerve, and being disjoined from the ninth and Intercostal, to which it adheres closely some way, runs streight down the Neck behind the internal jugular Vein, and at the external Side of the carotid Artery. As it is about to enter the Thorax, a large Nerve goes off from the eighth of each Side: This Branch of the Right-side

turns

turns round from the fore to the back Part of the Subclavian Artery, while the Branch of the Left-fide turns round the great Curve of the Aorta, and both of them mounting up again at the Side of the OE sophagus, to which they give Branches, are lost at last in the Larynx. These are called the RECURRENT Nerves, which we are defired to shun in the Operation of Bronchotomy, though their deep Situation protects them sufficiently. - The Muscles of the Larynx being in a good Meafure supplied with Nerves from the Recurrents. it is to be expected, that the cutting of them will greatly weaken the Voice, though it will not be entirely lost, so long as the superior Branches of the eight Pair are entire.—Why the recurrent Nerves rife fo low from the eight Pair to go round a large Artery, and to have fuch a long Courfe upwards, I know not.

The eight Pair at or near the Place where the recurrent Nerves go off from it, or frequently the Recurrents themselves, send off Nerves to the *Pericardium*, and to join with the Branches of the Intercostal that are distri-

buted to the Heart.

After these Branches are sent off, the Par vagum on each Side descends behind the great Branch of the Trachea, and gives numerous Filaments to the Lungs in going to the OEsophagus. The one of the Lest-side running on the Fore-part of the OEsophagus, communicates by several Branches with the right one in its Descent to be distributed to the Stomach: The right one gets behind the OEsophagus,

phagus, where it splits and rejoins several times before it arrives at the Stomach, to which it sends Nerves; and then being joined by one or more Branches from the Left-trunk, they run towards the cæliac Artery, there to join into the great semilunar Ganglion formed by the two Intercostals.

From the Distribution of this Par vagum, we may learn how tickling the Fauces, with a Feather or any such Substance, excites a Naufea and Inclination to vomit; ---- why Coughing occasions Vomiting, or Vomiting raises a Cough. Hence we fee how the nervous Althma and the Tuffis convulsiva, Chincough, are attended with a straitening of the Glottis; why Food difficult to digest occasions the Asthma to weakly People; and why Emetics have frequently cured the Afthma very speedily; -why an Attempt to vomit is fometimes in Danger of fuffocating Asthmatic People ; why the superior Orifice of the Stomach is so fenfible, as to be looked on as the Seat of the Soul by fome; - why People subject to Diftensions of the Stomach, have so often the Sensation of Balls in their Breasts and Throats; -why the Globus hystericus is so often attended with a violent Strangulation at the Glottis.

The NINTH PAIR of Nerves comes from the inferior Part of the Corpora pyramidalia, to go out of the Scull at their proper Holes of the occipital Bones. After their Egress they adhere for some way firmly to the eighth and Intercostal; and then sending a Branch, that in many Subjects is joined with

Branches of the first and second cervical Nerves, to be distributed to the thyroid Glands and Muscles on the Fore-part of the Trachea Arteria, the ninth is lost in the Muscles and Substance of the Tongue. Some have thought this Nerve, and others have esteemed the third Branch of the fifth Pair of Nerves to be the proper gustatory Nerve. I know no Observations or Experiments to prove either Opinion, or to assure us, that both Nerves do not serve for Tasting and for the Motion of the Tongue.—May not the Distribution of this Nerve to the Muscles below, as well as above the Os Hyoides, contribute to their acting more uniformly in depressing the Lower Jaw or Head?

The TENTH PAIR rises in separate

The TENTH PAIR rifes in separate Threads from the Sides of the spinal Marrow, to go out between the Os occipitis and first Vertebra of the Neck. After each of them has given Branches to the great Ganglion of the Intercostal, 8th, 9th, and 1st cervical Nerves, it is distributed to the streight, oblique, and some of the extensor Muscles of the Head. Whether the Name of the tenth of the Head, or of the sirst Vertebral, ought to be given to this Pair of Nerves, is of no such Consequence as to deserve a Debate, tho it has some

of the Marks of the spinal Nerves.

In the Description of the sixth Pair, I sollowed the usual Way of speaking among Anatomists, and called that the Beginning of the Intercostal Nerve which comes out of the Scull; and therefore shall here subjoin a cursory Description of this Nerve, notwithstanding its much larger Part is composed of Nerves coming

TIME

f

d

e

e

d

coming out from the *spinal Marrow*. There is no greater Incongruity in point of Method to fay, that the Nerve we are describing receives Additions from others that have not been described, than it is to repeat in the Description of a great many Nerves, that each of them gives Branches to form a Nerve which we are ignorant of; which is all the Difference between describing the Intercostal before or after the spinal Nerves.

The Branch reflected from the fixth Pair, joined possibly by some Filaments of the Ophthalmic Branch of the fifth, runs along with the internal Carotid Artery, through the crooked Canal formed for it in the temporal Bone, where the little Nerve is very foft and pappy, and in feveral Subjects divides and unites again, and is joined by one or more Branches from the fifth, particularly of its superior maxillary Branch, before it comes out of the Scull. May the Compression of this Nerve by the Carotid when stretched during the Systole, contribute to the Diastole of the Heart? As soon as the Nerve escapes out of this bony Canal, it is connected a little Way with the eighth and ninth; then separating from these, after seeming to receive additional Nerves from them, it forms a large Ganglion, into which Branches from the tenth of the Head, and from the first and fecond Cervical, enter. From this Ganglion the Nerves come out again small to run down the Neck along with the carotid Artery, communicating by Branches with the cervical Nerves, and giving Nerves to the Muscles that bend the Head and Neck. As the Intercostal is

Kk2

about to enter the Thorax, it forms another Ganglion from which Nerves are fent to the Trachea and to the Heart; these designed for the Heart joining with the Branches of the eighth, and paffing between the two great Arteries and the Auricles to the Substance of that Muscle. The Intercostal after this confisting of two Branches, one going behind, and the other running over the Fore-part of the Subclavian Artery, forms a new Ganglion where the two Branches unite below that Artery, and then descending along the Sides of the Vertebra of the Thorax, receives Branches from each of the dorfal Nerves; which Branches appearing to come out between the Ribs, have given the Name of Intercostal to the whole Nerve. Where the Addition is made to it from the fifth dorfal Nerve, a Branch goes off obliquely forwards; which being joined by such Branches from the fixth, seventh, eighth, and ninth Dorfal, an anterior Trunk is formed, and passes between the Fibres of the Appendix musculosa of the Diaphragm, to form, along with the other Intercostal and the other Branches of the eighth Pair, a large semilunar Ganglion fituated between the cæliac and superior mesenteric Arteries; the Roots of which are, as it were, involved in a Sort of nervous Net-work of this Ganglion, from which a great Number of very small nervous Threads run out to be extended on the Surface of all the Branches of those two Arteries, so as to be easily seen when any of the Arteries are stretched, but not to be raised from them by Dissection; and thus the Liver, Gall-Bladder, Duodenum, Pancreas, Spleen,

Spleen, Jejunum, Ilium, and a large Share of the Colon, have their Nerves fent from this great folar Ganglion or Plexus.—May the peristaltic Motion of the Intestines depend in some measure on the Passage of the Intercostal

Nerves through the Diaphragm?"

Several Fibres of this Ganglion, running down upon the Aorta, meet with other Nerves fent from the posterior Trunk of the Intercostal, which continues its Course along the Sides of the Vertebræ, they supply the Glandulæ renales, Kidneys and Testes in Men, or Ovaria in Women; and then they form a Network upon the inferior mesenteric Artery where the Nerves of the two Sides meet, and accompany the Branches of this Artery to the Part of the Colon that lies in the lest Side of the Belly, and to the Rectum, as far down as to the lower Part of the Pelvis.

The Intercostal continuing down by the Side of the Vertebræ of the Loins, is joined by Nerves coming from between these Vertebræ, and sends Nerves to the Organs of Generation and others in the Pelvis, being even joined with those that are sent to the inferior Ex-

tremities. 15 (1802)

The almost universal Connexion and Communication which this Nerve has with the other Nerves of the Body, may lead us to understand the following, and a great many more *Phænomena:* — Why tickling the Nose causes Sueezing. — Why the too great Quantity of Bile in the *Cholera* occasions Vomiting as well as Purging. — Why People vomit in Colicks, in Inflammations, or other Irritations of the K k 3

Liver, or of the Ducts going from it, and the Gall-bladder. Why a Stone in the Kidneys, or Ureters, or any other Cause irritating those Organs, should so much more frequently bring on Vomiting and other Diforders of the Stomach, than the Stone, or any other stimulating Cause in the Bladder does .- Why Vomiting is a Symptom of Danger after Child-birth. Lithotomy, and other Operations on the Parts in the Pelvis. Why the Obstructions of the Menfes are capable of occasioning Strangula. tions, Belching, Colicks, Stomach-aches, and even Convulsions in the Extremities. - Why Vesicatories applied from the Ears to the Clavicles of Children labouring under the Juffis Convultiva, are frequently of great Service. --- Why Worms in the Stomach or Guts excite an Itching in the Nofe, or Grinding of the Teeth. Why Irritations in the Bowels of the Belly occasion sometimes universal Convulfions of the Body.

The spinal Nerves rise generally by a Number of digregated Fibres from both the Fore and Back-part of the Medulla spinalis, and soon after form a little Knot or Ganglion, where they acquire strong Coats, and are extended into firm Cords. They are distinguished by Numbers, according to the Vertebrae from between which they come out; the superior of the two Bones forming the Hole through which they pass, being the one from which the Number is applied to each Nerve. There are generally said to be thirty Pair of them; seven of which come out between the Vertebrae of the Neck, twelve between those of

the

the Back, five between those of the Loins, and

fix from the falle Vertebræ.

The FIRST CERVICAL Pair of Nerves comes out between the first and second Vertebræ of the Neck; and having given Branch; es to join with the tenth Pair of the Head. the fecond Cervical and Intercostal, and to serve the Muscles that bend the Neck, it sends its largest Branches backwards to the Extensor Muscles of the Head and Neck; some of which piercing through these Muscles, run up on the Occiput to be lost in the Teguments here; and many Fibres of it advance so far forward, as to be connected with the Fibrils of the first Branch of the fifth Pair of the Head, and of the Portio Dura of the Auditory Nerve. Hence possibly it is that a Clavus hystericus changes suddenly sometimes from the Forehead to a violent Pain and Spalm in the Backpart of the Head and Neck.

The SECOND CERVICAL is foon joined by fome Branches to the ninth of the Head and Intercostal, and to the first and third of the Neck; then has a large Branch that comes out at the exterior Edge of the Sterno-massiodeus Muscle, where it joins with the Accessorius of the eighth Pair; and is afterwards distributed to the Platysma Myoides, Teguments of the Side of the Neck and Head, parotid Gland and external Ear, being connected to the Portio Dura of the Auditory Nerve, and to the first Cervical. The Remainder of this second cervical is spent on the Levator Scapulæ, and the Extensors of the Neck and Head. Generally a large Branch is here sent off to join

the

the Accessorius of the eighth Pair, near the fu.

perior Angle of the Scapula.

To the Irritation of the Branches of this Nerve it probably is, that, in an Inflammation of the parotid Gland, the Neck is pain. ed fo far down as the Clavicle, the Head is drawn towards the Shoulder of the affected Side, and the Chin is turned to the other Side.—In opening the external jugular Vein, no Operator can promise not to touch some of the cutaneous Branches of this Nerve with the Lancet; which occasions a sharp pricking Pain in the mean Time, and a Numnels of the Skin near the Orifice for some Time after.

The THIRD PAIR of the Neck passes out between the third and fourth cervical Vertebræ; having immediately a Communication with the second, and fending down a Branch, which, being joined by a Branch from the fourth cervical, forms the PHRENIC Nerve. This descending, enters the Thorax between the Subclavian Vein and Artery; and then being received into a Groove formed for it in the Pericardium, it has its Course along this Capfula of the Heart, till it is lost in the middle Part of the Diaphragm. The right Phrenic has a streight Course; but the left one is obliged to make a confiderable Turn outwards to go over the prominent Part of the Pericardium, where the Point of the Heart is lodged. Hence in violent Palpitations of the Heart, a pungent acute Pain is felt near the left Orifice of the Stomach — The middle of the Diaphragm scarce could have been supplied by any other Nerve which could have

had

had such a streight Course as the Phrenic has. If the Subclavian Artery and Vein have any Effect upon this Nerve, I do not know it.

The other Branches of the third cervical Nerve are distributed to the Muscles and Teguments at the lower Part of the Neck and Top of the Shoulder. No Wonder then that an Inflammation of the Liver or Spleen, an Abscess in the Lungs adhering to the Diaphragm, or any other Cause capable of irritating the Diaphragm, should be attended with a sharp Pain on the Top of the Shoulder, as well as Wounds, Ulcers, &c. of this Muscle itself.—If the Irritation of this Muscle is very violent, it may occasion that convulsive Contraction of the Diaphragm, which is called an Hiccough; and therefore an Hiccough in an Inflammation of the Liver has been justly declared to be an ill Symptom.

An Irritation of the thoracic Nerves which produces Sneezing, may sometimes free the phrenic Nerves from any Spasm they occasion; so that Sneezing sometimes takes away the Hiccough; and a Derivation of the Fluid of the Nerves any other Way may do the same Thing: Or the Hiccough may also be sometimes cured, by drawing up into the Nose the Smoak of burning Paper or other acrid Fumes, swallowing pungent or aromatic Medicines, and by a Surprize, or any other strong Application of the Mind in Thinking, or in distinguishing Objects: Or, when all these have sailed, it has been put away by the brisk Stimulus of a blistering Plaister applied to the Back.

The

The FOURTH CERVICAL Nerve, after sending off that Branch which joins with the third to form the Phrenic, and bestowing Twigs on the Muscles and Glands of the Neck, runs to the Arm-pit, where it meets with the FIFTH, SIXTH, and SEVENTH Cervicals, and FIRST DORSAL, that escape in the Interstices of the Musculi scaleni, to come at the Arm-pit, where they join, feparate, and re-join, in a Way scarce to be rightly expressed in Words; and after giving several confiderable Nerves to the Muscles and Teguments which cover the Thorax, they divide into feveral Branches, to be distributed to all the Parts of the superior Extremity. Seven of these Branches I shall describe under particular Names.

I. SCAPULARIS runs streight to the Cavitas semilunata of the upper Costa of the Scapula, which is a Hole in the recent Subject by a Ligament being extended from one Angle of the Bone to the other, giving Nerves in its Way to the Muscles of the Scapula. When it has passed this Hole, it supplies the Supra-spinatus Muscle; and then descending at the anterior Root of the Spine of the Scapula, it is lost in the other Muscles that lie on the Dorsum of that Bone.

the Axilla, to get below the Neck of the Head of the Os humeri, and to mount again at the Back-part of it; so that it almost surrounds the Articulation, and is distributed to the Muscles that draw the Arm back, and to those that raise

it up.

3. CUTA.

le

18

le

e

0

at-

al

1-

1-

e

ır

e

b-

ie

es

1.

ė

at

1,

e

at

d

le

e

3. CUTANEUS runs down the Fore-part of the Arm near the Skin, to which it gives off Branches; and then divides on the Inside of the Fore-arm into several Nerves, which supply the Teguments there, and on the Palm of the Hand.—In opening the Basilic Vein of the Arm, at the ordinary Place, the same Symptoms are sometimes produced, as in opening the external jugular Vein, and from a like Cause, to wit, from hurting a Branch of this cutaneous Nerve with the Lancet.

4. MUSCULO-CUTANEUS, or perforans Casserii, passes through the Coracobrachialis Muscle; and, after supplying the Biceps slexor cubiti and Brachiæus internus, passes behind the Tendon of the Biceps, and over the cephalic Vein, to be bestowed on the Teguments on the Outside of the Fore-arm and Back of the Hand.—This Nerve is sometimes hurt in opening the cephalic Vein, and causes Pain and Numbness for a short Time.

from the Axilla, under the Os humeri, and backward to the external Part of that Bone, supplying by the Way the extensor Muscles of the Fore-arm, to which it runs between the two brachiæi Muscles, and within the Supinator Radii longus.—At the Upper-part of the Fore-arm, it sends off a Branch, which accompanies the Supinator longus till it comes near the Wrist, where it passes obliquely over the Radius, to be lost in the Back of the Hand and Fingers.—The principal Part of this Nerve pierces through the supinator Radii brevis,

vis, to serve the Muscles that extend the Hand

and Fingers.

o. ULNARIS is extended along the Infide of the Arm, to give Nerves to the Mus. cles that extend the Fore-arm and to the Teguments of the Elbow: Towards the lower Part of the Arm, it flants a little backward to come at the Groove behind the internal Condyle of the Os humeri, through which it runs to the Ulna: In its Course along this Bone, it ferves the neighbouring Muscles and Teguments: and as it comes near the Wrist, it detaches a Branch obliquely over the Ulna to the Back of the Hand, to be lost in the convex Part of feveral Fingers. The larger Part of the Nerve goes streight forward to the internal Side of the Os pisiforme of the Wrist; where it fends off a Branch which finks under the large Tendons in the Palm, to go cross to the other Side of the Wrist, ferving the musculi lumbricales and interoffei, and at last terminating in the short Muscles of the Thumb and Forefinger. What remains of the ulnar Nerve after supplying the short Muscles of the Littlefinger, divides into three Branches; whereof two are extended along the Sides of the Sheath of the Tendons of the Flexors of the Littlefinger, to furnish the concave Side of that Finger; and the third Branch is disposed in the fame Way upon the Side of the Ring-finger next to the Little-finger.

When we lean or press on the internal Condyle of the Os humeri, the Numbness and Prickling we frequently feel, point out the Course of this Nerve. I have seen a Weakness

and

and Atrophy in the Parts which I mentioned this Nerve to be fent to, after a Wound in the

internal lower Part of the Arm.

7. RADIALIS accompanies the humeral Artery to the bending of the Elbow, serving the Flexors of the Cubit in its Way; then palfing through the pronator Radii teres Muscle, it gives Nerves to the Muscles on the Fore-part of the Fore-arm, and continues its Course near to the Radius, bestowing Branches on the circumjacent Muscles. Near the Wrist, it sometimes gives off a Nerve which is distributed to the Back of the Hand, and the convex Part of the Thumb and several of the Fingers, instead of the Branch of the Muscular. larger Part of this Nerve, passing behind the annular Ligament of the Wrist, gives Nerves to the short Muscles of the Thumb; and afterwards fends a Branch along each Side of the Sheath of the Tendons of the Flexors of the Thumb, Fore-finger, Mid-finger, and one Branch to the Side of the Ring-finger, next to the Middle one, to be loft on the concave Side of those Fingers.

Though the radial Nerve passes through the pronator Muscle, and the muscular Nerve seems to be still more unfavourably placed within the supinator brevis, yet the Action of these Muscles don't seem to have any Essect in hindering the Insluence of these Nerves, for the Fingers or Hand can be bended while Pronation is performing vigorously, and they can be extended

while Supination is exercised.

The Manner of the going off of these Nerves of the Fingers, both from the Ulnar and Ra-L 1 dial.

dial, is, that a fingle Branch is fent from the Trunk to the Side of the Thumb and Little. finger farthest from the other Fingers; and all the rest are supplied by a Trunk of a Nerve. which splits into two some Way before it comes as far as the End of the Metacarpus, to run along the Sides of different Fingers that are nearest to each other.

It might have been observed, that, in deferibing the posterior Branches of the ulnar and muscular Nerve, I did not mention the particular Fingers, to the convex Part of which they are distributed. My Reason for this Omission is, the Uncertainty of their Distribution; for though fometimes these posterior Branches go to the same Fingers, to the concave Part of which the anterior Branches of the Ulnar and Radial are fent, yet frequently they are distributed otherwise.

The Situation of these brachial Nerves in the Axilla, may let us fee, how a Weakness and Atrophy may be brought on the Arms by long continued Pressure of Crutches, or such other hard Substances on this Part; and the Course of them from the Neck to the Arm may teach us, how much better Effects Vesicatories, or stimulating nervous Medicines, would have, when applied to the Skin, covering the transverie Processes of the Vertebræ of the Neck, or at the Axilla, than when they are put between the Shoulders, or upon the spinal Processes, in Convulsions or Palsies of the superior Extremities, where a Stimulus is required.

The TWELVE DORSAL Nerves of each Side, as foon as they escape from between the Vertebræ, Vertebræ, send a Branch forward to join the Intercostal, by which a Communication is made among them all; and they soon likewise give Branches backwards to the Muscles that raise the Trunk of the Body, their principal Trunk being extended outwards to come at the Fursow in the lower Edge of each Rib, in which they run toward the anterior Part of the Thorax, between the internal and external intercostal Muscles, giving off Branches in their Course to the Muscles and Teguments of the Thorax.

The FIRST Dorfal, as was already observed, is particular in this, that it contributes to form the brachial Nerves; and that the two Branches of the Intercostal, which come down to the Thorax, form a considerable Ganglion with it.

The SIX lower dorfal Nerves give Branches to the Diaphragm and abdominal Muscles.

The TWELFTH joins with the first Lumbar, and bestows Nerves on the Musculus quadratus Lumborum and Iliacus internus.

May not the Communications of all these Nerves be one Reason, why the Parts they serve act so uniformly and conjunctly in Respiration, and conspire together in the convulsive Motions of Coughing, Sneezing, &c.?

The twitching Spaims that happen sometimes in different Parts of the Muscles of the Abdomen, by any Irritation on the Branches of the lower dorsal Nerves, are in Danger of occasioning a Mistake in Practice, by their Resemblance to the Colic, Nephritis, &c.

The Communications of these lower ones with the

the Intercostals, may serve to explain the violent Effort of the abdominal Muscles in a

Tenesimus and in Child-bearing.

As the Intercostal is larger in the Thorax than any where else, and seems to diminish gradually as it ascends and descends, there is Cause to suspect that this is the Trunk from which the superior and inserior Parts are sent as Branches.

The FIVE LUMBAR Nerves on each Side communicate with the Intercostal and with each other, and give Branches backwards

to the Loins.

The FIRST communicates with the last Dorsal, sends Branches to the abdominal Muscles, to the Psoas and Muscles, and to the Teguments and Muscles on the Fore-part of the Thigh; while its principal Branch joins with the other Nerves, to form the crural Nerve.

The SECOND LUMBAR Nerve passes through the Psoas Musicle, and is distributed nearly in the same Way as the former; as is

also the THIRD.

Branches of the fecond, third, and fourth, make up one Trunk, which runs along the Fore-part of the Pelvis; and passing in the Notch at the Fore-part of the great Hole common to the Os Pubis and Ischium, is spent on the Adductor Muscles, and on the Teguments on the Inside of the Thigh. This Nerve is called the OBTURATOR or POSTERIOR CRURAL NERVE.

By united Branches from the first, second, third, and fourth lumbar Nerves, a Nerve is formed that runs along the Psous Muscle, to escape escape with the external Iliac Vessels out of the Abdomen, below the tendinous Arcade of the external oblique Muscle. This Nerve, which is named the ANTERIOR CRURAL, is distributed principally to the Muscles and Teguments on the Fore-part of the Thigh. A Branch, however, of this Nerve runs down the Inside of the Leg to the upper Part of the Foot, keeping near to the Vena saphæna; in opening of which with a Lancet at the Ancle, the Nerve is sometimes hurr, and occasions sharp Pain at the Time of the Operation, and Numbness afterwards.

The Remainder of the fourth Lumbar and the fifth join in composing the largest Nerve of the Body; which is soon to be described.

Whoever attends to the Course of these lumbar Nerves, and of the spermatic Vessels and Nerves upon the *Psoas* Muscle with the oblique Passage of the *Ureter* over that Muscle, will not be surprised, that when a Stone is passing in this Canal, or even when it is inflamed, the Trunk of the Body cannot be raised erect, without great Pain; or that the Skin of the Thigh becomes less sensible, and the Thigh is drawn forward, and that the Testicle often swells and is drawn convulsively towards the Ring of the abdominal Muscles.

The SIX PAIR of the FALSE VERTE-BRÆ confift each of small posterior Branches sent to the Hips, and of large anterior Branch-

es.

The first, second, and third, after coming through the three upper Holes in the Fore-part of the Os sacrum, join together with the L 1 3 fourth

fourth and fifth of the Loins, to form the largest Nerve of the Body, which is well known by the Name of SCIATIC or ISCHIATIC Nerve: This, after fending large Nerves to the different Parts of the Pelvis, and to the external Parts of Generation and the Podex. as also to the Muscles of the Hips, passes behind the great Tuber of the Os Ischiem, and then over the Quadrigemini Muscles to run down near to the Bone of the Thigh at its Backpart, giving off Nerves to the neighbouring Muscles and Teguments. Some way above the Ham, where it has the Name of the Poplitæus Nerve, it fends off a large Branch that passes over the Fibula, and finking in among the Muscles on the anterior external Part of the Leg, runs down to the Foot, to be loft in the upper Part of the larger Toes, supplying the neighbouring Muscles and Teguments every where in its Paliage. The larger Branch of the Sciatic, after giving Branches to the Muscles and Teguments about the Ham and Knee, and fending a large cutaneous Nerve down the Calf of the Leg, to be lost at last on the Outlide of the Foot and upper Part of the lesser Toes, finks below the Gemelhis Muscle, and distributes Nerves to the Muscles on the Back of the Leg; among which it continues its Course, till passing behind the internal Malleolus, and in the internal Hollow of the Os calcis, it divides into the two Plantar Nerves: The internal of which is distributed to the Toes in the same Manner the radial Nerve of the Hand ferves the concave Side of the Thumb and Fingers; and the external Plantar

Plantar is divided and distributed to the Sole of the Foot and Toes, nearly as the ulnar Nerve is in the Palm of the Hand, and in the concave Part of the Fingers.

Several Branches of these Nerves, that serve the inferior Extremities, pierce thro

Mufcles.

By applying what was said of the Nerves in general to the particular Distribution of the Nerves of the inferior Extremities, we may see, how People with fractured Legs, especially where there are Splinters, should be subject to convulsive Startings of the fractured Member; why, upon tying the Blood-vessels in an Amputation of the Leg, the Patients should sometimes complain of violent Pain in their Toes; — why such Patients should also be troubled with Startings; — why, for a considerable Time after the Amputation of the diseased Limb, when the Suppuration is well advanced, they should complain of Pain in the Sore which occasioned the Amputation.

The FOURTH, which, with the two following, is much fmaller than the three inperior, foon is lost in the Vesica urinoria and In-

tellinusn rectum.

The FIFTH comes forward between the Extremity of the Os facrum and Coccygis, to be distributed principally to the Levatores Ani.

The SIXTH, which fome think to be only a Production of the Dura Mater, advances forward below the broad Shoulders of the first Bone of the Os Coccygis, and is lost in the Spincter Ani and Teguments covering it.

The Branches of the four last cervical Nerves,

and

and of the first Dorsal, which are bestowed on the Superior Extremities, and the two Crurals. with the Sciatic, which are distributed to the inferior Extremities, are much larger proportionally to the Parts they ferve, than the Nerves of the Trunk of the Body, and especially of the Viscera, are; and for a very good Reason, that in the most common necessary Actions of Life, a sufficient Quantity of Fluid, on which the Influence of Nerves feems to depend, may be supplied to the Muscles there, which are obliged to perform more frequent and violent Contractions than any other Parts do. The Size of the Nerves of the inferior Extremities feems larger proportionally than in the superior Extremities; the inferior Extremities having the Weight of the whole Body to fustain, and that frequently at a great Disadvantage. What the Effect is of the Nerves here being injured, we fee daily, when People happen, by fitting wrong, to compress the sciatic Nerve, they are incapable for some Time after to support themselves on the affected Extremity: And this is still more remarkable in the Sciatic or Hip-gout, in which the Member is not only weakened, but gradually shrivels and wastes.

OF THE TO A SALE

englishing as at the fe

THE COURT OF THE PARTY OF THE P

eyer M. Rios are deprived of a set water

Systole and DIASTOLE

in Rolling and a population

of animostif he find that is on

some of their Propositional Leve

Land Thomas and photom chimall

HEART,

And the Reciprocal Actions of its

Auricles and Ventricles.

THAT the Account I am to give of the alternate Motions of the Heart may be rightly understood, it will be necessary to premise the following Propositions, on which it depends.

1. All Muscles are furnished with Blood-velsels and Nerves, and their Action depends on the Instux of Blood, and of the nervous Fluid into their muscular Fibres; therefore, whenever

ever Muscles are deprived of a sufficient Quantity of either one or other of these Liquors. or of both, their Action is weakened or ceases. The Truth of this has been fully proved by the Experiments of compressing, tying and cutting the Nerves or Arteries of Muscles.

2. All Muscles are in a constant State of Action, as long as Blood and nervous Fluid are freely fupplied to them. This feems evident from the continued Contraction of the Sphineters of the Bladder and Anus, and of Muscles, whose Antagonists are cut asunder, or become

paralytic.

Though in these Propositions I have assumed a Fluid of the Nerves, for the Probability of which, fee the Arguments in the foregoing Treatise; yet to avoid a Dispute, it may be here remarked, that the following Account of the Heart's Motions will equally hold good, by supposing the Nerves to be solid Cords acting by Elasticity, Vibration, &c. if it is only granted, that their Action is necessary to the Contraction of Muscles, and that Compression is capable to hinder that Action; which Experiments plainly shew it is.

3. The Nerves of the Heart pass to it between the two Auricles or two Arteries, and

between the Auricles and Arteries.

4. The Coronary Arteries, which are the only ones that supply the Heart, rise from the Aorta immediately above the middle loofe Part of the semilunar Valves, in the same Height of the Aorta, as where the Angles of these Valves are fixed. geneal to has books to want all

and the continue there is the there where

5. The Edges of the semilunar Valves are duplicated with a muscular Corpuscle in the Middle, and muscular Cords running obliquely from that Corpuscle to the Insertion of the Valves into the Aorta:

The Truth of these three last Propositions

is evident from Dissections. A har , --

6. Before the Vessels in which the Circulation is performed can act, it is necessary to suppose them full of their Liquors; otherwise the whole vascular System could not be all at once put into Action, and the Motion of the Fluids sent out from the Heart could not be propagated to that in the returning Vessels; consequently the Circulation would be stopped almost as soon as

it this lame Time when we niged

If then both Auricles and Ventricles attempt, upon the first Communication of Motion, to contract, the Ventricles being stronger, will force their own Contraction, and hinder the Contraction of the Auricles, which must be, in the mean Time, much dilated by the Influx of Blood from the Veins, (Prop. 6.); and at this Time the Arteries are also distended by the Blood thrown out of the Ventricles; therefore the Cardiac Nerves lying between them, (Prop. 3.) are compressed, and prevented from exerting their Office. While this happens to the Nerves, the Blood rushing out of the left Ventricle into the Aorta, and thrusting the femilunar Valves outwards and upwards, diftends the great Artery confiderably, whereby the Valves are stretched, and their lunated Edges are brought to be streight and as high as their Angles; confequently these Edges, thus raised and

and pressed to the Sides of the Aorta, are made to cover the Orifices of the Arteria coronaria, which the Corpuscles will assist to do more adequately; and therefore, while they are in this Situation, no Blood can be sent to the Substance of the Heart. If then the Nerves do not exert their Ossice, and Access is denied to the Blood, this Muscle, the Heart must, (by Prop. 1.) be-

come paralytic or unactive.

The Auricles, which were attempting all this Time to contract themselves, will now, when the Resistance to them is removed, throw the Blood they contain into the Ventricles; and the Arteries that were violently distended while the Ventricles pushed the Blood into them, will at this same Time when the Ventricles cease to act, constrict themselves; and the Valvulæ semilunares will, by their Elasticity, affifted by the muscular Corpuscles and Cords, be depressed inwards. Since then, by the Constriction of the Auricles and Arteries, the Nerves of the Ventricles are freed from the Compression they suffered while the Auricles and Arteries were distended; and fince the Blood in the Aorta, pressed strongly by the contracting Sides of this Artery, and relifted by the Liquors which are to be thrown through all its Branches, is also pushed with great Force on the Orifice of the left Ventricle, thrusts back the femilunar Valves which stop its Entry into that Cavity, and impetuously rushes into the coronary Arteries: Since then, I fay, the Nerves of the Heart are again free from Compression, and the Blood is again fent to the Heart by its Arteries, its muscular Fibres must again contract.

tract, (by Prop. 2.). And thus, as long as these Causes continue to act and cease alternately, their Effects must be exerted in the same Manner; that is, as long as an Animal lives, the Heart must have an alternate State of Contraction, called its Systole, and of Relaxation, or Diastole; and the Auricles and Arteries must be dilated, while the Ventricles are contracting; and the Ventricles must be dilated, while the Auricles and Arteries are constricted; or, their Actions must always be reciprocal.

Mm

THE

And the result of the result o

- M. - .

tes Daylor, ou the wife of the

and the board of the second

of the solded for and mit

DESCRIPTION

OF THE HUMAN

Notes to Constant, and great Outstander of a constant of the c

LACTEAL SAC and DUCT.

THE Receptaculum Chyli of Pecquet, or Saccus Lacteus of Van Horne, is a membranous somewhat pyriform Bag, two thirds of an Inch long, one third of an Inch over in its largest Part when collapsed; situated on the first Vertebra of the Loins to the right of the Aorta, a little higher than the right emulgent Artery, behind the right inferior Muscle of the Diaphragm: It is formed by the Union of three Tubes, one from under the Aorta, the second from the Interstice of the Aorta and Cava, the third from under the Emulgents of the right Side. The Lacteal Sac, becoming gradually fmaller towards its upper Part, is contracted into a slender membranous Pipe, of about a MI m 2 Line

Line Diameter, which is generally named the THORACIC DUCT. This passes betwixt the muscular Appendices or inferior Muscles of the Diaphragm, on the right of, and somewhat behind the Aorta: then, being lodged in the cellular Substance behind the Pleura, it mounts between the Aorta and the Vena Azygos as far as the fifth Vertebra of the Thorax, where it is hid by the Azygos, as this Vein rifes forwards to join the descending or superior Cava; after which the Duct passes obliquely over to the left Side behind the OE fophagus, Aorta descendens, and great Curvature of the Aorta, until it reaches the left carotid Artery; behind which, on the left Side of the OE/ophagus, it runs to the Interstice of the first and fecond Vertebra of the Thorax, where it begins to separate from the carotid, stretching farther towards the left internal Jugular Vein by a circular Turn, whose convex Part is uppermost. At the Top of this Arch it splits into two for a Line and an half; the superior Branch receiving into it a large lymphatic Velfel from the cervical Glands. This Lymphatic appears, by blowing Air, and injecting Liquors into it, to have few Valves. When the two Branches are again united, the Duct continues its Course towards the internal Jugular Vein; behind which it descends, and, immediately at the lest Side of the Insertion of this Vein, enters the fuperior posterior Part of the left subclavian Vein, whose internal Membrane duplicated, forms a femilunar Valve that is convex externally, and covers two thirds of the Orifice of the Duct; immediately below this Orifice

Orifice, a cervical Vein from the Musculi sca-

The Coats of the Sac and Duct are thin transparent Membranes; from the Inside of which, in the Duct, small semilunar Valves are produced, most commonly in Pairs; which are so situated, as to allow the Passage of Liquors upwards, but oppose their Return in an opposite Course. The Number of these is generally ten or twelve.

This is the most simple and common Course, Situation, and Structure of the Receptaculum chyli and thoracic Duc; but having had Occasion to observe a Variety in these Parts, of different Subjects, I shall set down the most

remarkable of them.

The Sac is sometimes situated lower down than in the former Description; is not always of the same Dimensions; is not composed of the same Number of Ducts; and frequently appears to consist of several small Cells or Ducts,

instead of being one simple Cavity.

The Diameter of the Duct is various in most Bodies, and is seldom uniform in the same Subject; but frequently sudden Enlargements or Sacculi of it are observable.—The Divisions which Authors mention of this Duct are very uncertain. I have seen it divided into two, whereof one Branch climbed over the Forepart of the Aorta at the eighth Vertebra of the Thorax, and at the fifth slipped behind that Artery, to join the other Branch which continued in the ordinary Course.—The precise Vertebra, where it begins to turn to the left Side, is also uncertain.—Frequently it does

not split at its superior Arch; in which Case a large Sac is found near its Aperture into the Subclavian Vein. Generally it has but one Orifice; though I have feen two in one Body, and three in another: Nay, sometimes it divides into two, under the Curvature of the great Artery; one goes to the right, another to the left Subclavian Vein; and I have found this Duct discharging itself entirely into the right Subclavian. -- The lymphatic Vessel which enters its superior Arch, is often sent from the thyroid Gland.

Whether is not the Situation of the Receptaculum chyli fo much nearer the muscular Appendices of the Diaphragm in Men than in Brutes, defigned to supply the disadvantageous Course the Chyle must otherwise have in our

erect Posture?

עודי לי ביי ביי ביי ביי ביי אברי

Does not the Descent of the End of the Duct to the Subclavian Vein, and the opening of the lymphatic into the Top of the Arch, contribute to the ready Admission of the Chyle into. that Vein?

FINIS.

23 JY60

Errors of the Press.

PAge 2. Line antepenult, for cafily, read easily.-p. 5. l. 16. f. bluish, r. blueish. p. 21. 1. 23. f. their, r. these other. p. 37. 1. 7. for End, r. Ends.—p. 60. l. penult. f. cavarja, r. calvaria.—p. 89. l. ult. f. cia, r. ria.—p. 99. l. 26. f. Vertehra, r. Vertebra .-- p. 129. l. 13. f. of each Jaw, r. of each Side of each Jaw.-p. 131. l. 16. f. those, r. these.-p. 148. l. 29. f. Munducation, r. Manducation .- p. 161. l. 2. f. Bassinglossi, r. Bafioglossi .- p. 205. l. ult. f. Pictinis, r. Pectinis .-- p. 211. l. penult. f. Pectus cassum, r. Pectus, cassum. p. 225. l. ult. f. malum, granatum, r. malum granatum.-p. 237. 1. ult. f. Ayxuposions r. Ayxuposions ___ p. 241. l. penult. f. cana, r. canna. p. 247. l. 19. f. Longus Brevis, r. Longus, Brevis. p. 200. l. 20. f. Corpus, r. Carpus.—p. 388. l. 24. f. the other Branches, r. the Branches.

23 JY60